

Research Article

Analytical Techniques for a Rapid Determination of Nitrogen Content in Extra Virgin Olive Oil

Rizzo A and Telloli C*

Department of Nuclear Safety, Division of Security and Sustainability, Energy and Sustainable Economic Development (ENEA), Italian National Agency for New Technologies, Bologna, Italy

Abstract

A more attention to agri-food products was increase from the consumers not only because of their nutritional value but also for their origin (location) and authenticity, in order to practice more conscious consumption and to safeguard their health. Analytical chemistry played an important role in traceability for all products in the food market. Olive oil is an important agricultural product for the economy of the Mediterranean area, but it has undergone major alterations in quality in recent years. In Italy, some farmers worked in the olive oil sector have been subjected to legal investigations due to manipulations through mixture with olive oils from other countries. The aim of this study was to verify and develop a fast screening method to characterize specific parameters of olive oil, such as acidity and nitrogen content, and to highlight correlations with the pathogen *Xylella fastidiosa*.

Keywords: CN; Olive oil; Traceability; *Xylella fastidiosa*

*Corresponding author: Chiara Telloli, Department of Nuclear Safety, Division of Security and Sustainability, Energy and Sustainable Economic Development (ENEA), Italian National Agency for New Technologies, Martiri di Monte Sole 4, 40129, Bologna, Italy, Tel: + 39 0516098485; E-mail: chiara.telloli@enea.it

Citation: Rizzo A, Telloli C (2019) Analytical Techniques for a Rapid Determination of Nitrogen Content in Extra Virgin Olive Oil. J Agron Agri Sci 1: 005.

Received: December 18, 2018; **Accepted:** January 18, 2019; **Published:** February 01, 2019

Copyright: © 2019 Rizzo A and Telloli C. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Recently, food traceability is becoming an important factor to characterize the authenticity of agri-food products [1]. European Union established a legislation to guarantee the safety and quality of consumer products [2]. Olive Oil (OO) is one of the most representative products in Southern Europe and has nutritional and organoleptic qualities that enable it to have a high commercial value. It is important to note that European Union is the first producer of OO in the world (75.4%), followed by Tunisia (5.9%), Syria (4.8%), Turkey (4.6%), and Morocco (3.1%) [3]. In the European Union, Spain (52.4%), Italy (27.7%) and Greece (17.3%) represent 97% of OO producers [4].

The traceability of OO allows characterizing the identity, history and origin of the product in order to strengthen its food safety [5]. As a matter of that, Extra Virgin Olive Oil (EVOO) must follow specific characteristics as chemical, physical, biological or sensory nature to be included in the commercial market but it is difficult to verify the authenticity of OO due to the wide range of categories, cultivations, pedoclimatic conditions and working methods.

With the use of rapid analytical methods, it is possible to verify the presence of pollutants in the OO with the aim of eliminating the risk of fraud. A typical chemical-physical characteristic of the OO is the acidity, conventionally expressed as a percentage of free oleic acid. Acidity is useful to recognize extra virgin, virgin, lampante or refined olive oil. The Commission Implementing Regulation (EU) No 1348/2013 describes the guidelines for a correct identification of OO, and the Regulation (EC) No 1989/2003 and the International Olive Council (2015) describe the values and the analytical technique of titration to define the characteristics of OO to be introduced into the economic market [6-8].

In the years 2013-2015, in the main Italian regions producing EVOO, olive trees suffered a dangerous attack by *Xylella fastidiosa* pathogen, and a new legislation was adopted, regarding the “measures to prevent the introduction into and the spread within the Union of *Xylella fastidiosa*” [9]. In 2014, the bacterium, which grows in the xilema of the plant caused a reduction in the production of olives and consequently of EVOO [10-12]. The impact of this pathogen on the productivity of olive groves was devastating, despite that, the economic market did not suffer any losses, leaving the doubt that more or less legal methods were used to have the same quality of EVOO.

Recent studies shown that the presence of nitrogen from fertilizers could implement the metabolism of nitrogen of the plant [13,14]. Massive use of fertilizers could be used to restore fertile activity in olive trees during these years in the Italian region affected by the *Xylella fastidiosa*. For this reason, the presence of fertilizer could be a critical factor for the quality of OO and for the safety of the agro-food product, which is in contrast to the EU anti-fraud directives [15].

The object of this paper was to develop an analytical technique for determining traces of nitrogen in samples of EVOO from different Italian regions and several years (2013-2015). The analytical method was developed favoring quick analysis times, simple sample

treatment procedures, and limited use of reagents. The analytical technique considered is based on an elemental combustor, which allows complete oxidation of the matrix (eg., OO) by simultaneously separating species containing carbon and nitrogen.

Materials and Methods

Sampling

OO samples were collected in the period between 2013 and 2015 from small and medium Italian mills (Figure 1) with the exception of the samples from Albania (EVOO-006 and EVOO-010). The samples analyzed of Sunflower Oil (SFO), Corn Seeds (CO) and Olive Oil (OO) were obtained from virgin and refined oils from industrial certified processes, and EVOO from different mills declared as extra virgin olive by the producer, with the exception of the samples of Albania (EVOO-006 and EVOO-010), Perdifumo (SA) (EVOO-012) and Rossano (CS) (EVOO-016) that were produced by farmers for strictly family use with handmade techniques.




Figure 1: Map of Italy, where the origin of the samples analyzed is specified. On the top, map of the Mediterranean area with Albania region. Next to the names of origin of the olive oil samples, the production years are specified.

Table 1 shows the list of the OO samples collected. After collection, all samples were transported to ENEA (Italian National Agency for New Technologies, Energy and Sustainable Economic Development) laboratory in Bologna (Italy) and maintained at a constant conditions of temperature (laboratory room temperature: 22°C) until analyses were performed.

Titration

Titration allows calculating the degree of acidity expressed in percentage of free oleic acid, as described in the protocol of the International Olive Council (2015). The analytical method was based on a dilution of 0.1M sodium hydroxide solution (NaOH) starting from 1M ± 0.001 in a flask of 100.0 ml (± 0.1 ml). Diethyl ether / ethanol solution (95% v/v) was mixed in equal volume (100 ml) to which 0.3 ml of phenolphthalein was added. The procedure involved the use of an OO matrix of about 10 g. To reduce waste and environmental impact, a small amount of OO was analyzed reducing the volume of alcohol/ether mixture, since it was noticed that the result did not change. The matrixes were determined by weighing in a 250 ml flask the OO using an electronic balance KERN ABT (ARW Balance, Vicenza, Italy). The mixture of alcohol and ether (50 ml) was combined

and titration with 0.1M NaOH was carried out. When the solution has been reached, the mixture seemed to have an orange color due to the phenolphthalein presence. The titrated volume is recorded and the degree of acidity is calculated as the percentage of the weight of the free oleic acid (Equation 1):

$$\frac{V.M.MW_{\text{freeoleic acid}} \cdot 100}{1000.W} = \frac{V.M.MW_{\text{freeoleic acid}}}{W.10} \text{ @ equation 1}$$

V = Sample Volume

M = Molarity of the Titrant (NaOH)

MW_{free oleic acid} = Molecular Weight = 282.46 g/mol

W = Sample Weight.

Sample	Location	Year
EVOO-001	Italy Toscana: Pistoia (PT)	2013
EVOO-002	Italy Puglia: Barletta (BAT)	2013
EVOO-003	Italy Puglia: Manfredonia (FG)	2013
EVOO-004	Italy Puglia	2013
EVOO-005	COOP FIORE	2013
EVOO-006	Albania	2014
EVOO-007	Italy Toscana: Prato (PO)	2014
EVOO-008	Italy Puglia: Molfetta (BA)	2014
EVOO-009	COOP FIORE	2014
EVOO-010	Albania	2015
EVOO-011	Italy Marche: Colli del Tronto (AP)	2015
EVOO-012	Italy Campania: Perdifumo (SA)	2015
EVOO-013	Italy Campania: Pimonte (NA)	2015
EVOO-014	Italy Calabria: Catanzaro (CZ)	2015
EVOO-015	Italy Castrovillari (CS)	2015
EVOO-016	Italy Rossano (CS)	2015
EVOO-017	Italy Sardegna: Ales (OR)	2015
OO-001	Italy Toscana: Scandicci (FI)*	/
CO-001	Italy Emilia-Romagna: Modena (MO)*	/
SFO-001	COOP*	/

Table 1: List of the samples collected and analyzed (*industrial products, indicated the packaging area).

CN Vario Max Cube

Elemental analysis was carried out by an elemental combustor (Vario Max Cube, Elementar GmbH, Frankfurt and Germany), based on the Dumas combustion reaction, in order to determine the content of carbon and nitrogen and its ratio. The samples were prepared by weighing 250 ml of OO in stainless steel crucibles using the electronic balance KERN ABT.

The automatic sampler inserts the crucible into the combustion zone, where oxygen is injected directly on the sample. A second post combustion reactor converts possible byproducts of the combustion to carbon dioxide. Excess oxygen removal, conversion of nitrogen oxides to N₂ and removal of Sulphur is achieved in the tungsten reduction system. The analytical gases N₂ and CO₂ are completely separated by the purge and trap separation system. The gases are sent to the high performance Thermal Conductivity Detector (TCD) [16].

At the beginning of the analysis, standard samples (Aspartic acid) were analyzed for the calibration of the instrument.

Results

Table 2 shows the results of titration analysis to compare the acidity obtained from the averages of 3 repetitions, on that expressed on the Regulation (EC) No 1989/2003. EVOO from 2014 have high percentage of free oleic acid, respect the previous (2013) and next (2015), except for the red spots representing samples EVOO-006, EVOO-010, EVOO-012 and EVOO-016 produced by farmers for strictly family use. Samples EVOO-006, EVOO-010, EVOO-012 and EVOO-016 show the highest acidity, especially in the year 2015 (> 2%). This value not classify these types of oil as EVOO respect the Regulation (EC) No 1989/2003, but we remember that EVOO from Albania and Perdifumo were produced by farmers for strictly family use with handmade techniques. For this reason, we exclude the presence of pollutants or risk of fraud.

Table 2 also shows the average concentrations of carbon and nitrogen analyzed by CN Vario Max Cube. Carbon concentration is expressed in percentage, instead of nitrogen is expressed in Parts Per Million (PPM) in order to observe quantifiable nitrogen values. The results underline that carbon value is constant in all the samples, whereas nitrogen value is variable.

Figure 2 shows that nitrogen concentration is higher in the samples from 2013-2014, period characterized by the *Xylella fastidiosa* pathogen. In red, the Albania and Perdifumo samples show nitrogen concentration comparable to each other but lower respect to the Italian samples. SFO and CO show a similar nitrogen concentration of the EVOOs, whereas OO shows higher nitrogen concentration than EVOOs.




Figure 2: Nitrogen concentration (expressed in ppm) in the samples analyzed by CN Vario Max Cube.

Discussion

Titration shows that two EVOO samples, obtained with handmade methods, have a very high quantity of acidity; this anomaly could be related to the type of extraction. CN analysis proved to be a fast and accurate technique to characterize nitrogen and carbon concentration in EVOO samples: an analysis in 7 or 10 minutes. The high nitrogen concentration analyzed by CN could be related to the massive use of nitrogen fertilizer on olive trees.

EVOO samples produced in 2013 and 2014 years show higher nitrogen concentration compared to 2015. This could be related to the use of nitrogen-based fertilizers, in order to increase the yield of the olive oil product, strongly down because of the infestation due to *Xylella fastidiosa* pathogen.

	Acidity (% free oleic acid)	C (%)	C (dev std)	N (ppm)	N (dev std)
EVOO-001	0.28	77.38	0.08	405.5	32.44
EVOO-002	0.65	77.16	0.12	454.3	36.35
EVOO-003	0.54	77.20	0.08	568.1	45.45
EVOO-004	0.31	77.43	0.15	354.1	28.33
EVOO-005	0.42	77.32	0.18	396.0	31.68
EVOO-006	1.20	77.36	0.12	260.6	20.85
EVOO-007	1.30	77.05	0.08	478.2	38.26
EVOO-008	1.00	77.10	0.11	360.3	28.82
EVOO-009	0.64	77.33	0.14	366.7	29.34
EVOO-010	7.20	76.92	0.23	260.5	19.80
EVOO-011	0.34	77.06	0.14	120.0	24.56
EVOO-012	2.70	77.07	0.33	162.4	52.29
EVOO-013	0.92	77.01	0.14	101.2	8.10
EVOO-014	0.48	77.08	0.08	100.8	8.06
EVOO-015	0.45	77.09	0.02	136.7	15.28
EVOO-016	3.10	77.51	0.41	133.3	10.67
EVOO-017	0.56	77.05	0.18	108.1	1.13
OO-001	0.17	77.14	0.11	256.9	90.69
CO-001	0.17	77.28	0.08	156.8	21.31
SFO-001	0.22	77.22	0.12	126.7	21.58

Table 2: Averages of 3 repetitions of acidity analyzed by titration and carbon (expressed in %) and nitrogen concentration (expressed in ppm) analyzed by CN.

In 2013, in fact, the pathogen *Xylella fastidiosa* caused severe damage to all the olive trees in the Mediterranean area. In Puglia (South of Italy), *Xylella fastidiosa* infected 74,000 acres causing a major decrease in OO production during the years 2014 and 2015 [17]. This may cause a massive use of fertilizers by farmers. In accordance with this hypothesis, the Puglia Region has proposed a planning spray with insecticides to reduce the *Xylella fastidiosa* pathogen. Between the infestation and the effect of some frosts, the production of olive oil in the areas affected by the pathogen could be reduced by 60% [18]. On the other side indicated that an excessive use of nitrogen fertilizers on olive trees leads to a slow deterioration of sensory qualities [19].

Conclusion

The aim of this study was to verify and develop a fast screening method to characterize specific parameters of olive oil, such as acidity and nitrogen content, and to highlight correlations with infestations and / or parasites. The study showed that the volumetric analysis for the determination of the percentage of free oleic acid by titration is a useful method for a quick selection of the different oil samples: olive oil or extra virgin olive oil. It was also highlighted that two samples of olive oil produced by farmers for strictly family use with handmade techniques, show a higher quantity of acidity; this anomaly could be related to the extraction technique that could be the subject of future studies also through a self-production in the laboratory of small quantities of olive oil to verify the influence of the type of process on the final acidity.

Elemental analysis by CN Vario Max Cube was useful to determine the nitrogen content in samples of extra virgin olive oil; its applicability to this matrix and the information obtainable for the purposes of characterization of the oils has been studied. The analysis carried out on samples of extra-virgin olive oil using the elementary combustor proved to be a fast and precise technique to characterize the nitrogen and carbon content in olive oil. The innovation made by the instrument allows the use of samples without pre-treatments reducing analysis times and costs. The concentration of nitrogen, expressed in Parts Per Million (PPM), was higher in extra virgin olive oils produced in the years 2013 and 2014, compared to those of 2015. This could be due to the use of nitrogenous fertilizers in order to increase the yield of the product due to the infestation of the *Xylella fastidiosa* pathogen. It is interesting to note that the two highest values of N are related to two samples of 2013, coming from the same region Puglia, which was strongly affected by the infestation of the pathogen. The use of a new generation elementary combustor for olive oil screening can be useful for identifying trends and / or anomalies in the elemental composition of the matrix.

References

1. Pegasus (2002) General Principles and Requirements of Food Law - Regulation (EC) No 178/2002. Pegasus, London, UK.

2. Directive 2001/95/EC of the European Parliament and of the Council of 3 December 2001 on general product safety.

3. International Olive Oil Council (2016a) World Olive Oil Production. International Olive Oil Council, Madrid, Spain.

4. International Olive Oil Council (2016b) EU Olive Oil Production. International Olive Oil Council, Madrid, Spain.

5. Commission Implementing Regulation (EU) No 29 (2012) Marketing standards for olive oil.

6. Commission Implementing Regulation (EU) No 1348 (2013) Regulation (EEC) No 2568/91 on the characteristics of olive oil and olive-residue oil and on the relevant methods of analysis.

7. Regulation (EC) No 1989 (2003) Regulation (EEC) No 2568/91 on the characteristics of olive oil and olive-pomace oil and on the relevant methods of analysis.

8. International Olive Council (2015) Determination of free fatty acids, cold method. International Olive Council, Madrid, Spain.

9. Commission Implementing Regulation (EU) 789 (2015) Measures to prevent the introduction into and the spread within the Union of *Xylella fastidiosa*.

10. Fernández-Escobar R, Beltrán G, Sánchez-Zamora MA, García-Novelo J, Aguilera MP, et al. (2006) Olive Oil Quality Decreases with Nitrogen Over-fertilization. Hort Science 41: 215-219.

11. Burbank LP, Ortega BC (2018) Novel amplification targets for rapid detection and differentiation of *Xylella fastidiosa* subspecies fastidiosa and multiplex in plant and insect tissues. J Microbiol Methods 155: 8-18.

12. Monteiro MP, Clerici JH, Sahoo PK, Cesar CL, de Souza AA, et al. (2017) Stiffness signatures along early stages of *Xylella fastidiosa* biofilm formation. Colloids Surf B Biointerfaces 159: 174-182.

13. Lominadze S, Nakashidze N (2016) The influence of nitrogen fertilizers on nitrate accumulation in leaves of orange Washington Navel. Annals of Agrarian Science 14: 233-236.

14. Li Y, Sun Y, Liao S, Zou G, Zhao T, et al. (2017) Effects of two slow-release nitrogen fertilizers and irrigation on yield, quality, and water-fertilizer productivity of greenhouse tomato. Agr Water Manage 186: 139-146.

15. De Lange E (2013) Report on the food crisis, fraud in the food chain and the control thereof (2013/2091(INI)). Committee in the Environment, Public Health and Food Safety, A7-0434/2013.

16. <http://www.elementar.de/en/products/organic-elemental-analysis/vario-max-cube.html>

17. Bucci EM (2018) *Xylella fastidiosa*, a new plant pathogen that threatens global farming: Ecology, molecular biology, search for remedies. Biochem Biophys Res Commun 502: 173-182.

18. Giliberto J (2018) *Xylella*: il batterio minaccia la Terra di Bari, la Puglia ricorre agli insetticidi. Corriere della Sera.

19. Vossen PM (2007) Organic Olive Production Manual. UCANR Publications, California, USA.



Journal of Anesthesia & Clinical Care
Journal of Addiction & Addictive Disorders
Advances in Microbiology Research
Advances in Industrial Biotechnology
Journal of Agronomy & Agricultural Science
Journal of AIDS Clinical Research & STDs
Journal of Alcoholism, Drug Abuse & Substance Dependence
Journal of Allergy Disorders & Therapy
Journal of Alternative, Complementary & Integrative Medicine
Journal of Alzheimer's & Neurodegenerative Diseases
Journal of Angiology & Vascular Surgery
Journal of Animal Research & Veterinary Science
Archives of Zoological Studies
Archives of Urology
Journal of Atmospheric & Earth-Sciences
Journal of Aquaculture & Fisheries
Journal of Biotech Research & Biochemistry
Journal of Brain & Neuroscience Research
Journal of Cancer Biology & Treatment
Journal of Cardiology: Study & Research
Journal of Cell Biology & Cell Metabolism
Journal of Clinical Dermatology & Therapy
Journal of Clinical Immunology & Immunotherapy
Journal of Clinical Studies & Medical Case Reports
Journal of Community Medicine & Public Health Care
Current Trends: Medical & Biological Engineering
Journal of Cytology & Tissue Biology
Journal of Dentistry: Oral Health & Cosmesis
Journal of Diabetes & Metabolic Disorders
Journal of Dairy Research & Technology
Journal of Emergency Medicine Trauma & Surgical Care
Journal of Environmental Science: Current Research
Journal of Food Science & Nutrition
Journal of Forensic, Legal & Investigative Sciences
Journal of Gastroenterology & Hepatology Research
Journal of Gerontology & Geriatric Medicine
Journal of Genetics & Genomic Sciences
Journal of Hematology, Blood Transfusion & Disorders
Journal of Human Endocrinology
Journal of Hospice & Palliative Medical Care
Journal of Internal Medicine & Primary Healthcare
Journal of Infectious & Non Infectious Diseases
Journal of Light & Laser: Current Trends
Journal of Modern Chemical Sciences
Journal of Medicine: Study & Research
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology
Journal of Neonatology & Clinical Pediatrics
Journal of Nephrology & Renal Therapy
Journal of Non Invasive Vascular Investigation
Journal of Nuclear Medicine, Radiology & Radiation Therapy
Journal of Obesity & Weight Loss
Journal of Orthopedic Research & Physiotherapy
Journal of Otolaryngology, Head & Neck Surgery
Journal of Protein Research & Bioinformatics
Journal of Pathology Clinical & Medical Research
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance
Journal of Physical Medicine, Rehabilitation & Disabilities
Journal of Plant Science: Current Research
Journal of Psychiatry, Depression & Anxiety
Journal of Pulmonary Medicine & Respiratory Research
Journal of Practical & Professional Nursing
Journal of Reproductive Medicine, Gynaecology & Obstetrics
Journal of Stem Cells Research, Development & Therapy
Journal of Surgery: Current Trends & Innovations
Journal of Toxicology: Current Research
Journal of Translational Science and Research
Trends in Anatomy & Physiology
Journal of Vaccines Research & Vaccination
Journal of Virology & Antivirals
Archives of Surgery and Surgical Education
Sports Medicine and Injury Care Journal
International Journal of Case Reports and Therapeutic Studies

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>