

Review Article

Common Physiological Disorders of White/Irish Potato (*Solanum tuberosum*) Tubers Produced in Swaziland: A Review

Kwanele A Nxumalo^{1*}, Michael T Masarirambi¹, Mzwandile Mabuza², Tavagwisa Muziri³ and Tariro Masarirambi⁴

¹Horticulture Department, University of Swaziland, South Africa

²Crop Production Department, University of Swaziland, Swaziland, South Africa

³Horticulture Department, Midlands State University, Gweru, Zimbabwe

⁴LA26 UNISWA Senior Staff Housing, University of Swaziland, South Africa

Abstract

Physiological disorders of white/Irish potato tubers are abnormalities of the tubers which are not caused by infectious diseases/insects, or animals. Potato tuber abnormalities occur as a result of environmental stress, nutritional deficiencies or excess on the plant. Information was obtained through informal surveys in the four agro-ecological zones of Swaziland, review of existing literature and interviews of key farmers and producers of the potato crop. Physiological disorders of potato tubers encountered in this study included black heart, brown core, cold damage, greening, growth cracks, loose skin, enlarged lenticells, malformation, sprouting and vascular browning. Ways to alleviate various potato tuber disorders are suggested.

Introduction

The potato (*Solanum tuberosum*) is widely grown in the world and contributes significantly not only to food security but to nutritional security [1-3]. It is a vegetable crop characterised by a shallow and superficial root system and has been shown to be a heavier nutrient feeder taking up nutrition until maturity [4].

The potato also known as the white or 'Irish' potato is a member

*Corresponding author: Kwanele A Nxumalo, Horticulture Department, University of Swaziland, PO Luyengo, M205, Swaziland, South Africa, Tel: +268 7612 8026; E-mail: aknxumalo@uniswa.sz

Citation: Nxumalo KA, Masarirambi MT, Mabuza M, Muziri T, Masarirambi T (2017) Common Physiological Disorders of White/Irish Potato (*Solanum tuberosum*) Tubers Produced in Swaziland: A Review. J Agron Agri Sci 1: 001.

Received: November 19, 2017; Accepted: December 22, 2017; Published: January 09, 2017

of the solanaceae family thought to have originated in South America in the cool highlands of the Andes, Spanish explorers are believed to have introduced it to Europe and it later spread throughout the world and to the African continent [1,5,6]. In its habitat of origin, it is a cool season perennial. It is susceptible to freezing temperature; optimum temperatures for maximum growth are 10 to 15°C. The plant develops from rhizomes - seed pieces, it is characterized by compound leaves with opposite leaflets. When the plant flower, apical dominance is eliminated and the plant forms lateral branches [5].

The plant grows up to 60 to 120cm in height. The plant is a vine or haulm. Flower colour is cultivar dependent; it can be purple, white or yellow. The potato plant produces fruits similar to tomato and contains seeds which are mostly used for breeding purposes. The plant exhibits a high degree of heterozygosity because it is a polyploidy [7]. The underground portion of the stem has buds which give rise to lateral slender underground stems called stolons. It is the stolons which enlarge to form tubers. Tubers form at the time of flowering and they have buds called 'eyes' from where sprouting occurs to give rise to the vegetative plant [1,5,6,8].

In Southern Africa where Swaziland is located potatoes can successfully grow in the highlands or Highveld all year round, elsewhere like the Lowveld during the cool season or winter months [8,9]. Small tubers called 'sets' are used for propagation where planting can be on ridges or flatland at a depth of 15 to 30cm at a spacing of 90x30cm or 50x30cm depending on cultivar. The most suitable soils to grow potatoes are sandy loams which are deep, fertile, friable and well drained. The ideal pH is 4.8 to 6.5. Tuberization is promoted by Short Days (SD) and cool temperatures. Compact and poorly drained soils result in production of deformed tubers which is a physiological disorder. High temperatures inhibit tuberisation and it has been reported that tubers are not formed at 30°C and above [1,5,6,8].

Swaziland although still a developing country is classified under the middle income group, however it is still faced with poverty problems and struggling with the HIV and AIDS pandemic [10]. There are high hopes of the 2022 vision by that time the country would have leapfrogged to a near fully developed country. The backbone of the country's economy is largely dominated by agriculture whose contribution to the Gross Domestic Product (GDP) is about 11% [11]. Potato is a horticultural crop, (horticulture is a subsector of the agricultural industry) which is a popular vegetable in Swaziland. During production potato tubers are affected by a number of diseases and physiological disorders which affect produce quality. Most of the physiological disorders are often caused by environmental stress resulting from unpredictable and uncontrollable weather conditions and or mineral deficiencies usually micronutrients [12,13].

Physiological or non-pathogenic disorders in general

Physiological disorders can be divided into groups: genetic predisposition, nutrient imbalance and watering disorders [8,13,14]. Both Genetic (G) and Environmental (E) interactions are involved and this complex interplay of (GXE) factors are poorly understood for most disorders and in some cases contradictory results have been reported

apart from numerous names for a particular disorder in many cases [14]. Physiological disorders of vegetables are mainly caused by changing environmental conditions such as moisture, temperature, unbalanced soil nutrients, excess or inadequacy of certain soil minerals, poor drainage or extremes of soil pH [12,13,15-17]. Several ways can be employed to alleviate the various physiological disorders affecting potato tubers produced in the Kingdom of Swaziland.

Consumers of pre-packed and loose potatoes increasingly desire potatoes of good quality and skin finish. These should have a bloom, which is good and shiny skin with no blemishes. Attention has been focussed on postharvest skin diseases with little attention on physiological disorders. Black heart, brown centre and hollow heart are the most commonly reviewed disorders of potato. Sprouting, netting and loose skin are some of the physiological disorders affecting potato skin finish thereby affecting marketing for either seed or for home consumption.

The objective of this review was to identify common physiological disorders of potato produced under Swaziland environmental conditions in the four agro-ecological zones and suggest ways of alleviating these adverse conditions by situational/environmental manipulation and use of locally available resources.

Methodology

The study was a qualitative research. Information was sought through desk review of existing literature and informal surveys in the four agro-ecological zones of the country were carried out. Brief interviews of key farmers in potato producing areas were conducted. Samples of potato crops found on sale in the markets were observed, disorders were identified and described.

Agro-ecological zones of Swaziland

The country has four geographical zones with distinct topography, geology, soils, vegetation and climatic patterns (Figure 1). In the west is the Highveld, which is mountainous and has a vegetation of mainly commercial forests with the bulk of the land being used for subsistence farming [11,18]. It experiences a temperature range of 4.5 to 33°C [19]. It has rivers, waterfalls and gorges with some protected and natural areas including Malolotsha, Hawane and Phophonyane [20].

The Middleveld is characterized with temperatures ranging from 2.5 to 37.2°C [19]. This region has fertile valleys which favour intensive farming. It has the most diversely cultivated and heavily populated area in the country [11]. Protected nature reserve areas include Mantenga and Milwane [20]. Further east, there is the Lowveld with the largest area coverage of 40% of the country and is drought prone [19]. There is the Western Lowveld which is underlain by sandstone/claystone and the Eastern Lowveld which is underlain by basalt [19]. It has a vegetation of shrubs, and mean temperature ranges from 2.6 to 41.8°C with the bulk of commercial farms growing crops under irrigation, including the three sugar estates in the country and citrus fruit plantations. The nature reserves in the area are: Mlawula, Hlane, Shewula, Mbuluzi, Simunye and Nisela game reserves [11,18]. The fourth region is the escarpment called Lubombo plateau with an altitude of 600m above sea level and climatic conditions similar to the Middleveld [19]. Given the mountainous topography of the Lubombo plateau, only one eighth of the land is arable and the rest is suitable for animal grazing [19].




Figure 1: Ecological zones of Swaziland.

Common potato tuber physiological disorders encountered in Swaziland's four agro-ecological zones are reviewed in this paper.

Findings and Discussion

Data on findings of numerous physiological disorders of potato tubers were not quantified but are presented in table 1.

Interviews with farmers on potato physiological disorders

Different responses were obtained from different farmers from the 4 different agro-ecological zones of Swaziland, mostly in places where there is high potato production. Most of these physiological disorders are common to the potato production areas, but some are found in specific areas. Black-heart physiological disorder is common to all farmers' produce due to the fact that most of them do not have proper storage facilities after harvesting their produce. However, farmers in the Highveld not only experience this physiological disorder due to poor storage facilities, but also due to excessive rains received in the region, yet most of the soils drain poorly, thus promoting black-heart physiological disorder.

Brown core and hollow heart physiological disorders are very common in the Highveld of Swaziland, especially in places like Motshane, Nkhamba, Sgangani, Nkoyoyo and Mnyokane that are very prone to frost. Growth cracks are more experienced from potatoes grown in the Lower Middleveld and Vuvulane in the Lowveld where there is potato production in the cool season. In 2015 during the El Nino drought, even farmers in the Highveld and the upper Middleveld also experienced a lot of this physiological disorder on their produce due to excessive heat received and shortage of irrigation water. Enlarged lenticels are most seen from potato produced in the Highveld and Middleveld, especially in those places that have waterlogged soils. These places in include: EZulwini, EBuka, Mnyokane, Motshane to name a few. Farmers in the Highveld that plant potatoes in late March just before the winter season begin in May experience enlarged frost damage to the plants. Most of them have shifted to planting the potatoes in February so that when frost comes, the plants will be able to withstand it.

Name of physiological disorder	Plant part affected	Symptom	Cause	Management
Black heart	Tubers- internal	<ul style="list-style-type: none"> The inside of the tubers are greyish black to black and the affected tissue is normally well defined The affected tissue is hard and leathery 	<ul style="list-style-type: none"> Develops as a result of oxygen shortage or an oversupply of CO₂ which causes the tissue to die, especially if the oxygen shortage is accompanied by high temperatures Black heart can also occur in the potato field if high temperatures follow excessive rain which causes an oxygen shortage in the soil When harvested tubers are left on the land for a long time When seed planted by hand lie in furrows for a long time before being covered by soil, especially when it is very hot 	<p>After harvesting do not leave tubers in the sun on trailers under tarpaulins</p> <ul style="list-style-type: none"> Ensure sufficient ventilation during storage and transportation Avoid fields with poor drainage if rain is received in the after season Make sure that lifted tubers are picked up as soon as possible Cover seed tubers in furrows as soon as possible after planting
Brown core and hollow heart	Tubers-internal	The cavities can appear lengthwise or diagonal in the tuber and also have irregular shapes.	Brown core develops when tubers are very small and temperature is low (<15°C), especially during tuber initiation up to the tuber reaching 50 g. Cells die, turn brown and can easily tear apart.	<p>Select cultivars that are less prone to the disorder.</p> <ul style="list-style-type: none"> Avoid over-irrigation Avoid a low plant population If brown centre occurs regularly in early plantings, consider planting later when temperatures are slightly higher Endeavour to maintain uniform conditions throughout the season by managing fertilisation and irrigation
Cracking/ Growth cracks	Tubers-external	<ul style="list-style-type: none"> Growth cracks vary in depth and length, but normally occur in the length of the tuber. 	<p>Growth cracks occur during varying soil moisture conditions;</p> <ul style="list-style-type: none"> Tubers crack when a dry period is followed by heavy rains or over-irrigation. Moisture uptake causes a quick increase in tuber moisture and growth and consequently an increase in tuber size Uneven plant population, over fertilisation with nitrogen and nutrient imbalances contribute to the occurrence of growth crack Application of large amounts of nitrogen at one time after tuber formation 	<ul style="list-style-type: none"> Avoid cultivars that are prone to crack if growth cracks regularly occur and cause loss in income Where possible ensure uniform growing conditions, even plant population, good irrigation scheduling and good fertilisation practices
Enlarged lenticels	Tubers-external	<ul style="list-style-type: none"> Raised, white callus-like tissue on the surface of the tubers The size may vary from inconspicuous to large 	Enlarged lenticels are formed under conditions of oxygen shortage and excessive moisture in the soil surrounding the tubers, or in storage as the cells beneath the lenticels expand and the cell mass breaks through the corky layer.	<ul style="list-style-type: none"> Avoid fields that are prone to water logging Reduce irrigation two weeks prior to harvesting by irrigating when the plant available water is 40-50%
Frost damage	Tubers-internal	Exposure for short periods to temperatures around 2°C - 0°C can result in grey or reddish tissue. The tissue can also turn dark grey or black.	Exposure of tubers to temperatures that vary from just above freezing point (2°C) or below freezing-point.	<p>Do not leave harvested tubers on the land over night, especially if low temperatures are expected.</p> <ul style="list-style-type: none"> Avoid cultivars that tend to bear shallow in regions with low temperatures at the end of the growing season Do not store seed potatoes on the farm if the correct conditions are not available. Isolate stacks by covering it with grass and/or hessian bags, but preferably store it in a storeroom with temperature control and ventilation If seed potatoes must be transported in winter, the consignment must be properly covered and it must be done during the day Ensure that mechanisms are in place for proper temperature control during cold storage Harvest plantings with tubers close to the soil surface before the first cold front is expected
Greening	Tubers-external	<ul style="list-style-type: none"> Greening of the tissue immediately under the skin The intensity may differ and may also be accompanied by sunburn 	Greening occurs when tubers are exposed to light (sun or artificial).	<ul style="list-style-type: none"> Avoid cultivars that are prone to bear shallowly Avoid high nitrogen application, especially with cultivars that are prone to form long stolons Plant in well prepared soils If possible, irrigate to prevent cracks occurring during dry weather conditions If tubers are exposed after foliage die-off, it can be ridged to cover it with soil. Do not expose tubers to light for long periods after harvesting Use packaging materials that provide sufficient protection against light transmission if potatoes are exposed to light for more than a few days. White paper transmits more light than brown paper

Internal brown fleck	Tubers-internal	Irregular brown flecks that primarily occur in the vascular bundle ring.	The primary cause of brown fleck is localised shortages in Calcium (Ca). Environmental factors that lead to brown fleck include air and soil temperature, soil types and soil moisture.	<p>Ensure that sufficient Ca is available for uptake where and when tubers are formed. Where granular fertilizer is used, the Ca must be available in the whole tuber zone.</p> <ul style="list-style-type: none"> Opting for planting times to avoid high temperatures during tuber formation and harvesting, reduces the risk associated with brown fleck Avoid stress conditions at the beginning of the growing season through proper fertilisation and irrigation practices
Loose skin	Tubers-external	<ul style="list-style-type: none"> The skin is totally or partially removed from the tubers leading to the underlying tuber tissue being exposed. 	<ul style="list-style-type: none"> Excessive application of nitrogen (N) late in the season and wet soil may delay ripening and promote loose skin. 	<ul style="list-style-type: none"> Top-growth must be completely dead 14 to 21 days prior to harvesting to promote skin setting Ensure that the harvester is properly set to prevent tuber damage
Malformation	Tubers-external	Tubers form secondary tubers, some on short stolons and others directly on an eyelet.	<ul style="list-style-type: none"> Malformed tubers form when an interruption in growth is experienced during the growing season as a result of high temperature, often accompanied by moisture stress and nutrient shortage 	<ul style="list-style-type: none"> Avoid production in periods when heat waves generally occur Avoid cultivars that are not adaptable to high temperatures, or plant it during cooler times of the year Ensure uniform growing conditions
Netting	Tubers-external	<ul style="list-style-type: none"> Shallow fissures that give the skin a distinctive net like appearance. 	<p>It is related to climatic conditions and physiological factors in the skin of young tubers.</p> <ul style="list-style-type: none"> Moisture stress leads to an increased intensity of the disorder 	Where possible tubers should not be exposed to varying soil moisture conditions.
Sprouting	Tubers-external	Potatoes start to sprout when it is harvested, especially when tubers have been left in the soil for long.	<p>Sprouting is promoted by high temperature before harvesting.</p> <p>Sprouting on the land is normally an indication that the cultivar is not adapted to the cultivation practices of a specific area.</p>	Evaluate new cultivars for at least three years in the production area and follow normal cultivation practices to identify cultivars that are not totally adapted.
Thumb nail cracks	Tubers-external	<ul style="list-style-type: none"> Mechanical impact manifests often as thumb nail cracks on tubers. 	<ul style="list-style-type: none"> The cracks can develop after a slight impact or injury and initially only occur in the skin (periderm) of the tuber without bruising to the underlying tissue. 	<ul style="list-style-type: none"> To prevent the occurrence of thumb nail cracks, tubers should not be harvested or handled when it is cold. The temperature of tuber tissue should be >9°C The crop should be fertilised optimally to ensure that sufficient Ca and Mg are present in the cell-walls
Vascular browning	Tubers-internal	<ul style="list-style-type: none"> Vascular browning of variable intensity develops at the stolon end, but in serious cases it can extend throughout the whole vascular ring Vascular browning appears as speckled, light brown to brown and even dark brown stripes 	<p>Physiological vascular browning is generally associated with a combination of low soil moisture and sudden foliage die-off as a result of chemicals, frost or mechanical removal</p> <ul style="list-style-type: none"> Vascular browning can also be caused through infection by tuber-borne pathogens such as <i>Verticillium</i> and <i>Fusarium</i> and leaf roll virus 	Irrigation prior to foliage die-off decreases the incidence of vascular browning, irrespective of the method of foliage die-off.

Table 1: Common physiological disorders of potato tubers found in Swaziland and their management.

Greening physiological disorder is common to potatoes produced in all the 4 agro-ecological zones of Swaziland, especially following heavy rains that wash the soil and expose the potato tubers. This physiological disorder can also be experienced on potatoes that are about to be harvested especially when you have planted varieties like Mondial and Sifra. Exposing potatoes to sunlight can also lead to the occurrence of this physiological disorder, as most of our farmers do not have proper storage facilities that can deter light to enter the harvested produce.

Internal brown fleck is common where potatoes are grown in soils lacking calcium. This is more common in the Highveld where there is a lot of nutrient leaching due to excessive rains in most cases. Loose skin and thumbnail damage are most common to potatoes that are harvested using machinery. These physiological disorders are also prone to places like the Highveld, especially in soils lacking calcium and magnesium. Malformation is not very common to potatoes grown in the Highveld, but it is common to potatoes grown in the Lowveld and lower Middleveld especially if the farmers did not plant during the winter season.

Netting and Sprouting are more common to most potatoes grown in the Lowveld. This is due to the fact that this region often receives

excessive heat and dry spells, thus promoting the occurrence of these physiological disorders. If the farmers did not adhere to the recommended planting time of potatoes, these physiological disorders become more prevalent. Vascular browning is most common to all potatoes produced in the 4 agro-ecological zones of Swaziland as it is caused by soil pathogens in most cases.

Commonly grown potato varieties in Swaziland's four agro-ecological zones were Mondial (most common/popular), Sifra, Avalanche and BP1. Susceptibility of a given variety to a particular physiological disorder is dependent on Genetics (G) X Environment/ Agro-ecological zone (E) X Management (M). Severe weather events due to climate change are likely to worsen the physiological disorders of potato tubers.

Encountered disorders are described and various ways to alleviate them are suggested and discussed. It is proposed going forward that the disorders are quantified and losses due to these disorders are determined and their impact on the industry recorded.

Growth cracks

This disorder is characterized by cracks on the tuber and the cracks

vary in length and depth. The tuber tissue is exposed however in some cases the skin is later formed. The causes of growth cracks include variation in soil moisture conditions, for example when a dry period is followed by ones watering or heavy rains leading to increased uneven growth and subsequent cracking. Similar conditions have been reported in tomato [21-23] a fellow member of the solanaceous family like potato. Other causes of growth cracks include uneven plant population, over fertilization with nitrogen at one time after tuber formation and nutrient imbalances [24]. Growth cracks disorder may be overcome by ensuring uniform irrigation through irrigation scheduling, uniform growing conditions and appropriate fertilizer management. Proper diagnosis is important so as not to confuse with cracks caused by herbicide or pathogen damage. The use of cultivars which are less prone to cracking is advised Model cultivar which is widely grown in the Southern African region may be avoided because it is more prone to cracking [7,24].

Thumb nail cracks

Thumb nail cracks as the name suggests appear as cracks made by the thumb nail. Mechanical impact is thought as the primal causal agent. A slight impact or injury to the periderm (skin) of the tuber can cause cracks, without bruising of the underlying tissue [24]. Chilling temperatures of $<10^{\circ}\text{C}$ makes tubers more prone to cracks. Low humidity conditions aggravate the condition where the crack desiccates and leads to thumb nail crack and subsequent lower tuber quality. Cracks also provide ingress for diseases entry pathogens like soft rots bacteria. The minerals Magnesium (Mg) and Calcium (Ca) have been implicated. Susceptible Cultivars (cvs) have been found to have low Mg and Ca content of cell walls [25]. Magnesium and Calcium are components of pectin of the cell wall. To avoid tubers succumbing to thumb nailcracks the crop should be harvested and handled under cool conditions. On potato plant nutrition there is need for optimal fertilization with Ca and Mg in order to get relatively strong tuber cell walls [24,25].

Enlarged lenticels

Enlarged lenticels disorder manifests as white callus like tissue on the surface of tubers especially when tubers have been removed from wet soil. Under dry conditions the raised white callus tissue desiccates and become sunken. When brushed the raised lenticels are removed and under favourable conditions of suberization wound healing occurs. Possible tuber infection by pathogens is possible before wound healing such that in warm weather subsequent soft rot may occur. Conditions of restricted oxygen supply and excessive moisture conditions around tubers may cause enlarged lenticels to occur [24,26]. Unwanted enlarge lenticels conditions of potato tubers may be avoided in many ways [25]. These include avoiding water logged conditions and reducing irrigation water supply by 40-50% two weeks before harvesting. Harvested tubers with the physiological disorder must be dried as soon as possible before packing and where post-harvest washing with chlorine treated water should be done at a $\text{pH} > 7$. On the other hand seed potatoes from waterlogged land must be kept away from those from other conditions in the event of subsequent moist conditions prevailing [25].

Greening

This disorder is characterized by a green colour on the tuber surface, indicating the accumulation of chlorophyll. The intensity may

differ and may also be accompanied by sunburn [24,27,28]. Greening not only affects external quality, but causes bitter taste and can pose a health risk as a result of Glycoalkaloid *essolan* in which can be moderately toxic for humans, if consumed in large quantities [24-26]. White skinned varieties like BP1, Avalanche, Sifra and Mondial often green more readily than the red or russet varieties. This physiological disorder is caused by potato tuber exposure to light in the field, resulting in formation of a green pigmentation on the potato. This occurs when sunlight directly contacts tubers growing at or near the soil surface or reaches tubers through cracks in the soil surface. Greening typically affects a limited part of the tuber [7,29]. The same effect can be caused by extended exposure to lowlight levels in storage or on store shelves. In this case, the result typically is a lighter, more diffuse colouration on the entire tuber. Excessive nitrogen application early in the season may cause stolons to grow longer than normal, thus tubers are exposed to light on the land, and exposure after harvesting will aggravate greening [24,30].

Preventative steps to control this physiological disorder include proper seeding depth, hilling to cover exposed tubers, reducing the time potatoes are exposed to natural light and eliminating exposure to artificial light during storage. If possible, irrigate to prevent cracks occurring during dry weather conditions. Avoid high nitrogen application, especially with cultivars that are prone to form long stolons. Variety choice should also be considered. Use brown packaging materials that provide sufficient protection against light transmission if potatoes are to be exposed to light for more than a few days during marketing [7,24,31,32].

Blackheart

Black heart is often confused with soft rot. This physiological disorder is very common in large tubers. A diagnostic method is to cut longitudinally the potato to see its symptoms. Black heart symptoms differ from soft rot symptoms in that tissue with soft rot is soft and watery, and can often develop on the surface of the tuber. With secondary infection of tubers, a distinctive unpleasant odour becomes present [24,28].

Blackheart is a tuber condition that results from asphyxiation of the tissues of the tuber. This condition occurs in storage or transit when tubers are subjected to very high temperatures are stored where the ventilation is so poor that the supply of oxygen is inadequate, or results from a combination of both factors [33,34]. An oversupply of carbon dioxide whereby the oxygen supply levels are reduced can be another cause [25,26]. Black heart can also occur in the potato field if high temperatures follow excessive rain which causes an oxygen shortage in the soil. The internal symptom is a dark discoloration, greyish to purplish or inky black. Generally, this discoloration is restricted to the heart of the tuber. Usually, the discoloured tissues are sharply set off from healthy tissues. They are firm and leathery if they have dried out [31,35]. In advanced stages affected tissues dry-out and shrink, forming cavities. When tubers are cut soon after injury, the exposed tissues are of normal colour. Shortly after exposure to air, however, they turn pink, then grey, purplish, or brown, and finally become black. Sometimes only grey or brown discoloration is found. This occurs when tubers have been heated to above 55°C or when they have been deprived of all oxygen for considerable periods of time, as in waterlogged soils or in flooded storage pits. The external symptoms of blackheart, which rarely occur, are moist areas on the tuber surface.

These areas may be purplish for a short time but, characteristically, these discoloured areas are brown or black [1,29].

To control blackheart, it is recommended that high-temperature storage should be avoided and storage areas should be well ventilated. Seed tubers in furrows should be covered as soon as possible after planting and poorly draining fields should be avoided. After harvesting potatoes during hot weather, the tubers should be promptly removed from the soil surface. To prevent oxygen shortages, tubers should not be stored in solid piles higher than 1.8 meters, even at low temperatures [1,7,24,25,29].

Internal brown spot

This physiological disorder is often confused with zebra chip. Internal brown spot is frequently called by other names such as internal necrosis, internal brown fleck, internal rust spot, internal browning, and, in part, sprain [24]. Affected tubers do not display foliage or external symptoms of brown spot. Internal brown spot is characterized by irregular, dry, brown spots or blotches scattered throughout the flesh of affected tubers. These spots are not restricted to the water vessels, as in vascular diseases, but are often found in the central parts of the tubers, in the vascular bundle ring. The spots are a result of groups of dead cells that are free from bacteria and fungi [24,36]. The primary cause of brown spot is shortages of calcium in the soil which results in loss of integrity of cell membranes and cell walls under stress conditions. Environmental factors that can lead potato tuber to brown spot include: soil temperature, soil types and soil moisture. Uptake of water and nutrients happens very slowly through the roots if air temperature is high but the soil temperature is low, resulting in water and nutrients being withdrawn from the tubers causing stress in tuber cells. The roots end-up not performing optimally, resulting in insufficient water and calcium uptake, thus leading to the loss of membrane integrity which causes cells to die and turn brown [24-26,28]. Internal brown spot occurs most often in tubers grown in sandy soils because their water holding capacity is low and soil temperature tend to increase quickly; however, it has also been reported in potatoes grown on muck soils [7,30,31]. A similar disorder in tomatoes caused by lack of calcium is called blossom end rot.

Other factors that are implicated with this defect are lack of available soil moisture, particularly late in the growing season; lack of water at some critical stage in the growth of the plant because of either poor soil texture or alternating wet and dry weather; and excessive evaporation of moisture from the foliage. Internal brown spot can also develop during storage if the calcium content is low and if tubers were harvested under high temperature conditions [24,28,32].

To control this physiological disorder, favourable cultural conditions, such as proper irrigation practices and covering the tubers with five centimetres or more of soil, reduce the severity of internal brown spot. Sufficient calcium must be available for uptake where and when tubers are formed. Opting for planting times to avoid high temperatures during tuber formation and harvesting, and ensuring calcium uptake is not limited by an imbalance between nutritional elements or unfavourable pH reduces risks associated with internal brown spot [24,25,28,31,32].

Malformation

Malformation is often called second growth or knobby tubers. This physiological disorder may result in tubers exhibiting different

malformation depending on when the growing malformation took place, i.e., dumbbell, elongated, bottleneck or tubers with multiple knob-like protuberances. This physiological disorder is often confused with herbicide damage. Malformation is a condition produced in potatoes when high temperatures accompanied by moisture stress and nutrient shortage during midseason, resulting to second growth occurring on the main or primary tuber at sites where eyes are located [25,26]. This abnormal growth may also be a result of irregular irrigation, especially when overhead irrigation methods like sprinklers and pivots are used. No cultivar is resistant to malformation, but cultivars with an elongated tuber shape (Mondial and Sifra) are more subject to knobiness than other varieties [24,29]. Other factors that play a role in occurrence of malformation include poor plant population, single haulm plants, incidence of diseases such as black scurf as well as size and age of seed potatoes and insufficient fertilisation [7,24,27].

To control secondary growth (knobby tubers, there is need to; provide uniform growing conditions through out the growing season. Planting of potatoes should be done during the cooler times of the year. Avoid production in periods when heat waves generally occur usually during mid-summer. Irrigation should be maintained uniformly throughout the growing season especially in the warm areas like the Lowveld of Swaziland. Avoid planting those varieties that tend to be excessively knobby, such as Mondial and Sifra and plant varieties (BP1 and Avalanche) that produce round tubers [24-27].

Cold damage

Cold damage can be confused with soft rot symptoms. Symptoms are somewhat variable depending on the temperature, exposure period, and potato cultivar. Symptoms appear on warming after frost due to probably extreme weather events, and due to climate change which may manifest in the Highveld of Swaziland. Externally, the tuber appears wrinkled and flabby, whereas internal tissues generally turn bluish grey to black in response to cold temperatures [24]. Chilling damage to tubers occurs at temperatures below 3°C or below freezing point. Seed potatoes stored under conditions of insufficient protection against low temperatures and cold winds, e.g., stacked under trees, are prone to cold damage [24,28,29]. Internal symptoms include diffuse smoky grey areas inside the tubers. Tubers or portions of tubers that have been frozen at temperatures below -1°C are soft and watery. Frozen tissues tend to disintegrate and will eventually dry out. Symptoms of chilling and freezing can, but do not always, occur in the same tuber [28,29]. To manage this physiological disorder, harvested tubers should not be left on the field overnight, especially if low temperatures are expected as in the cool season in the Highveld of Swaziland. Do not dig if a frozen crust is present on the ground. Frost protection should be provided in storage and in transit. Cultivars that tend to bear shallow in areas with low temperatures at the end of the growing season should be avoided. Harvest tubers close to the soil surface first before the first cold front is expected [24,28,29].

Brown core/centre and hollow heart

Brown core and hollow heart are internal non-infectious physiological disorders that occur in potato tubers. These physiological disorders are commonly known for having a region of cell death in the pith of the tuber resulting in browning of tissue. Neither of these disorders has been reported as harmful and neither affects the tuber's taste or nutrition. Brown core frequently precedes the development

of hollow heart, but both these disorders often occur separately [24,28,29].

Brown centre and hollow heart arise at a higher incidence when growing conditions abruptly change during the season, such as when potato plants recover too quickly after a period of environmental or nutritional stress. When the tubers begin to grow rapidly, the tuber pith can die and/or pull apart, leaving a void in the centre. Brown centre and hollow heart effects likely form during tuber initiation but could also form during tuber bulking [24,37]. Incidence of brown core and hollow heart also increases with periods of stress because of high or low moisture level, more so if the heavy water applications follow a period of stress because of low moisture levels. If the disorder occurs during the early part of the season, then it is most often preceded by brown centre and forms in the stem-end of the tuber, while late-forming hollow heart [28,37].

Brown core and hollow heart can negatively affect tuber quality as these disorders make cut fresh market tubers unattractive and can result in most potatoes failing to make the grade. To prevent these physiological disorders it is important to select varieties that are known to be resistant and by planting when soil temperature reaches adequate levels to lessen the occurrence of these physiological disorders. Potato should be larger and less aged to reduce brown centre and hollow heart because of increased stem number per seed piece. Other practises that can reduce incidence of brown core and hollow heart include: achieving recommended stand establishment, avoiding planting skips, and applying multiple split fertiliser applications especially nitrogen [24,28,37].

Vascular browning

Vascular browning is a physiological disorder that causes an unacceptable vascular browning and can cause economic losses to the grower. The browning can be found in tubers in the potato field as well as in storage during the first one to two months. Symptoms of vascular browning include: vascular browning developing at the stolon end and vascular browning appears as speckled, light brown to brown and even dark stripes. Immature tubers are more susceptible, but vascular browning is common in tubers approaching maturity [24,28]. This physiological disorder is associated with a combination of low soil moisture and sudden foliage die-off as a result of chemicals, frost or mechanical removal, or a combination of these factors. This physiological disorder can be accelerated by high temperature during foliage die-off, resulting in the seriousness of discolouration. Tuber-borne pathogens such as *Verticillium* and *Fusarium* and leaf roll virus can also induce vascular browning. This physiological disorder can be managed by irrigating prior to foliage die-off, irrespective of the method of foliage die-off [24,28,36,38-40].

Spouting

When potato tubers are harvested, they normally undergo a rest period through dormancy, an adaptation mechanism they use to survive unfavourable environmental conditions [41]. This rest period is where potato derives its usefulness for cooking, processing as well as for seed. Initially, the period is regulated by endodormancy. Following the endodormancy period, sprouting can occur anytime if not controlled. However, its timing and regulation has been an area of study for a long time. Sprouting results in quality loss through remobilisation of starch and proteins. As a consequence, sprouting is

accompanied by remobilisation of nutrients and shrivelling. Although still vaguely understood, literature has linked it to sucrose, phytohormones, levels of ethylene and ABA [41]. Attempts have been made to reduce sprouting disorder. These include regulation and control of sucrose within the tuber, regulation of hormones signalling sprouting and use of sprout inhibitors such as chlorpropham (isopropyl N-chlorophenyl carbamate, CIPC). In addition, management of environmental conditions promoting sprouting, including temperature control continues to be useful in reducing sprouting. Pre and post-harvest conditions controlling sprouting which need attention include day length during the growth of tubers, nutrients, water supply and temperatures [28,37].

Netting

Netting in potatoes relates to a condition characterised by a network of fine cracks or fissures in the outer layers of the potato outermost skin, also known as periderm. Potato periderm comprise of the three major layers, viz: the phellem, phellogen and phelloderm. Netting tends to affect the phellem, which is the outermost layer of the periderm. The disorder has been reported to start as faint lines and creases which eventually become netted in appearance in about 2.5 to 5 months post harvesting. A number of factors have been linked to this disorder. Immature periderm has been found to be susceptible to physiological damage [42]. Variety, soil texture, depth of planting, soil nutrition, soil temperature, water stress, water logging and length of growing season have been cited to contribute to phellem thickness and maturity and hence development of netting disorder. Conditions that promote thick skins also predispose potato to skin cracking characterised by resetting and netting. Water stress tends to increase netting while depth of placement of potato tubers also increased incidences of netting. Storage temperature was also found to influence netting disorder development, with low temperatures delaying netting. Literature on resetting is very scanty, so are control methods [28,37].

Loose skin

Skin set is critical stage of potato growth which ensures that the potato develops a skin that protects the potato from damage during harvesting and handling. However, this process may require long periods to sufficiently develop, up to 40 days [43]. When it comes to skin separation the activity of the phellogen comes into play [44]. Physical maturity is therefore important in determining the strength of the phellem and hence the potato in storage. During maturation of potato, various processes including suberization and integration in the phellem of fatty acid biopolymers and waxes occur, strengthening the skin of potato [43].

Management of skin physiological disorders

Current recommendations for the management of skin disorders include ensuring that tuber maturation is not interrupted. Often, farmers use the vine maturity to determine harvesting and hasten to harvest once vines show signs of maturity. Time should be allowed for the phellem to develop and set before harvesting can be done. This should be followed by adequate curing after harvest at optimum temperatures will help reduce these disorders. Vine desiccants when applied should be allowed sufficient time before potatoes is harvested, normally about three weeks [43]. Other practices such as adequate irrigation during tuber growth and maturation, appropriate fertiliser application

and soil conditioning will promote development of strong skin and reduce incidences on netting and loose skin [44].

Conclusion

Various physiological disorders of potato tubers occur under Swaziland agro-climatological conditions in all agro-ecological zones of the Kingdom. Climate change conditions due to global warming often result in extreme weather events and are likely to increase the incidence and severity of physiological disorders of potato tubers subsequently affecting potato tuber quality, nutritional and security contribution. There is urgent need to educate farmers on positive identification of common physiological disorders and ways of alleviating them including use of appropriate cultivars adapted to the four agro-ecological zones. Once awareness has been acquired by researchers, extension officers and local farmers then the 2022 visions becomes near reach so will be the Sustainable Development Goals (SDS) pertaining to health, nutrition and food security.

References

1. Miguel, C., 1985. Production and Utilization of Potatoes in the World. Academic press, London, UK.
2. Jackson SD (1999) Multiple signalling pathways control tuber induction in potato. *Plant Physiology* 119: 1-8.
3. Karim MR, Hanif MM, Shahidullah SM, Rahman AHMA, Akanda AM, et.al. (2010) Virus free seed potato production through sprout cutting technique under net-house. *African Journal of Biotechnology* 9: 5852-5858.
4. Masarirambi MT, Mandisodza FC, Mashingaidze AB, Bhebhe E (2012) Influence of Plant Population and Seed Tuber Size on Growth and Yield Components of Potato (*Solanum tuberosum*). *International Journal of Agriculture and Biology* 14:545-549.
5. Yamaguchi M (1983) *World Vegetables. Principles, Production and Nutritive Values*, AVI Publishing, Westport, Connecticut, USA.
6. Peirce LC (1987) *Vegetables: Characteristics, Production and Marketing*. John Wiley & Sons Inc., New York, USA.
7. Bohl WH, Johnsoneds SB (2010) *Commercial Potato Production in North America: The Potato Association of America Handbook*. The Potato Association of America, Ann Arbor, Michigan, USA.
8. Norman JC (1992) *Tropical Vegetable Production*. Arthur H Stockwell Ltd. Elms Court, Ifracombe, Devon, UK.
9. Rice RP, Rice LW, Tindall HD (1987) *Fruit and Vegetable Production in Africa*. MacMillan Publishers Ltd. London and Basingstoke, UK.
10. Zwane PE, Masarirambi MT (2009) Kenaf (*Hibiscus cannabinus*) and allied fibres for sustainable development in Swaziland. *Journal of Agriculture & Social Sciences* 5: 35-39.
11. Thompson CF (2009) *Swaziland Business Year Book*. Christina Forsyth Thompson, Mbabane, Swaziland, South Africa.
12. Masarirambi MT, Mhazo N, Oseni TO, Shongwe VD (2009) Common Physiological Disorders of Tomato (*Lycopersicon esculentum*) Fruit Found in Swaziland. *Journal of Agriculture & Social Sciences* 5: 35-39.
13. Masarirambi MT, Oseni TO, Shongwe VD, Mhazo N (2011) Physiological disorders of Brassicas/Cole crops found in Swaziland: A review. *African Journal of Plant Science* 5: 8-14.
14. Verma P (2009) Physiological Disorders of Vegetable Crops. In: Russo VM (ed.). Alfa Beta Technical Solutions. *International Journal of Vegetable Science* 16: 1-170.
15. Ceponis MJ, Cappellini RA, Lightner GW (1987) Disorders in cabbage, bunched broccoli and cauliflower shipments to the New York market, 1972-1985. *Plant Disease* 71: 1151-1154.
16. Chiang MS, Chong C, Landry BS, Crete R (1993) Cabbage (*Brassica oleracea* subsp *Capitata* L.) In: Kaloo G, Bergh BO (eds.). *Genetic Improvement of Vegetable Crops*. Pergamon Press, Oxford, UK.
17. Jarvis WR, McKeen CD (1991) *Tomato Diseases*. Agriculture Canada Publication, Ottawa, Canada.
18. Government of Swaziland (2007) *Swaziland Review*. Ministry of Commerce and Industry. Government of Swaziland, Mbabane, Swaziland, South Africa.
19. Dlamini GM, T Lupapa (1995) Swaziland: Country report to the FAO International technical conference on plant genetic resources. FAO, Rome, Italy.
20. Government of Swaziland (2005) *The Swaziland National Biodiversity Strategy and Action Plan*. Government of Swaziland, Mbabane, Swaziland, South Africa.
21. Peet MM (1992) Fruit cracking in tomato. *Hort Technology* 2: 216-223.
22. Wien HC (1997) *The Physiology of Vegetable Crops*. CAB International, New York, USA.
23. Dorais M, Papadopoulos AP, Gosselin A (2001) Greenhouse tomato fruit quality. *Wiley online library, Horticulture Reviews* 5: 239-319.
24. Potatoes South Africa (2016) *Physiological tuber disorders*. Potatoes South Africa, Pretoria, South Africa.
25. Hiller LK, Thornton RE (2008) *Managing Physiological Disorders*. In: Johnson DA, Powelson ML (eds.). *Potato Health Management: Plant health management series*. APS Press, USA. Pg no: 261.
26. Hiller LK, Koller DC, Thornton RE (1985) *Physiological Disorders of Potatoes*. In: Li PH (ed.). *Potato Physiology*. Academic Press, New York, USA. Pg no: 586.
27. Stark JC, Love SL (2006) *Potato Production Systems*. Educational Communications, University of Idaho, Moscow, USA.
28. Selman LN, Andrews N, Stone A, Mosley A (2008) What's Wrong with my Potato Tubers? Diagnosing tuber abnormalities in western Oregon and Washington. Oregon State University, Oregon, USA.
29. Snowden AL (1990) *A Colour Atlas of Post-harvest Diseases and Disorders of Fruits and Vegetables: General introduction and fruits*. Wolfe, London, UK.
30. Zotarelli L, Dittmar PJ, Roberts PD, Noling JW, Wells B (2012) *Potato Production*. In: Zotarelli L, Dittmar PJ, Roberts PD, Noling JW, Wells B (eds.). *Vegetable Production Handbook of Florida*, University of Florida, Institute of Food and Agricultural Sciences, Gainesville, USA.
31. Paul L (1985) *Production and Storage of Potatoes*. John Wiley and Sons, London, UK.
32. Zaag PV, Demagante AL, Ewing EE (1990) Influence of plant spacing on potato (*Solanum tuberosum* L.) morphology, growth and yield under two contrasting environments. *Potato Research* 33: 313-323.
33. Lipton WJ (1967) Some effects of low-oxygen atmospheres on potato tubers. *American Potato Journal* 44: 292-299.
34. Reeve RM (1968) Further histological comparisons of back spot, physiological internal necrosis, black heart and hollow heart in potatoes. *American Potato Journal* 45: 391-401.
35. Stevenson W, Loria R, Franc G, Weingartner D (2001) *Compendium of potato diseases*. American Phytopathological Society, St. Paul, MN, USA.
36. Rich AE (1983) *Potato Diseases (1stedn)*. Academic Press, New York, USA.

37. Bussan AJ (2007) The Canon of Potato Science: 45. Brown Centre and Hollow Heart. *Potato Research* 50: 395-398.
38. Hooker WJ (1981) *Compendium of Potato Diseases*. APS Press, St. Paul, MN, USA.
39. Rich AE (1981) Verticillium Wilt In: Hooker WJ (ed.). *Compendium of Potato Diseases*. International Potato Center, St. Paul, MN, USA. Pg no: 62-63.
40. Davis JR, Huisman OC, Westermann DT, Hafez SL, Everson DO, et al. (1996) Effects of green manures on Verticillium Wilt of potato. *Phytopathology* 86: 444-453.
41. Sonnewald S, Sonnewald U (2014) Regulation of potato tuber sprouting. *Planta* 239: 27-38.
42. Lulai EC (2007) The Canon of Potato Science: 43. Skin-set and wound-healing / suberisation. *Potato Research* 50: 387-390.
43. Bussan AL, Sabba RP, Drillas MJ (2009) Tuber maturation and potato storability: Optimising skin set, sugars and solids. University of Wisconsin-Extension, Cooperative Extension Cooperative Extension Publishing, Madison, Wisconsin, USA.
44. British Potato Council (2006) Preserving potato skin finish during storage: Research Review. British Potato Council, UK.