

Research Article

Effect of Genetic Factors on Atopic and Non Atopic Asthmatic and Allergic Rhinitis Saudi Children in Taif Area

Yousri M Hussein^{1,2*}, Abd Alrahman N Alghamdy³, Haytham A Dahlawi¹, Rana G Zaini¹, Abdulmajeed A Almaleki¹, Ali Kh Al-Zahrani⁴, Mazen M Almeahadi¹ and Emad T Ahmed⁵

¹Department of Medical Laboratories, Taif University, Saudi Arabia

²Department of Medical Biochemistry, Zagazig University, Zagazig, Egypt

³Department of Community Medicine, Taif University, Taif, Saudi Arabia

⁴Department of Pediatrics College of Medicine, Taif, Saudi Arabia

⁵Department of Physical Therapy, Taif University, Saudi Arabia

Abstract

Objectives

To assess the value of serum Interleukin-13 (IL-13) levels as an immunological marker in atopic upper respiratory diseases, to clarify its differences in atopic and non atopic bronchial asthma and to determine the role of an Interleukin -13 receptor alpha 1 (IL-13 R α_1) gene Single Nucleotide Polymorphism (SNP) (A1398G) in the pathogenesis of these diseases.

Methods

Seventy-five patients were compared with 25 age-matched healthy volunteers. Serum total immunoglobulin E (Ig E) and IL-13 levels were measured by enzyme-linked immune sorbent assay and the IL-13R α_1 gene (A1398 G) was screened by specific polymerase chain reaction.

Results

There was a non significant association between G allele frequencies of the IL-13R α_1 (1398) gene polymorphism (42%, 38% and 30% for atopic asthma, non atopic asthma, and allergic rhinitis, respectively) as compared to in controls. There were a significant increase in the serum level of total IgE & IL-13 towards heterozygous AG and homozygous GG than homozygous AA in atopic asthma, non atopic asthma, and allergic rhinitis patients. There was a significant increase in the serum level of total IgE & IL-13 towards homozygous GG than heterozygous AG in atopic asthma ($p=0.035$), non atopic asthma ($p=0.014$), and allergic rhinitis patients ($p=0.003$) for IgE and ($p<0.001$) in all groups for IL-13 as shown by LSD test.

*Corresponding author: Yousri M Hussein, Department of Medical Laboratories, Taif University, Saudi Arabia, Tel: +966 201001752426; E-Mail: yousrihussein@hotmail.com

Citation: Hussein YM, Alghamdy AAN, Dahlawi HA, Zaini RG, Almaleki AA, et al. (2017) Effect of Genetic Factors on Atopic and Non Atopic Asthmatic and Allergic Rhinitis Saudi Children in Taif Area. J Allergy Disord Ther 3: 007.

Received: April 26, 2017; Accepted: June 02, 2017; Published: June 19, 2017

Conclusion

Serum interleukin (IL) 13 can be used as an immunological marker in atopic upper respiratory diseases and to differentiate between atopic and non atopic bronchial asthma.

Keywords: Immunoglobulin E (Ig E); Interleukin -13 Receptor Alpha 1 (IL-13 R α_1); Single Nucleotide Polymorphism (SNP)

Introduction

Asthma is associated with atopy and with IgE-mediated inflammation of the airways, which occurs via release of Interleukin (IL)-4 and IL-13, both of which have been included in the pathogenesis of asthma in multiple human and animal studies [1]. Human IL-13 is a protein with a molecular mass of 13 KDA, which has four α helical bundles (α A, α B, α C and α D) [2]. In Saudi Arabia, the prevalence of asthma is higher than in other Arab countries, with substantial regional variations [3-5]. Saudi Arabia has increasing prevalence of childhood atopic asthma. Allergic disease was reported to be very common in primary school-aged children in different areas of Saudi Arabia, including Taif city, with figures closer to the highest risk regions in the world [6-9].

It has an important role as an effector molecule in asthma through multiple mechanisms, including induction of production of immunoglobulin (IgE) by B-cells [10], attraction of eosinophils to the airway [11], metaplasia of goblet cell and increase mucus secretion, and airway remodeling [12]. A wealth of data supports a role for IL-13 in mediating asthma pathology. IL-13 produces its effect via receptor which is heterodimeric and composed of two membrane proteins, [13] IL-4R α and either a low affinity IL-13R α_1 or high affinity IL-13R α_2 . [14] IL-13 binds to IL-13 binding chain (IL-13R α_1) at low affinity in the absence of IL-4R α , whereas in the presence of IL-4R, the site becomes of high affinity [15]. IL-13R α_1 is widely expressed and has been detected on nearly every cell tested except human T cells, while, the human IL-13R α_1 gene is present on the X q24 chromosome possibly suggesting a role in X-linked immune disease [16].

The C-terminal alpha helix D, which is one of the four alpha helices that constitutes IL-13, contains key residues for binding with both IL-13R α_1 and IL-13R α_2 , whereas IL-13 interaction with IL-4R α is mediated via helices A and C [3]. Analysis of crystal structures of ternary complexes composed of IL-13 or IL-4 binding with IL-13R α_1 and IL-4R α chain gives idea about shared receptor interactions with distinct cytokine, explaining the different affinities of cellular responses to IL-4 and IL-13 [16]. Association of the IL-13R α_1 /IL-4R receptor complex by IL-13 results in stimulation of a variety of signal transduction pathways. Upon forming a dimer of IL-13R α_1 with IL-4R α , JAK1 and Tyk-2 kinases become phosphorylated and activated which lead to phosphorylation of tyrosine residues on the IL-4R α and IL-13R α_1 chains [17].

Polymorphisms in these receptor molecules contribute to the genetic effects on asthma susceptibility and atopic diseases as allergic rhinitis. For example, asthma susceptibility was significantly elevated in Korean children by gene-gene interaction between IL-4 T-590C and IL-4R α Gln 551 Arg alleles [18-19] and between IL-13 C-1112T and IL-4R α Ser478 Pro alleles in Dutch population [20]. Also, a non

coding polymorphism in IL-13Ra1 (A+1398G) was linked to increased IgE levels in British population [21].

On the basis of the important role of the IL-13/IL-4 pathway in atopy and asthma, it was hypothesized that genetic variation in IL-13Ra1 may lead to the development and/or predict severity of asthma and atopy [22-24].

Patients and Methods

The study was conducted by the Clinical Laboratory Department Faculty of Applied Medical Sciences and Pediatric Departments at the Faculty of Medicine Taif University, Saudi Arabia.

Participants

Seventy-five patients and 25 healthy controls were studied. None of the participants received antihistamine, systemic or topical corticosteroids in the 3 weeks prior to clinical evaluation and they all underwent skin prick testing. The patients were divided into 3 groups: 1) those with atopic bronchial asthma (n=25), 2) those with non atopic bronchial asthma (n=25) and 3) those with allergic rhinitis (n=25), who met the criteria defined by Meltzer [25]. Asthma phenotypes and bronchial responsiveness were determined by a pediatric physician. Following the American Thoracic Society guidelines [26], asthma was confirmed by a history of chest tightness and wheezing during the previous 12 months, a greater than 12% reversibility of FEV1 spontaneously or after b2-agonist inhalation, and/or a methacholine provocation test result with a PC20 less than 16 mg/ml. Atopy was defined by a positive skin prick test (wheal diameter ≥ 3 mm) to at least one of 12 common aeroallergens [18]. Non-atopy was defined by negative skin prick test to the 12 common aeroallergens. Then on atopic control subjects had no history of asthma or other allergic diseases, negative skin prick tests, normal total IgE values (≤ 100 IU/ml), normal lung function tests, and no airway hyper responsiveness (PC20 >16 mg/ml). All the patients and healthy individuals gave their written consent before blood sample collection. In addition to a full history and clinical examination, stool and urine analysis was performed in all cases to exclude factors that could influence measurements.

Collection of blood samples

6 ml of blood was taken from each participant under complete aseptic conditions and divided into 2 portions; 1.5 ml of whole blood was collected in sterile EDTA-containing tubes for DNA extraction, and the rest was left for 30 to 60 minutes for spontaneous clotting at room temperature and then centrifuged at 3000 rpm for 10 minutes. Serum samples were separated into another set of tubes and kept frozen at -20°C for determination of total IgE and IL-13.

Determination of serum IL-13

Serum IL-13 was measured by sandwich Enzyme-Linked Immune Sorbent Assay (ELISA) (KOMABIOTECH Inc., Gayang Technotown, Korea), with the minimum detectable dose established as 1 pg/ml.

Total IgE measurements

Total serum IgE levels were also measured by sandwich ELISA (General Biologicals Corporation, Hsinchu, Taiwan), with assay sensitivity for total IgE of 5.0 IU/ml.

Detection of (A1398G) gene polymorphism

Genomic DNA was extracted from whole blood using abios-pingenomic DNA purification kit (Bioflux, Bioer, R&D Department,

Ferrotec, China). Participants were genotyped for the A1398G SNP in genomic DNA by PCR based restriction fragment length polymorphism [21]. Fragments were amplified in 50 μ l reaction mixtures containing 10 μ l genomic DNA, 30 μ l one step PCR mixture (1 unit Taq polymerase, 10 mM KCl, 10 mM (NH₄)₂SO₄, 20 mM TrisHCl (pH-8.75), 0.1% Triton X-100, 0.1 mg/ml BSA and 200 μ M dNTPs) and 1 μ l of each primer (BioBasic Inc., Ontario, Canada) and 8 μ l DdH₂O. The thermo cycling conditions consisted of an initial denaturation at 95°C for 12 min, followed by 35 cycles of denaturation at 95°C for 30 s, annealing for 2 min at 45°C and extension at 72°C for 40 s, followed by a final extension step at 72°C for 5 min, using a Gene-Amp PCR PTC-100 thermal cyclers (MJ Research, Inc., Watertown, Massachusetts, USA).

The primer sequences used for the IL-13Ra₁ (1398) gene polymorphism were as follows:

5' - TCA GTC ATG GAG ATA ATT TA 3' (sense)

5' - TGA GCT GCC TGT TTA TAA AT 3' (antisense)

The products were digested using MseI (New England Biolabs), which digested the +1398A allele into 85 and 45 bp fragments and yielded a single 130 bp band for the +1398G allele. The 20 μ l of PCR products were digested with 5 U of the restriction enzyme at 65°C for 16 h, and separated on 3% agarose gel stained with ethidium bromide.

Data analysis

Statistical analysis was conducted using version 11 of the statistical package SPSS for Windows [27]. A P value of <0.05 was considered statistically significant.

Results

(A1398G) Frequencies

The frequencies of the AA, AG, and GG genotypes of IL-13Ra1 gene polymorphism were, respectively, 52%, 32%, and 16% in controls, 40%, 36%, and 24% in atopic asthma patients, (Figure 1) 44%, 36%, and 20% in non atopic asthma patients, and 56%, 28%, and 16% in allergic rhinitis patients. The X² values for atopic asthma, non atopic asthma and allergic rhinitis were 3.4, 1.35, and 41, respectively, and the presence of these genotypes was not significantly associated with the presence of atopic conditions or with non atopic asthma compared to the control group (P=0.182, P=0.51, and P=0.812, respectively) (Table1).

	Control Group	Atopic Asthma Group	Non Atopic Asthma Group	Allergic Rhinitis Group
	No Percent	No Percent	No Percent	No Percent
AA	13 52%	10 40%	11 44%	14 56%
AG	8 32%	9 36%	9 36%	7 28%
GG	16 4%	24 6%	5 20%	4 16%
X ²		3.4	1.35	0.41
p		0.182(>0.05)	0.5 (>0.05)	0.812(>0.05)

Table 1: Genotype Frequencies for IL-13Ra₁ gene polymorphism in all studied group.

As compared to control group (statistical significance, P<0.05). P values calculated using the X² test.

1398G allele frequencies

There was non significant association between G allele frequencies of the 1398 polymorphism in the atopic asthma, non atopic asthma,




Figure 1: Genotype frequencies for IL-13Rα1 gene polymorphism in atopic asthma group.

and allergic rhinitis groups compared to the control group ($P > 0.05$ for each group) (Table 2). In atopic asthma OR=1.5; 95% CI (0.8-2.7), non atopic asthma (OR=1.3; 95% CI; 0.7-2.3), and allergic rhinitis (OR=0.9; 95% CI, 0.5-1.65) (Table 2).

Group	No percent	Odds Ratio (95% CI)
Control group	16 32%	
Atopic asthma	21 42%	1.5 (0.8-2.7)
Non atopic asthma	19 38%	1.3 (0.7-2.3)
Allergic rhinitis	15 30%	0.9 (0.5-1.65)

Table 2: Odds ratio for 1398 G allele in studied groups.

Association between (A1398G) polymorphism and IL-13 and IgE Levels

We studied the association between the parameters measured and the allelic variants of the (A1398G) polymorphism by analysis of variance and showed that there was a significant increase in the serum level of total IgE & IL-13 towards heterozygous AG and homozygous GG than homozygous AA in atopic asthma ($p < 0.001$), non atopic asthma ($p < 0.001$), and allergic rhinitis patients ($p < 0.001$) as shown by LSD test but not in control group. There was a significant increase in the serum level of total IgE & IL-13 towards homozygous GG than heterozygous AG in atopic asthma ($p = 0.035$), non atopic asthma ($p = 0.014$), and allergic rhinitis patients ($p = 0.003$) for IgE and ($p < 0.001$) in all groups for IL-13 as shown by LSD test. But not in control group (Table 3).

IL-13 and IgE levels

There was a highly significant increase of serum total IgE & IL-13 as regard atopic asthma group ($p < 0.001$), and allergic rhinitis group ($p < 0.001$) as compared with the control group and in atopic asthma and allergic rhinitis when compared with non atopic asthma ($p < 0.001$). There was no statistically significant difference between non atopic asthma and control group, or between atopic asthma and allergic rhinitis groups as regard serum total IgE & IL-13 ($p > 0.05$) (Table 4).

Discussion

The prevalence of asthma, allergic rhinitis, has increased greatly over the past years [28]. Atopy is characterized by the interaction between multiple genetic and environmental factors. It has been well established that genetic factors have an important effect on the risk of developing atopic disease and several genome-wide searches have proved linking atopy to loci on multiple chromosomes [29,30]. IL-13 produces its effect through its receptor, a hetero dimer composed of IL-4Ra and IL-13Ra1. Firstly, IL-13 binds to the IL-13Ra1 chain on

the surface of cells with an affinity of approximately 10^{-8} to 10^{-9} M in both mouse and human systems [31]. Upon IL-13 association with IL-13Ra1, IL-4Ra is recruited to form the high affinity ($\sim 10^{-10}$ to 10^{-11} M) receptor complex [23-31].

However, IL-13 responsiveness has been reported in the absence of IL-4Ra, suggesting the presence of alternate receptor forms. AHR, airway mucus production, and lung eosinophilia were distinguished in mice with transferred OVA-specific IL-13 producing T cells but lacking IL-4Ra, but not in those lacking STAT6 [32]. Also, in mice lacking IL-13 airway inflammation, fibrosis, and mucus cell hyperplasia, were diminished [33], but persisted in animals deficient in IL-4 [34] or IL-4Ra [33], suggesting that IL-13 may produce its effects through an IL-4Ra-independent pathway in this asthma model [15-24,33-35].

Three fibronectin type III sub units (D1, D2, and D3) compose the extracellular r portions of IL-13Ra1 [36]. Significant impairment of the IL-13 response may be due to mutation of IL-13 and IL-13Ra, as found in mutation of Leu 319 and Tyr 321 in the D2 and D3 respectively of IL-13Ra [36]. IL-13 induced signals or responses was not able to be mediated through IL-13Ra1 lacking the intracellular domain, supporting the possibility that IL-13Ra1 is required for signaling [37].

We reported that there was a no significant association between IL-13Ra1 polymorphism (+A1398G) and the susceptibility of atopic asthma, non atopic asthma or allergic rhinitis in Egyptian children while hetero- or homozygosity for the risk allele of IL-13Ra1 A+1398G was significantly associated with increased total IgE & IL-13 levels in children with atopic asthma, non atopic asthma and allergic rhinitis. Konstantinidis et al., [38] reported that no association between IL-13Ra1 (A+1398G) and asthma susceptibility or severity or with the development of atopic phenotype in Caucasian families while, IL-13Ra1 polymorphism (A+1398G) included in the control of IgE production.

Kim et al., [39] found that IL-13Ra1 (+A1398G) were not linked to the susceptibility of asthma or atopic asthma and that hetero- or homozygosity for the risk allele of IL-13Ra1 A+1398G was significantly associated with increased total IgE levels in children with atopic asthma, non atopic asthma and allergic rhinitis. Heinzmann et al., [21] reported that non coding polymorphism in IL-13Ra1 (A+1398G) was associated with increased level of IgE in British population with restriction among different ethnic groups as in Japanese people. This may suggest that this non-coding polymorphism of IL-13Ra1 has a functional effect for the binding of IL-13, or alternatively that the IL-13Ra1 A+1398G polymorphism is associated with as yet undiscovered polymorphisms in the regulatory or coding regions of the gene encoding IL-13Ra1. Another possibility is that IL-13Ra1 could have additional unknown signaling functions that are impacted by this polymorphism.

Collectively, these studies indicate that IL-13Ra1 is likely to play a critical role not only in binding but also in signaling of IL-13. We reported also a highly significant increase of serum total IgE & IL-13 levels in atopic asthma and allergic rhinitis groups as compared with the control group and in atopic asthma and allergic rhinitis when compared with non atopic asthma. There was no statistically significant difference between non atopic asthma and control group or between atopic asthma and allergic rhinitis groups and there is a positive correlation between IL-13 and IgE levels in these groups. This finding consistent with observations by EL-Helaly et al. [40], Turato et al., [41] and Gergen et al., [42] who reported from National health and nutrition examination survey (2005, 2006) that total IgE level

Groups	Measurement	AA	AG	GG	P ^b value
Atopic asthma	Serum IL-13 pg/ml	13.73 ± 0.46	15.98 ± 0.5	18.87 ± 1.06	<0.001
	Serum IgE IU/ml	244.8 ± 42.6	360.66 ± 10.67	394.3 ± 5.83	<0.001
Nonatopic asthma	Serum IL-13 pg/ml	4.37 ± 0.54	6.38 ± 0.69	7.88 ± 0.21	<0.001
	Serum IgE IU/ml	68.63 ± 9.99	99.66 ± 6.65	111.4 ± 2.19	<0.001
Allergic rhinitis	Serum IL-13 pg/ml	14.12 ± 0.82	16.82 ± 0.69	19.26 ± 1.0	<0.001
	Serum IgE IU/ml	255.2 ± 27.27	337 ± 14.47	384.5 ± 12.44	<0.001

Table 3: Association between IL-13Rα1 (A1398G) variant and Interleukin (IL) 13 levels and Immunoglobulin (IgE) levels.

b. calculated using analysis of variance

Measurement	Control	Atopic asthma	Non atopic asthma	Allergic rhinitis
IgE IU/ml	70.1 ± 18.5	322.32 ± 71.126*	88.36 ± 19.84	298.8 ± 56.85*
IL-13 pg/ml	5.828 ± 1.2	15.78 ± 2.1*	5.79 ± 1.5	15.7 ± 2.1*

Table 4: Serum Interleukin (IL) 13 and Immunoglobulin (Ig) E Levels in all Groups studied.

*- P value <0.05

predicted asthma among atopic subjects but not among non atopic individuals. Elevation of serum IL-13 in asthma group agreed with Lee et al., [43] who evidenced that the expression of IL13, IL4, and IL5 were increased in acute asthmatic patients so they may be deeply involved in the pathogenic process of asthma [44].

Yang et al., [45] suggested that anti-IL-13 monoclonal antibody inhibits airway hyper responsiveness, inflammation and airway remodeling in a chronic mouse model of asthma. These findings also confirmed with Kumar et al., [33] and Follettie et al., [46] who concluded that inhibition of IL-13 has considerable potential as a therapeutic strategy in chronic asthma. Thom et al., [47] optimized the affinity of a human IL-13-neutralizing antibody, a therapeutic candidate for the treatment of asthma, more than 150-fold. Ippoliti et al., [48] [who observed a significant reduction in asthma and rhinitis scores in the immunotherapy group compared with the placebo group associated with a significant decrease in IL-13 after 6 months of therapy.

We can explain the non significant difference in IL-13 level between atopic asthma and allergic rhinitis that both conditions have similar mechanisms and underlying pathogenesis, many of the cells, mediators, cytokines, and neurotransmitters involved in the biology of asthma and rhinitis are the same [22]. Indeed, up to 45% of patients with asthma have allergic rhinitis while 93.5% of those with allergic rhinitis were also asthmatic [49]. Feleszko et al., [50] found a positive correlation between IL-13 levels and serum IgE concentrations in children with allergic asthma. This can be explained by that the most important inducers of the release of IgE are IL-4 and IL-13 [51-54]. These cytokines stimulate transcription of the gene for the epsilon class of the constant region (Cε) of the immunoglobulin heavy chain [22,51,55-56]. In the other hand, Hussein et al., [57] denied this correlation suggesting that IL-13 role in the pathogenesis of allergic diseases was not only by inducing IgE production but also by other mechanisms. This hypothesis was concordant with animal models where mechanisms responsible for asthma and allergic rhinitis were independent of IgE levels [58].

However, other studies doubt on the role of serum total IgE as an important indicator of atopic diseases [59,60] found that both atopic and non atopic asthmatics had raised serum total IgE levels. The difference may be due to different population who were included in the study or to the different age groups or due to the severity of the

disease. Future work will be required to identify gene-gene and gene-environment interactions on a genome-wide level, with the aim of fully understanding the genetic risk factors for asthma and atopy, the pathogenesis of these common diseases, and perhaps describing new treatment strategies aimed at changing IL-13 signaling via IL-4Rα/IL-13Rα1.

Conclusion

The significant association of hetero- or homozygosity for the risk allele of IL-13Rα1 A+1398G with increased total IgE and IL-13 levels in children with atopic asthma and allergic rhinitis may suggest that this non-coding polymorphism of IL-13Rα1 has a functional effect for the binding of IL-13, or alternatively that the IL-13Rα1 A+1398G polymorphism is associated with as yet undiscovered polymorphisms in the regulatory or coding regions of the gene encoding IL-13Rα1.

Acknowledgment

Thanks to all staff members in Clinical Laboratory Department Faculty of Applied Medical Sciences and in Pediatric Department Faculty of Medicine, Taif University, Saudi Arabia.

References

- Flostrup H, Swartz J, Bergstrom A, Alm JS, Scheynius A, et al. (2006) Allergic disease and sensitization in Steiner school children. *J Allergy Clin Immunol* 117: 59-66.
- Moy FJ, Diblasio E, Wilhelm J, Powers R (2001) Solution structure of human IL-13 and implication for receptor binding. *JMB* 310: 219-230.
- Al-Dawood K (2000) Epidemiology of bronchial asthma among schoolboys in Al-Khobar city, Saudi Arabia: cross-sectional study. *Croat Med J* 41: 437-441.
- Al-Frayh AR, Hasnain SM (2007) Prevalence of bronchial asthma in children in Saudi Arabia. *World Allergy Organization Journal* 167-168.
- Al Frayh AR, Shakoor Z, Gad El Rab MO, Hasnain SM (2001) Increased prevalence of asthma in Saudi Arabia. *Ann Allergy, Asthma Immunol* 86: 292-296.
- Al-Makoshi A, Al-Frayh A, Turner S, Devereux G (2013) Breastfeeding practice and its association with respiratory symptoms and atopic disease in 1-3-year-old children in the city of Riyadh, central Saudi Arabia. *Breast Feed Med* 8: 127-133.
- Alshehri MA, Abolfotouh MA, Sadeg A, Al Najjar YM, Asindi AA, et al. (2000) Screening for asthma and associated risk factors among urban school boys in Abha city. *Saudi Med J* 21: 1048-1053.

8. Sabry EY (2011) Prevalence of allergic diseases in a sample of Taif citizens assessed by an original Arabic questionnaire (phase I) A pioneer study in Saudi Arabia. *Allergol immunopathol* 39: 96-105.
9. Nahhas M, Bhopal R, Anandan C, Elton R, Sheikh A (2012) Prevalence of allergic disorders among primary school-aged children in Madinah, Saudi Arabia: two-stage cross-sectional survey. *PLoS One* 7: 36848.
10. Defrance T, Carayon P, Billian G, Guillemot JC, Minty A, et al. (1994) Interleukin 13 is a B cell stimulating factor. *J Exp Med* 179: 135-143.
11. Wardlaw AJ (2001) Eosinophil trafficking in asthma. *Clin Med (Lond)* 1: 214-218.
12. Richter A, Puddicombe SM, Lordan JL, Bucchieri F, Wilson SJ, et al. (2001) The contribution of interleukin (IL)-4 and IL-13 to the epithelial-mesenchymal trophic unit in asthma. *Am j Respir Cell and Mol Biol* 25: 385-391.
13. Madhankumar AB, Mintz A, Debinski W (2002) Alanine-scanning mutagenesis of alpha-helix D segment of interleukin-13 reveals new functionally important residues of the cytokine. *J Biol Chem* 277: 43194-43205.
14. Akdis CA, Akdis M (2009) Mechanisms and treatment of allergic disease in the big picture of regulatory T cells. *J Allergy Clin Immunol* 123: 735-746.
15. LaPorte SL, Juo ZS, Vaclavikova J, Colf LA, Qi X, et al. (2008) Molecular and structural basis of cytokine receptor pleiotropy in the interleukin-4/13 system. *Cell* 132: 259-272.
16. Hussein YM, Ahmad AS, Ibrahim MM, Elsherbeny HM, Shalaby SM, et al. (2011) Interleukin 13 receptors as biochemical markers in atopic patients. *J Investig Allergol Clin Immunol* 21: 101-107.
17. Nelms K, Keegan AD, Zamorano J, Ryan JJ, Paul WE (1999) The IL-4 receptor: signaling mechanisms and biologic functions. *Annu Rev Immunol* 17: 701-738.
18. Lee SL, Wong W, Lau YL (2004) Increasing prevalence of allergic rhinitis but not asthma among children in Hong Kong from 1995 to 2001 (Phase 3 International Study of Asthma and Allergies in Childhood). *Pediatr Allergy Immunol* 15: 72-78.
19. Hussein YM, Awad HA, Shalaby SM, Ali AS, Alzahrani SS (2012) Toll-like receptor 2 and Toll-like receptor 4 polymorphisms and susceptibility to asthma and allergic rhinitis: A case-control analysis. *Cell Immunol* 274: 34-38.
20. Howard TD, Koppelman GH, Xu J, Zheng SL, Postma DS, et al. (2002) Gene-gene interaction in asthma: IL4RA and IL13 in a Dutch population with asthma. *Am J Hum Genet* 70: 230-236.
21. Heinzmann A, Mao XQ, Akaiwa M, Kreomer RT, Gao PS, et al. (2000) Genetic variants of IL-13 signalling and human asthma and atopy. *Hum Mol Genet* 9: 549-559.
22. Hussein YM, El-Tarhouny SA, Shalaby SM, Mohamed RH, Hassan TH, et al. (2011) Interleukin-13 receptor A1 gene polymorphism and IL-13 serum level in atopic and non-atopic Egyptian children. *Immunol invest* 40: 523-534.
23. Hussein Y, Allah SA, Mahmoud S, Ahmed A (2006) Impact of IL-13 gene mutations in atopic diseases.
24. Hussein Y, Abou El YM, Mahmoud Y, Mahmed H, Rasha L (2006) Some biochemical markers in atopic patients and PCR-based assay for detection of R576 IL-4 receptors allele gene. *Zagazig University Medical Journal* 12: 4307-4321.
25. Meltzer EO (1988) Evaluating rhinitis: clinical, rhinomanometric, and cytologic assessments. *J Allergy Clin Immunol* 82: 900-908.
26. [No authors listed] (1995) Standardization of Spirometry, 1994 Update. American Thoracic Society. *Am J Respir Crit Care Med* 152: 1107-1136.
27. Kirkwood B, Sterne AJC (1989) *Essentials of Medical Statistics*, (2nd edn). Blackwell Scientific Publication, London, England.
28. Sengler C, Lau S, Wahn U, Nickel R (2002) Interactions between genes and environmental factors in asthma and atopy: new developments. *Respir res* 3: 7.
29. Vercelli S, Ferriero G, Sartorio F, Stissi V, Franchignoni F (2009) How to assess postsurgical scars: a review of outcome measures. *Disabil Rehabil* 31: 2055-2063.
30. Mobasher A, El-Shahat H, Gabry M, Affara N, Hussein Y (2004) Eosinophilic Cationic Protein (ECP) in bronchial asthma, Role of treatment. *Egypt J Chest Dis & Tub* 53: 17-26.
31. Andrews AL, Holloway JW, Puddicombe SM, Holgate ST, Davies DE (2002) Kinetic analysis of the interleukin-13 receptor complex. *J Biol Chemistry* 277: 46073-46078.
32. Mattes J, Yang M, Siqueira A, Clark K, MacKenzie J, et al. (2001) IL-13 induces airways hyperreactivity independently of the IL-4R alpha chain in the allergic lung. *J Immunol* 167: 1683-1692.
33. Kumar RK, Herbert C, Yang M, Koskinen AML, McKenzie ANJ, et al. (2002) Role of interleukin-13 in eosinophil accumulation and airway remodelling in a mouse model of chronic asthma. *Clin Exp Allergy* 32: 1104-1111.
34. Foster PS, Ming Y, Matthei KI, Young IG, Temelkovski J, et al. (2000) Dissociation of inflammatory and epithelial responses in a murine model of chronic asthma. *Lab Invest* 80: 655-662.
35. Hussein YM, Shalaby SM, Nassar A, Alzahrani SS, Alharbi AS, et al. (2014) Association between genes encoding components of the IL-4/IL-4 receptor pathway and dermatitis in children. *Gene* 545: 276-281.
36. Arima K, Sato K, Tanaka G, Kanaji S, Terada T, et al. (2005) Characterization of the interaction between interleukin-13 and interleukin-13 receptors. *J Biol Chem* 280: 24915-24922.
37. Orchansky PL, Ayres SD, Hilton DJ, Schrader JW (2005) An interleukin (IL)-13 receptor lacking the cytoplasmic domain fails to transduce IL-13-induced signals and inhibits responses to IL-4. *J Biol Chem* 272: 22940-22947.
38. Konstantinidis AK, Barton SJ, Sayers I, Yang IA, Lordan JL, et al. (2007) Genetic association studies of interleukin-13 receptor alpha1 subunit gene polymorphisms in asthma and atopy. *Eur Respir J* 30: 40-47.
39. Kim H-B, Lee Y-C, Lee S-Y, Jung J, Jin H-S, Kim (2006) Gene-gene interaction between IL-13 and IL-13Ralpha1 is associated with total IgE in Korean children with atopic asthma. *Journal of Human Genetics* 51: 1055-1062.
40. El-Helaly N, El-Wan A, Kamel Y, Nabih M, Mahmoud H (2009) Eosinophil-Derived Neurotoxin Versus Immunoglobulin E as Biomarkers for Evaluation of Bronchial Asthma. *J Biol Chem* 9: 165-169.
41. Turato G, Barbato A, Baraldo S, Zanin ME, Bazzan E, et al. (2008) Nonatopic children with multitrigger wheezing have airway pathology comparable to atopic asthma. *Am J Respir Crit Care Med* 178: 476-482.
42. Gergen PJ, Arbes SJ Jr, Calatroni A, Mitchell HE, Zeldin DC (2009) Total IgE levels and asthma prevalence in the US population: Results from the National Health and Nutrition Examination Survey 2005-2006. *J Allergy Clin Immunol* 124: 447-453.
43. Lee YC, Lee KH, Lee HB, Rhee YK (2001) Serum levels of interleukins (IL)-4, IL-5, IL-13, and interferon-gamma in acute asthma. *J Asthma* 38: 665-671.
44. Abdel-Mawla MY, Mostafa Y, Abuel-Majd Y, Attwa R (2009) Detection of R576 interleukin-4 receptor an allele gene, serum interleukin-4, and eosinophilic cationic protein in atopic dermatitis patients. *Indian J Dermatol* 54: 31-35.
45. Yang G, Volk A, Petley T, Emmell E, Giles-Komar J, et al. (2004) Anti-IL-13 monoclonal antibody inhibits airway hyperresponsiveness, inflammation and airway remodeling. *Cytokine* 28: 224-232.
46. Follettie MT, Ellis DK, Donaldson DD, Hill AA, Diesl V, et al. (2006) Gene expression analysis in a murine model of allergic asthma reveals overlapping disease and therapy dependent pathways in the lung. *Pharmacogenomics J* 6: 141-152.
47. Thom G, Cockcroft AC, Buchanan AG, Candotti CJ, Cohen ES, et al. (2006) Probing a protein-protein interaction by *in vitro* evolution. *Proceedings of the National Academy of Sciences of the United States of America* 103: 7619-7624.

48. Ippoliti F, De Santis W, Volterrani A, Lenti L, Canitano N, et al. (2003) Immunomodulation during sublingual therapy in allergic children. *Pediatr Allergy Immu* 14: 216-221.
49. Cengizlier MR, Misirlioglu ED (2006) Evaluation of risk factors in patients diagnosed with bronchial asthma. *Allergol Immunopathol (Madr)* 34: 4-9.
50. Feleszko W, Zawadzka-Krajewska A, Matysiak K, Lewandowska D, Peradzyńska J, et al. (2006) Parental tobacco smoking is associated with augmented IL-13 secretion in children with allergic asthma. *J Allergy Clin Immunol* 117: 97-102.
51. Hussein YM, Alzahrani SS, Alharthi AA, Ghonaim MM, Alhazmi AS, et al. (2014) Association of serum cytokines levels, interleukin 10 -1082G/A and interferon- γ +874T/A polymorphisms with atopic asthma children from Saudi Arabia. *Cell Immunol* 289: 21-26.
52. Hussein YM, Shalaby SM, Zidan HE, Sabbah NA, Karam NA, et al. (2013) CD14 tobacco gene-environment interaction in atopic children. *Cell Immunol* 285: 31-37.
53. Hussein YM, Alzahrani SS, Alharthi AA, Alhazmi AS, Ghonaim MM, et al. (2016) Gene Polymorphism of Interleukin-4, Interleukin-4 Receptor and STAT6 in Children with Atopic Dermatitis in Taif, Saudi Arabia. *Immunol Invest* 45: 223-234.
54. Zahran F, Hussein YM, Ashour E, Sayed M, Shalaby SM, et al. (2013) Interleukin-4 and interleukin-4 receptor alpha polymorphisms in atopic dermatitis: A case-control study. *BioChemistry: An Indian Journal* 7: 89-96.
55. Marinkovich VA (2001) Allergy and IgE antibodies. *N Engl J Med* 344: 1332-1333.
56. Hussein PY, Zahran F, Ashour Wahba A, Ahmad AS, Ibrahim MM, et al. (2010) Interleukin 10 receptor alpha subunit (IL-10RA) gene polymorphism and IL-10 serum levels in Egyptian atopic patients. *J Investig Allergol Clin Immunol* 20: 20-26.
57. Hussein YM, Zahran F, El-Zaher AA, El Tarhouny SA, Shalaby SM (2012) Interleukin-10 gene polymorphism and its blood level as biochemical markers among Egyptian atopic patients. *J Cell Sci Ther* 3: 8.
58. Hussein YM, El-Tarhouny SA, Shalaby SM, Mohamed RH, Hassan TH, et al. (2011) Interleukin-13 Receptor A1 Gene Polymorphism and IL-13 Serum Level in Atopic and Non-atopic Egyptian Children. *Immunological Investigations* 40: 523-534.
59. Aliprantis AO, Wang J, Fathman JW, Lemaire R, Dorfman DM, et al. (2007) Transcription factor T-bet regulates skin sclerosis through its function in innate immunity and via IL-13. *Proc Natl Acad Sci USA* 104: 2827-2830.
60. Bettiol J, Bartsch P, Louis R, De Groote D, Gevaerts Y, et al. (2000) Cytokine production from peripheral whole blood in atopic and nonatopic asthmatics: relationship with blood and sputum eosinophilia and serum IgE levels. *Allergy* 55: 1134-1141.