



Review

Implications of Allergic Reactions to Soybean Oil

San Miguel-Hernández A^{1*}, San Miguel-Rodríguez A¹, Martín-Armentia B¹, Martín-Armentia S², Armentia-Medina A³, Lozano-Estevan C⁴ and Iglesias-Peinado I⁵

¹Department of Clinical Analysis Service and Research Unit, Rio Hortega University Hospital, Valladolid, Spain

²Department of Pediatric Service, Centro Delicias II, Valladolid, Spain

³Department of Allergy, University Hospital Rio Hortega, Valladolid, Spain

⁴Department of Pharmacy, Universidad Alfonso X el Sabio, Spain

⁵Department of Pharmacy, Universidad Complutense, Spain

Abstract

Soybeans oil both cold-pressed and fully refined oils have been shown to contain proteins. Although most publications suggest that refined oils do not induce allergic reactions in sensitive individuals, adverse reaction to soy oil in an infant has been reported. Is used in cooking and food formulations. The presence of protein in soy oil depends on the degree of refining, as for other seed oils.

There seems to be agreement that fully refined soybean oils are not allergenic but our groups recently have found cases of anaphylaxis due to soybean oil presented in generic drugs and anaphylaxis during the night that was caused by an unexpected exposure to soybean oil included in their pillows. Skin prick tests and determination of specific Immunoglobulin IgE to soy were negative but molecular analysis by microarrays resolved the diagnosis.

Keywords: Allergic reactions; *Glycine max*; Soy bean; Soy bean oil

***Corresponding author:** San Miguel-Hernández A, Department of Clinical Analysis Service and Research Unit, Rio Hortega University Hospital, C/San Antonio de Padua 1, 1º Dcha. 47003, Valladolid, Spain, Tel: +34 620800117; E-mail: asanmi@saludcastillayleon.es

Citation: Miguel-Hernández AS, Miguel-Rodríguez AS, Martín-Armentia B, Martín-Armentia S, Armentia-Medina A, et al. (2018) Implications of Allergic Reactions to Soybean Oil. J Allergy Disord Ther 4: 008.

Received: October 3, 2018; **Accepted:** November 16, 2018; **Published:** December 3, 2018

Copyright: © 2018 San Miguel-Hernández A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

Soy (soybean) (*Glycine max*) is an edible legume belonging to the Fabaceae family. The seed contains around 20% oil and 38 to 40% protein. Consumption of soy, widespread in Asia and the US, has increased in Europe during the past years particularly [1,2].

More than 90% of food allergies are caused by eight foods: cow's milk, hen's egg, soy, peanuts, tree nuts, wheat, fish, and shellfish. The diagnostic workup for a patient with suspected food allergy includes a detailed medical history, physical examination, food allergy screening tests, and responses to an elimination diet and an oral food challenge. None of the screening tests, alone or in combination, can definitely diagnose or exclude a food allergy [3]. Novel microarrays diagnostic methods including those that focus on immune responses to specific food proteins or epitopes of specific proteins are under active study.

Soy bean is the world's most important legume. Protein and soy oils are widely used in animal and human nutrition, pharmaceutical industry and in different manufactures textiles [1,2]. Soybean oil is used in cooking and food formulations. The presence of protein in soy oil depends on the degree of refining, as for other seed oils.

There is a certain consensus that fully refined soybean oils are not allergenic but our groups recently have found cases of anaphylaxis due to soybean oil presented in generic drugs and anaphylaxis during the night that was caused by an unexpected exposure to soybean oil included in their pillows.

Clinical Properties of Soy Bean Oil

Soy is a leguminous but it is also an oilseed with a high fat content, 20% dry matter, the second highest content among all food legumes (the highest oil content is found in peanut, 48%; then chickpea, 5%, and other food legume species in the range of 1-3.6%) [4]. thus, the average composition of the soybean is (% dry weight): fat 21% (located mainly in the cotyledon and in the hypocotyls), protein 40.3%, carbohydrates 33.9% and ash 4.9% [5].

Table 1 shows the composition of crude and refined soybean oil. Soybean oil consist primarily of triglycerides (>99% in refined oils), some phospholipids and some minor compounds such as phytosterols and tocopherols. Triglycerides are formed mostly by three unsaturated fatty acids (65.3% of all TG, wt) or two unsaturated FA (31% wt) [6]. Thus, soy oil stands out by a low content in saturated fats (15.5%, mainly palmitic acid) and high in unsaturated fats (85.5%) (Particularly oleic acid, 21% and polyunsaturated acids, 58.5%). Among the polyunsaturated fats, soy oil is rich in n-6 like linoleic acid (53% in weight) and in n-3 like linolenic acid (8% in weight). There are some variations in the composition because of varietal and seed maturation temperature effects, and feed of animal (Table 2) [7].

Phospholipids like lecithin are present in crude soybean oil (1.5-2.5%) and are eliminated during oil refining (during degumming). In fact, lecithin is a co-product of soybean and corn oil processing (Table 1) [7].

The unsaponifiable fraction of soy oil is 0.3-1.6%. The main phytosterols (0.13-0.33%) in soy oil are β -sitosterol (1317 mg/kg), campesterol and stigmasterol (with around 50% of the β -sitosterol each) [8]. Phytosterols have healthy effects in the diet, decreasing cholesterol absorption in the intestine [9-12].

Soy oil has tocopherols, minor components with antioxidant activities, with the γ being in the highest concentrations, close to 1000 mg/kg (crude oil), followed by δ (around 400 mg/kg) and α (66.5-90.7 mg/kg, according to the variety) [5]. Tocopherols are decomposed during vegetable oil processing but a large proportion is left in the finished oils. The content of mixed tocopherols is quite high and range between 550 and 1000 μ g/g in Refined, Bleached and Deodorized (RBD) oils [12]. This value is above the optimum range of 400-600 μ g/g for antioxidant activity [10]. The tocopherol content in several oils, including soybean, may be adequate to protect the oils against oxidation under ambient conditions, while in others like olive and peanut oils the levels (110 and 200 μ g/g, respectively) are not sufficient [11].

Components	Crude oil	Refined oil
Triglycerides (%)	95-97	> 99
Phospholipids (%)	1.5-2.5	0.003-0.045
Unsaponifiable matter (%)	1.6	0.3
Phytosterols	0.33	0.13
Tocopherols	0.15-0.21	0.11-0.18
Hydrocarbons	0.014	0.01
Free fatty acids (%)	0.3-0.7	< 0.05
Trace metals		
Iron (ppm)	1-3	0.1-0.3
Copper (ppm)	0.03-0.05	0.02-0.06

Table 1: Average compositions for crude and refined soybean oil.

Adapted from Pryde [12].

Soy oil has a light yellow color and a soft flavor [8]. The fatty acid profile of soybean oil results in low oxidative stability that limits the uses of soybean oil in food products and industrial applications, and can be modified easily by genetic engineering, thus obtaining oil with a low content of saturated fats and high of unsaturated fats like oleic or linoleic acid, for example [9].

Soy oil has also a low stability to frying comparing to other edible fats. In addition, due to the presence of furanic acids, this oil oxidizes rapidly in the presence of light. For this reason, the storage in the dark is recommended [8].

Soy Bean Oil as Source of Allergens

Both cold-pressed and fully refined oils have been shown to contain proteins (0.35-0.78 mg/kg) [3]. Although there is a certain consensus that fully refined soybean oils are not allergenic and most publications suggest that refined oils do not induce allergic reactions in sensitive individuals, an adverse reaction to soy oil in an infant has been reported [13-15].

IgE-binding proteins with MWs of 53 and 58 kDa were identified in three unrefined soybean oils [16]. A 56 kDa allergenic protein was also found in cold-pressed and deodorised soybean oils, which was later identified as soybean β -amylase (7S globulin), together with the 20 kDa allergen Kunitz Trypsin Inhibitor (KTI) [17,18].

The protein profile of the cold-pressed soy oil was similar to that of soy flour, with seven bands in a wide molecular range (94-14 kDa) [19]. The soy lecithin seed maturation protein P34 from the 7S globulin fraction (35 kDa) and β -amylase (56 kDa) were identified.

Hypersensitivity Reactions to Generic Drug-Containing Soybean Oil

Over the last years the use of generic drugs has increased in Spain and in the European Union due to their minor economic cost. The main regulatory request to market these products is their equivalence to the original compounds.

	Soy (<i>Glycine max</i>)	Sunflower (<i>Helianthus annuus</i>)	Peanut (<i>Arachis hypogaea</i>)	Canola (<i>Brassica napus</i>)
Oil content of the seed (% in weight)	18-23	25-30	42-52	Aprox. 40
Solidification point (°C)	-8(-18)	-18(-20)	-2-(+3)	0(-2)
Saturated fatty acids:				
Palmitic acid (16:0)	10	6.5	10	4
Stearic acid (18:0)	5	5	3	1.5
Eicosanoic acid (20:0)	0.5	0.5	1.5	0.5
Docosanoic acid (22:0)	0	0	3	0
Unsaturated fatty acids:				
Oleic acid (18:1)	21	23	41	63
Linoleic acid (18:2, n-6)	53	63	35.5	20
Linolenic acid (18:3, n-3)	8	<0.5	0	9
Eicosenic acid (20:1) and Eicosadienoic acid (20:2)	3.5	1	1	1
Erucic acid (22:1)	0	0	-	0.5

Table 2: Composition in fatty acids (% in weight) and properties of some edible oils.

Adapted from Belitz [8].

Soy bean is an additive of some drugs. Hypersensitivity reactions to soy bean oil as a result of drug intake have been scarcely reported [20-22]. Recent studies on analytic investigation on protein content in refined seed oils suggest that fully refined seed oils should be taken into account in the context of allergic reactions and would benefit of further toxicological studies [22].

The observations suggest that an immediate hypersensitivity to drug-containing soy oil may cause anaphylaxis reaction in patients previously sensitized to this legume. The first report of possible anaphylaxis after a drug containing soybean oil implicated the drug propofol [20]. Since propofol contains both egg lecithin and soybean oil, its use is contraindicated in patients with hypersensitivities to these component. Several other drugs may have a food component, resulting in contraindications and warnings in product labelling.

Several commercially important refined vegetable oils are derived from plants which are recognized as potent food allergens (peanut, soy). Full refining of oils results in the almost complete removal from oils of protein, which is responsible for allergic reactions. However, it is uncertain whether the minute amounts remaining could provoke allergic reactions in highly susceptible individuals. This has led to a vigorous debate about the safety of refined oils and specifically whether to label each oil individually because of the potential risk of allergenicity [23-25].

Until active principles are clearly indicated in drug labellings, excipients and other minor additives are not included and only defined as excipients. Since the introduction of generic drugs to the pharmaceutical market a debate exists whether they are well-investigated and of high quality. There is some uncertainty about whether evidence of bioequivalence is enough to guarantee efficacy and safety of generic drugs [25]. Food allergy consumers depend on ingredient labels for allergen avoidance [23]. However the drug labelling may not indicate the form or source of the allergen, and individuals who currently avoids foods may presented severe allergic symptoms after intake of drugs with minimal protein content such as soy oil, soy lecithin or lysozyme [26]. There are now reliable assays for the determination of soybean proteins in processed foods that may be applicable in drugs [27, 28].

Over the last years the use of generic drugs has increased in the European Union due to their minor economic cost. The main regulatory request to market these products is their equivalence to the original compounds. We described two women that presented anaphylaxis after intake of generic omeprazole. The two patients had previously tolerated original non-generic omeprazole. Dot-blot assay revealed that the serum of the two patients was positive to generic omeprazole and soy. The diagnosis of soy allergy should not be excluded in cases of drug hypersensitivity. We suggest to tests soy in all hypersensitivity reaction to generic drugs [29].

Although incidences of anaphylactic reactions induced by proton pump inhibitors of H₂ are rare, they can life threatening. Skin prick tests and oral challenge tests may be useful for the diagnosis, but immunological test never confirm the presence of specific IgE antibodies to active principles of these drugs [23].

Soy is a clinical relevant allergenic source. The diagnosis of soy allergy should not be excluded in cases of drug hypersensitivity. We suggest to tests soy in all hypersensitivity reaction to generic drugs [28].

New studies described new protein involve in oil bodies fractions of several seed and nuts mainly, which are recognized by IgE antibodies from patients sensitized to these, and which cause allergic reactions ranging from oral allergy syndrome to anaphylaxis. These works used standard prick test and specific IgE determinations, which provided negative results. Although isolated reports have shown sensitization to the agent tested using prick-by-prick tests prepared with saline, in most cases specific techniques to conserve the lipid fractions of the extracts were required [29].

In addition, they have not only identified oleosins as sensitizing lipid fractions (fractions Ara h 10 and 11 of peanuts) but also establish that these are gastro-resistant allergens. This might explain the cases of anaphylaxis reported in the literature and the existence of cross-reactivity with other plant foods such as buckwheat, which has been tested by RAST inhibition techniques [30].

We hypothesize that negative results might be due to the lipid nature of the allergen in question and, therefore, the difficulties inherent in its processing. Interestingly, most allergenic extracts are obtained by water-soluble methods that could eliminate these lipids derivatives of the allergenic extract. The role of lipid derivatives in food allergies requires deeper investigation.

Soy Bean Oil as a Hidden Allergen: Anaphylaxis Due to Pillows

Protein and soy oils are widely used in animal and human nutrition, pharmaceutical industry and very recently in manufacturing of pillows. We studied cases of anaphylaxis during the night that was caused by an unexpected exposure to soy included in their pillows. The patients suffered anaphylaxis during their sleep, near daily. They presented nocturnal episode of dyspnoea, erythema and facial edema, scattered wheals, malaise, palpitations and intense palmar itching. Emergency treatment with adrenaline was necessary. The symptoms did not appear when the patients sleep in hotels or in other houses.

Today, pillows are stuffed primarily with materials such as polyester (a synthetic), feathers, down, or a combination of the latter two. There are recently made generations of pillows, soy-based. There are also internationally patented the *Aloe vera* injected in the pillow core, the use of natural soy oils in our compositions, and many other innovative products, most of them used in babies pillows.

The only common finding among these patients was that they have bought a viscoelastic pillow before their symptoms began. We obtained industrial information of soy-oil composition in the matrix of these pillows. Consequently, we performed a later allergic study. All patients presented Skin Prick Tests (SPT) pricks and IgE determination (ImmunoCAP assay, Phadia, Uppsala, Sweden) to conventional aeroallergens and foods (soy included) negative [29,30].

Component-resolved diagnosis and microarray technology have been recently introduced into clinical allergy practice, and may be particularly useful in food-sensitized allergic patients and in patient with anaphylaxis of hidden allergens. After informed consent, new SPT with nut, hull and oil soy and microarrays (ISAC®) were performed. The results of SPT to soy different nut extracts were negative in all patients. Only oil soy revealed a small wheal (12 mm², histamine 19.6 mm²). Nevertheless microarray-based IgE detection assay (ISAC®) revealed in all patients hypersensitivity to β-conglycinin (nGly m 5). Their symptoms disappear after elimination of their

pillows [29]. They eliminated all source of soy in their room (pillows) and the ingestion of any food that could be manufactured with this oil. They have been revised each six months during a period of 10 years without relapse.

Exposure to soy antigens has been associated with asthma in community outbreaks and in some workplaces. Recently, 135 Soy Flake Processing Workers (SPWs) in a Tennessee facility were evaluated for immune reactivity to soy. Allergic sensitization to soy was common and was five times more prevalent than in Health Care Worker controls (HCWs) with no known soy exposure. The prominent proteins that bound SPW IgE were identified by nanoUPLC MS/MS analysis to be the high molecular weight soybean storage proteins, β -conglycinin (Gly m 5), and Glycinin (Gly m 6) [31].

High molecular weight soybean storage allergens, Gly m 5 and Gly m 6, may be respiratory sensitizers in occupational exposed workers but in other patients no professionally exposed. Soy protein has shown great potential for use in biobased adhesives. β -Conglycinin is a major component of soy protein; it accounts for 30% of the total storage protein in soybean seeds. β -Conglycinin was isolated and purified, and its subunits' (β , α') physicochemical and adhesive properties were characterized [32].

Soy is also used in pharmaceutical products as soymorphins or drug-excipients [14,22]. Oral allergy syndrome to soy milk is classified as a phenotype of Pollen-Food Allergy Syndrome (PFAS). As causative antigens, Gly m 4 (Bet v1 homolog, 17 kD) and oleosin (23 kD), have been reported. Recently two cases of Pollen Food Allergy Syndrome (PFAS) to soy milk have been reported [15]. Both cases showed positive reactions to soy milk in Skin Prick Tests (SPT) and to Gly m 4 in specific serum immunoglobulin E antibody. When they measured specific serum IgE antibody of soy-related proteins using microarray analysis, both cases showed a positive reaction for Bet v1. Based on the array results, the authors diagnosed the two cases as PFAS to Gly m 4.

We performed protein microarray analysis and found it useful as a screening test for immediate allergy, such as PFAS. The reason of negative SPT and IgE to soy in the tests performed to our patients may be due to the acidic subunits of glycinin and β -conglycinin, major soybean storage proteins, appear to be absent or present in much reduced amounts in these techniques. Immunoblots with soy-allergic sera indicate alteration, reduction, or loss of IgE-binding in the commercial extracts as compared to extracts of soy flour. Preparation methods appear to be partially responsible for the variable allergen content in commercial soybean skin test extract [33].

In summary, soy is recognized as one of the "big 8" food allergens. IgE antibodies from soybean-sensitive patients recognize more than 15 soybean proteins [34]. Some of these food allergens can be found as a hidden allergen like pillow composition. Sensitization to the soybean allergens Gly m 5 or Gly m 6 is potentially indicative for severe allergic reactions to soy, like our patients suffered from [35-37].

Recent Advances in the Control of Allergic Reaction to Soybean

Recently, some global regulatory agencies started requiring quantitative analysis of individual allergens, including unproven allergens, as part of the risk assessment for genetically engineered soy beans

[38]. Identification of allergenic globulins as percutaneously sensitized soybean allergens have been demonstrated through epidermal application of crude soybean extract [39]. Glycinin and β -conglycinin can induce intestinal damage [40].

Measures as fermentation with microorganisms boiling and pressure cooking genetical modification and other measures can degrade major allergens and achieve a reduction of soybean hypersensitivity. Removal of soybean oil in intralipid for total parenteral nutrition have been also proposed [41-45].

Conclusion

Soy by itself is not usually an important food in the diet, however, it is used in many products. The elimination of all foods that include soy is essential for allergic patients. It is recognized as one of the major food allergens. IgE antibodies to soy appear in sensitive patients and have been recognized up to more than 15 soy proteins. Some of these food allergens can be found as a hidden allergen, as is the case with the composition of the pillows and their inclusion in certain pharmaceutical products. Sensitization to soy allergens Gly m 5 or Gly m 6 is potentially indicative of severe allergic reactions to soy.

References

- European Food Safety Authority (2014) Scientific Opinion on the evaluation of allergenic foods and food ingredients for labelling purposes. EFSA Journal 12: 3894.
- Wang T (2002) Soybean Oil. In: Gunstone FD (ed.). Vegetable Oils in Food Technology: Composition, Properties, and Uses. Blackwell, New Jersey, USA. Pg no: 337.
- Hidalgo FJ, Zamora R (2006) Peptides and proteins in edible oils: Stability, allergenicity, and new processing trends. Trends Food Sci Tech 17: 56-63.
- Pattee HE, Salunkhe DK, Sathe SK, Reddy NR (1983) Legume lipids. In: Arora SK (ed.). Chemistry and Biochemistry of Legumes: Edward Arnold. Experimental Agriculture 20: 95.
- Anwar F, Kamai GM, Nadeem F, and Shabir G (2015) Variations of quality characteristics among oils of different soybean varieties. Journal of King Saud University-Science 28: 332-338.
- Dergal SB (2006) [Química de los alimentos] (Food chemistry) (4thedn). Pearson Educacion, México.
- Lusas EW (2000) Oilsdeeds and oil-bearing materials. In: Kulp K, Ponte JG (eds.). Handbook of Cereal Science and Technology, Revised and Expanded. CRC Press, New York, USA.
- Belitz HD, Grosch W, Schieberle P (2009) Food Chemistry. Springer-Verlag Berlin Heidelberg, Germany.
- Clemente TE, Cahoon EB (2009) Soybean oil: genetic approaches for modification of functionality and total content. Plant Physiol 151: 1030-1040.
- Frankel EN, Cooney PM, Moser HA, Cowan JC, Evans CD (1959) Effect of antioxidants and metal in activators in tocopherol-free soybean oil. Eur J Lipid Sci Technol 10: 1036-1039.
- Frankel EN (1996) Antioxidants in lipid foods and their impact on food quality. Food Chem 57: 51-55.
- Pryde EH (1980) Composition of soybean oil. In: Erickson DR, Pryde EH, Brekke OL (eds.). Handbook of Soy Oil Processing and Utilization. AGRIIS, Rome, Italy.

13. Taylor SL, Baumert JL, Zruizinga AG, Remington BC, Crevel RW, et al. (2014) Establishment of Reference Doses for residues of allergenic foods: report of the VITAL Expert Panel. *Food Chem Toxicol* 63: 9-17.
14. Crevel RW, Baumert JL, Baka A, Houben GF, Knulst AC, et al. (2014) Development and evolution of risk assessment for food allergens. *Food Chem Toxicol* 67: 262-276.
15. Moneret-Vautrin DA, Morisset M, Flabbee J, Kanny G, Kirch F, et al. (2002) Unusual soy oil allergy. *Allergy* 57: 266-267.
16. Paschke A, Zunker K, Wigotzki M, Steinhart H (2001) Determination of the IgE-binding activity of soy lecithin and refined and non-refined soybean oils. *J Chromatogr B Biomed Sci Appl* 756: 249-254.
17. Errahali YJ, Thomas LD, Keller TC, Lee HJ (2013) Inhibition by New Glucocorticoid Antedrugs [16 α , 17 α -d] Isoxazoline and [16 α , 17 α -d]-3'-Hydroxy-Iminoformyl Isoxazoline Derivatives of Chemotaxis and CCL26, CCL11, IL-8, and RANTES Secretion. *J Interferon Cytokine Res* 33: 493-507.
18. Errahali Y, Morisset M, Moneret-Vautrin DA, Kanny G, et al. (2002) Allergen in soy oils. *Allergy* 57: 648-649.
19. Martin-Hernandez C, Bénet S, Obert L (2008) Determination of Proteins in Refined and Nonrefined Oils. *J Agric Food Chem* 56: 4348-4351.
20. Hofer KN, McCarthy MW, Buck ML, Hendrick AE (2003) Possible anaphylaxis after propofol in a child with food allergy. *Ann Pharmacother* 37: 398-401.
21. Frémont S, Errahali Y, Bignol M, Metche M, Nicolas JP (2002) [Allergenicity of oils]. *Allerg Immunol (Paris)* 34: 91-94.
22. Ramazzotti M, Mulinacci N, Pazzagli L, Moriondo M, Manao G, et al. (2008) Analytic investigations on protein content in refined seed oils: implications in food allergy. *Food Chem Toxicol* 46: 3383-3388.
23. Stefanaki ECh, Vovolis V, Letsa I, Koutsostathis N (2008) Anaphylactic reaction to Omeprazole. *Am J Gastroenterol* 103: 1581-1583.
24. Crevel RW, Kerkhoff MA, Koning MM (2000) Allergenicity of refined vegetable oils. *Food Chem Toxicol* 38: 385-393.
25. Simone E, Weiis CC, Furlong TJ, Sicherer SH (2005) Impact of ingredient labelling practices on food allergic consumers. *Ann Allergy* 95: 426-428.
26. Tschibitscher D, Platzer P, Baumgärtel C, Müllner M (2008) [Generic drugs: quality, efficacy, safety and interchangeability]. *Wien Klin Wochenschr* 120: 63-69.
27. Artesani MC, Donnanno S, Cavagni G, Calzone L, D'Urbano L (2008) Egg sensitization caused by immediate hypersensitivity reaction to drug-containing lysozyme. *Ann Allergy* 101: 105.
28. Morishita N, Kamiya K, Matsumoto T, Sakai S, Teshima R, et al. (2008) Reliable enzyme-linked immunosorbent assay for the determination of soybean proteins in processed foods. *J Agric Food Chem* 56: 6818-6824.
29. Dueñas-Laita A, Pineda F, Armentia A (2009) Hypersensitivity reactions to generic drug containing soybean oil. *N Engl J Med* 361: 1317-1318.
30. Armentia A, Pineda F, Martín B, San Miguel A, Martín Gil FJ, et al. (2013) Anaphylaxis caused by hidden soybean allergens in pillows. *J Allergy Clin Immunol* 131: 228-230.
31. Green BJ, Cummings KJ, Rittenour WR, Hettick JM, Bledsoe TA, et al. (2011) Occupational sensitization to soy allergens in workers at a processing facility. *Clin Exp Allergy* 41: 1022-1030.
32. Mo X, Wang D, Sun XS (2011) Physicochemical Properties of β and $\alpha'\alpha$ Subunits Isolated from Soybean β -Conglycinin. *J Agric Food Chem* 59: 1217-1222.
33. Kaneko K, Iwasaki M, Yoshikawa M, Ohnata K (2010) Orally administered soymorphins, soy-derived opioid peptides, suppress feeding and intestinal transit via gut mu(1)-receptor coupled to 5-HT(1A), D(2), and GABA(B) systems. *Am J Physiol Gastrointest Liver Physiol* 299: G799-805.
34. Yagami A, Inaba Y, Kuno Y, Suzuki K, Tanaka A, et al. (2009) Two cases of pollen-food allergy syndrome to soy milk diagnosed by skin prick test, specific serum immunoglobulin E and microarray analysis. *J Dermatol* 36: 50-55.
35. Krishnan HB, Kim WS, Jang S, Kerley MS (2009) All three subunits of soybean beta-conglycinin are potential food allergens. *J Agric Food Chem* 57: 938-943.
36. Holzhauser T, Wackermann O, Ballmer-Weber BK, Bindslev-Jensen C, Scibilia J, et al. (2009) Soybean (*Glycine max*) allergy in Europe: Gly m 5 (beta-conglycinin) and Gly m 6 (glycinin) are potential diagnostic markers for severe allergic reactions to soy. *J Allergy Clin Immunol* 123: 452-458.
37. Herian AM, Bush RK, Taylor SL (1992) Protein and allergen content of commercial skin test extracts for soybeans. *Clin Exp Allergy* 22: 461-468.
38. Lu M, Jin Y, Cerny R, Bllmer-Weber B, Goodman RE (2018) Combining 2-DE immunoblots and mass spectrometry to identify putative soybean (*Glycine max*) allergens. *Food Chem Toxicol* 116: 207-215.
39. Murakami H, Ogawa T, Takafuta A, Yano E, Zaima N, et al. (2018) Identification of the 7S and 11 S globulins as percutaneously sensitizing soybean allergens as demonstrated through epidermal application of crude soybean extract. *Biosci Biotechnol Biochem* 82: 1408-1416.
40. Peng C, Cao C, He M, Shu Y, Tang X, et al. (2018) Soybean Glycinin- and β -Conglycinin-Induced Intestinal Damage in Piglets via the p38/JNK/NF- κ B Signaling Pathway. *J Agric Food Chem* 66: 9534-9541.
41. Yang A, Zuo L, Cheng Y, Wu Z, Li X, et al. (2018) Degradation of major allergens and allergenicity reduction of soybean meal through solid-state fermentation with microorganisms. *Food Funct* 9: 1899-1909.
42. Cabanillas B, Cuadrado C, Rodriguez J, Dieguez MC, Crespo JF, et al. (2018) Boiling and Pressure Cooking Impact on IgE Reactivity of Soybean Allergens. *Int Arch Allergy Immunol* 175: 36-43.
43. Lu M, Jin Y, Ballmer-Weber B, Goodman RE (2018) A comparative study of human IgE binding to proteins of a genetically modified (GM) soybean and six non-GM soybeans grown in multiple locations. *Food Chem Toxicol* 112: 216-223.
44. Selb R, Wal JM, Moreno FJ, Lovik M, Mills C, et al. (2017) Assessment of endogenous allergenicity of genetically modified plants exemplified by soybean - Where do we stand? *Food Chem Toxicol* 101: 139-148.
45. Nicklas RA (2013) Lack of allergenic soy in intralipid for total parenteral nutrition. *Ann Allergy Asthma Immunol* 111: 423.



- Journal of Anesthesia & Clinical Care
Journal of Addiction & Addictive Disorders
Advances in Microbiology Research
Advances in Industrial Biotechnology
Journal of Agronomy & Agricultural Science
Journal of AIDS Clinical Research & STDs
Journal of Alcoholism, Drug Abuse & Substance Dependence
Journal of Allergy Disorders & Therapy
Journal of Alternative, Complementary & Integrative Medicine
Journal of Alzheimer's & Neurodegenerative Diseases
Journal of Angiology & Vascular Surgery
Journal of Animal Research & Veterinary Science
Archives of Zoological Studies
Archives of Urology
Journal of Atmospheric & Earth-Sciences
Journal of Aquaculture & Fisheries
Journal of Biotech Research & Biochemistry
Journal of Brain & Neuroscience Research
Journal of Cancer Biology & Treatment
Journal of Cardiology & Neurocardiovascular Diseases
Journal of Cell Biology & Cell Metabolism
Journal of Clinical Dermatology & Therapy
Journal of Clinical Immunology & Immunotherapy
Journal of Clinical Studies & Medical Case Reports
Journal of Community Medicine & Public Health Care
Current Trends: Medical & Biological Engineering
Journal of Cytology & Tissue Biology
Journal of Dentistry: Oral Health & Cosmesis
Journal of Diabetes & Metabolic Disorders
Journal of Dairy Research & Technology
Journal of Emergency Medicine Trauma & Surgical Care
Journal of Environmental Science: Current Research
Journal of Food Science & Nutrition
Journal of Forensic, Legal & Investigative Sciences
Journal of Gastroenterology & Hepatology Research
Journal of Gerontology & Geriatric Medicine
Journal of Genetics & Genomic Sciences
Journal of Hematology, Blood Transfusion & Disorders
Journal of Human Endocrinology
Journal of Hospice & Palliative Medical Care
Journal of Internal Medicine & Primary Healthcare
Journal of Infectious & Non Infectious Diseases
Journal of Light & Laser: Current Trends
Journal of Modern Chemical Sciences
Journal of Medicine: Study & Research
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology
Journal of Neonatology & Clinical Pediatrics
Journal of Nephrology & Renal Therapy
Journal of Non Invasive Vascular Investigation
Journal of Nuclear Medicine, Radiology & Radiation Therapy
Journal of Obesity & Weight Loss
Journal of Orthopedic Research & Physiotherapy
Journal of Otolaryngology, Head & Neck Surgery
Journal of Protein Research & Bioinformatics
Journal of Pathology Clinical & Medical Research
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance
Journal of Physical Medicine, Rehabilitation & Disabilities
Journal of Plant Science: Current Research
Journal of Psychiatry, Depression & Anxiety
Journal of Pulmonary Medicine & Respiratory Research
Journal of Practical & Professional Nursing
Journal of Reproductive Medicine, Gynaecology & Obstetrics
Journal of Stem Cells Research, Development & Therapy
Journal of Surgery: Current Trends & Innovations
Journal of Toxicology: Current Research
Journal of Translational Science and Research
Trends in Anatomy & Physiology
Journal of Vaccines Research & Vaccination
Journal of Virology & Antivirals

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>