

## Research Article

# Envenomings Caused by Venomous Animals in Roraima: A Neglected Health Problem in the Brazil's Northernmost State

Wállex Matias Pedroso Souza<sup>1</sup>, Gabriel Alexandre Silva<sup>1</sup>, Felipe Augusto Cerni<sup>2</sup>, Isadora Sousa de Oliveira<sup>2</sup>, Umberto Zottich<sup>1</sup>, Bruna Kempfer Bassoli<sup>1</sup> and Manuela Berto Pucca<sup>1\*</sup>

<sup>1</sup>Medical School, Federal University of Roraima, Boa Vista, Brazil

<sup>2</sup>Department of Physics and Chemistry, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, Brazil

### Abstract

Accidents caused by venomous animals are an important global neglected disease with high impact in Brazilian Amazon. The substantial numbers of envenomings registered in the Amazon region can be explained by the optimal conditions for the venomous fauna (e.g climate), however, the numbers are underestimated owing to extensive under-reporting. This study carried out an epidemiological study of envenomings caused by venomous animals in the state of Roraima, the Brazil's Northernmost state within a typical vegetation of Amazon rainforest. Envenomings by venomous animals between 2013 and 2016 were collected from 15 municipalities of Roraima and statistically analyzed using different parameters. Data were collected from the National Notifiable Diseases Information System (SINAN). During the studied period, Roraima presented 2,239 envenomings within venomous animals, with most of them registered in the capital Boa Vista. In addition, male victims between 20 and 39-years old prevailed. Regarding the length of time between the accident and therapy (i.e antivenom administration), 72.7% of the victims were treated in less than 6 hours. Fortunately, most of the Roraima envenomings were classified as mild, with exception to snakebites, in which victims were associated to present moderate envenomings, although deaths were considered rare. The high incidence of envenomings caused by venomous animals in Roraima, as well as the induced-moderate severity in victims of snakebites, contribute to the knowledge of the local reality, which is needed to advocate measures that aim to reduce the accidents by venomous animal attacks

\*Corresponding author: Manuela Berto Pucca, Medical School, Federal University of Roraima, Boa Vista, Brazil, Tel: +55 4581924482; E-mail: manu.pucca@ufr.br

**Citation:** Souza WMP, Silva GA, Cerni FA, de Oliveira IS, Zottich U, et al. (2019) Envenomings Caused by Venomous Animals in Roraima: A Neglected Health Problem in the Brazil's Northernmost State. J Toxicol Cur Res 3: 011.

**Received:** July 14, 2019; **Accepted:** July 29, 2019; **Published:** August 07, 2019

**Copyright:** © 2019 Souza WMP, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and promote therapeutic measures, such as supplying of specific antivenoms in places where they are most required.

**Keywords:** Amazon region; Epidemiological report; Envenoming; Roraima; Venomous animals

### Introduction

Accidents caused by venomous animals are considered a global public health problem due to high number of cases and the complexity of their clinical evolution [1]. Envenomings caused by these animals are classified as Neglected Tropical Diseases (NTD), since they affect almost exclusively low-income people, deprived of political power, and living in developing regions [2-4]. Snakebites leads the severity of envenomings, being responsible for most of disabilities and premature deaths worldwide. According to the World Health Organization (WHO), it is estimated that 7,400 people every day are bitten by snakes, resulting in 2.7 million cases of snakebite envenomings in the continent and about 81,000 to 138,000 deaths each year [4].

In Brazil, despite the magnitude regarding the incidence of the venomous animals' accidents, there is no uniformity of them through the regions [5]. While North and Central-West regions present snakebite prevalence, in the Northeast and Southeast regions scorpion stings prevailed [6,7]. On the other hand, in the South region, spiders are responsible for the highest number of accidents [5]. These geographic variations reflect the different degrees of human occupation in each region, the ecological imbalance caused by disordered urban growth, and the particular habitat that the animal can be adapted (e.g *Tityus serrulatus* scorpion demonstrated easy adaptation with high proliferation in most of Brazilian urban areas) [8-10]. It is important to note that only after June 1985 (during a high crisis and low production of antivenoms in the country), the notification of accidents with snakes became mandatory in the country. Later (1988), reporting scorpionism and araneism also became mandatory, which results in the National Program for Control Accidents by Venomous Animals (*Programa Nacional de Controle de Acidentes por Animais Peçonhentos*) [11].

The state of Roraima is the Brazil's Northernmost state, presenting a typical vegetation of Amazon rainforest, besides being able to encompass favorable climatic conditions (equatorial and tropical) to shelter a wide biodiversity in almost all its territory, similar to the other Brazilian states of the North region [5,12,13]. Together, all these conditions possibly justify why the Amazon region presents the highest incidence of snakebites. Interestingly, the Amazon stable temperature during all the year (average of 27°C) present relevant effect upon the reproduction of reptilians based on the inability of keeping themselves warm [14,15].

Despite the long history and high prevalence of accidents with venomous animals in Brazil, there are still few epidemiological studies regarding envenomings in the country [14,16], especially in the state of Roraima. Although underreported accidents are a problem in all Brazil, Roraima can be highlighted due to the scarcity of

research investments and high number of Indians (more than 40% of the state is considered Indian areas), which potentiate underreporting data. Thus, the real statistics that provide a complete understanding of the current picture of the venomous animals' accidents in Roraima is inconsistent and underestimated. In this sense, the present study describes an epidemiological analysis of venomous animals' accidents occurred in the Roraima state.

## Materials and Methods

The epidemiological data on envenomings for the period from 2013 to 2016 were obtained from National Notifiable Diseases Information System (SINAN), a public information system, available online by the Ministry of Health (see <http://portalsinan.saude.gov.br/>). Different variations were analyzed including year of incidence (2013, 2014, 2015, and 2016), municipality's incidence (Alto Alegre, Amajari, Boa Vista, Bonfim, Cantá, Caracarái, Caroebe, Iracema, Mucajaí, Normandia, Pacaraima, Rorainópolis, São João da Baliza, São Luiz and Uiramutã), age group of victims, sex of victims, venomous animal responsible for the accident (snake, spider, scorpion, caterpillar, bees, others, and unknown), number of snakebite accidents according to the genus (*Bothrops*, *Crotalus*, *Micrurus*, *Lachesis*, non-venomous snake and unknown), interval between accident and therapeutics (0-6 hours, 6-24 hours, and unknown), and envenoming severity (healing and deaths).

Data was obtained using Tabwin32 (version 4.14) and statistical analyzes were performed to verify the normal distribution (Shapiro Wilk's test) and homogeneity (Levene and Brown-Forsythe test) of the results, followed by parametric tests (in the presence of these conditions) or non-parametric tests (in the absence of these conditions).

Comparison between the different victims' age groups of accidents caused by venomous animals, different species of animals responsible for the accidents, and number of accidents caused by the different snake genus were performed by One-way of Analysis of Variance (ANOVA) followed by Tukey or Kruskal-Wallis test. The difference between the occurrence of accidents by venomous animals between the female and male patients was evaluated by Student's t-test for independent samples. Possible association between the type of accident and the time for the therapeutic intervention, as well as between the type of accident and the severity of the envenoming was evaluated by the Chi-Square or Fisher's Exact test. Statistical analysis was performed using the Excel Software (Microsoft Office 2013, Redmond, WA, USA), Graph Pad Prism 5.0 (La Jolla, San Diego, CA, USA), or Statistica 12.0 software (StatSoft). Statistical significance level was set at 5% ( $p < 0.05$ ).

## Results

During 2013 to 2016, a total of 2,239 cases of accidents involving venomous animals was reported in the state of Roraima. The 2014 year showed the highest number of accidents (589) followed by 2016, 2013, and 2015 with 583, 546, and 521 reported cases, respectively (Figure 1).

Concerning Roraima municipalities, the highest accidents numbers were registered in Boa Vista (state capital), which presented 1,048 victims through all the analyzed period, corresponding to 46.8% of the total number of accidents among the 15 municipalities analyzed (Table 1). Other municipalities also presented relevant number of envenomings during the four years analyzed, however presenting lower

victims than the state capital. Alto Alegre was the second city with highest number of victims during the analyzed period, with a total of 434 cases, corresponding to 19.38%. Amajari (110 cases), Mucajaí (100 cases), Uiramutã (96 cases), and Rorainópolis (77 cases), also presented important relevance. On the other hand, Cantá, the municipality with the lowest envenoming victims, only held 5 cases during 2013. Interestingly, during the years of 2014, 2015, and 2016, Cantá did not registered any envenoming. The analysis of the victims' ages reveals that victims with 20 to 39-years old present the highest envenoming numbers (2013-192, 2014-186, 2015-199, and 2016-182 victims), followed by victims with 40 to 59 (2013-108, 2014-124, 2015-61, and 2016-118 cases), and with 15 to 19-years old (2013-65, 2014-77, 2015-66 and 2016-75 cases). Elderly group (*i.e* more than 70 years-old) presented fewest envenoming victims (Figure 2).





Figure 1: Envenomings in Roraima (2013-2016). This figure is in color in the electronic version.


Table 1: Distribution of envenomings by Roraima municipalities (2013-2016).

Roraima Municipalities	2013	2014	2015	2016	Total
Alto Alegre	95	108	115	116	434
Amajari	29	18	35	28	110
Boa Vista	250	262	263	273	1,048
Bonfim	10	21	4	9	44
Cantá	5	-	-	-	5
Caracarái	19	21	11	12	63
Caroebe	15	16	10	12	53
Iracema	11	17	11	12	51
Mucajaí	20	23	17	40	100
Normandia	6	3	3	6	18
Pacaraima	22	13	15	13	63
Rorainópolis	22	35	5	15	77
São João da Baliza	18	13	8	10	49
São Luiz	7	7	8	6	28
Uiramutã	17	32	16	31	96
<b>TOTAL</b>	<b>546</b>	<b>589</b>	<b>521</b>	<b>583</b>	<b>2,239</b>

Regarding victim's gender, there was a significant predominance ( $p < 0.05$ ) of men in all the analyzed years, presenting at least two-thirds of the total of the victims (2013-377 versus 169; 2014-377 versus 212; 2015-338 versus 183, and 2016-384 versus 198 envenomings case, male and female, respectively) (Figure 3).




**Figure 2:** Roraima age-related number of envenomings (2013-2016). Data was analyzed by Kruskal-Wallis test. Values of  $p < 0.05$  were considered statistically significant. Age range 15-19 was statistically significant with 80 or more (#). Age range 20-39 was statistically significant with 65-69, 70-79 and 80 or more (\*). Age range 40-59 was statistically significant with 70-79 and 80 or more (&). This figure is in color in the electronic version.



**Figure 3:** Roraima sex-related number of envenomings (2013-2016). Sex-related (male versus female) number of accidents were analyzed by Student's t-test. Values of  $p < 0.05$  were considered statistically significant (\*). ND means not determined. This figure is in color in the electronic version.

Regarding the animals responsible for envenomings there were a significant predominance ( $p < 0.05$ ) of snakebites (2013-351, 2014-332, 2015-331, and 2016-326 cases) in comparison to other venomous animals. On the other hand, accidents triggered by scorpions appear in second position with more modest numbers than snakes (2013-62, 2014-80, 2015-81, and 2016-102 cases), followed by bees (2013-52, 2014-66, 2015-48, and 2016-81 cases), spiders (2013-22, 2014-27, 2015-23, and 2016-30 cases), and caterpillars (2013-9, 2014-11, 2015-4, and 2016-13 cases) (Table 2). In respect to snakes genera,

*Bothrops* genus presented the highest number of envenomings (2013-247, 2014-233, 2015-119, and 2016-233 cases), followed by *Crotalus* (2013-34, 2014-22, 2015-26 and 2016-326 cases), *Lachesis* (2013-33, 2014-29, 2015-15 and 2016-22 cases), and *Micrurus* genera (2013-1, 2014-3, 2015-2 and 2016-4 cases) (Figure 4). Nevertheless, the statistical significance ( $p < 0.05$ ) was only observed between *Bothrops*, *Micrurus*, and non-venomous snakes. It is important to emphasize that in a high number of accidents the snake genus was not determined (unknown).



**Figure 4:** Total number of snakebites in Roraima according to the animal genus (2013-2016). Data was submitted to analysis of Kruskal-Wallis test. \*indicates statistical difference ( $p < 0.05$ ) among them. ND means not determined. NV means non-venomous. This figure is in color in the electronic version.

The time between accident and medical care (antivenom therapy) was also evaluated. After excluding cases in which the venomous animal was not identified, a total of 1,319 (72.71%) victims were treated in less than 6 hours after the accident, meanwhile low number of victims were treated with more than 6 hours (21.3%) (Table 3). Interestingly, the statistical analysis demonstrated that ophidian accidents are associated with 78% of patients that received therapeutic intervention after 6 hours of the accident. On the other hand, accidents caused by spiders are associated with 6.2% of patients that received therapeutic intervention after 6 hours of the accident.

The victims' envenoming severity was also explored in this study. The analysis of all the reported cases (including those 198 cases where the severity was not stratified) reveals that 1,280 envenomings (57.1%) were classified as mild, 597 (26.6%) as moderate, and 164 (7.3%) as severe (Table 4). Notably, ophidian envenomings are high associated to the development of moderate pathological manifestations in victims, while other animals are associated to progress as mild envenomings ( $p < 0.05$ ).

**Table 2: Number of envenomings in Roraima according to the different venomous animals (2013-2016).**

Accidentyear	Snake	Spider	Scorpion	Caterpillar	Bees	Others	ND	Total
2013	351	22	62	9	52	36	14	546
2014	332	27	80	11	66	53	20	589
2015	331	23	81	4	48	32	2	521
2016	326	30	102	13	81	23	8	583
<b>Total</b>	<b>1,340<sup>a</sup></b>	<b>102<sup>b</sup></b>	<b>325<sup>c</sup></b>	<b>37<sup>b</sup></b>	<b>247<sup>c</sup></b>	<b>144<sup>b</sup></b>	<b>44</b>	<b>2,239</b>

Data was analyzed by ANOVA followed by Tukey's post hoc test. Values of  $p < 0.05$  were considered statistically significant. Different superscript letters indicate statistical difference. ND means not determined.

**Table 3: Number of envenomings in Roraima concerning therapeutical intervention time (2013-2016).**

Venomous Animal	0-6 Hours	6-24+ Hours	ND	Total
Snake	775	373	194	1,342 <sup>a</sup>
Spider	52	30	20	102 <sup>a</sup>
Scorpion	232	26	67	325 <sup>b</sup>
Caterpillar	24	2	11	37 <sup>b</sup>
Bee	147	28	72	247 <sup>b</sup>
Others	89	17	37	143 <sup>b</sup>
Unknown	17	2	24	43
<b>Total</b>	<b>1,336</b>	<b>478</b>	<b>425</b>	<b>2,239</b>

Data was analyzed by Chi-square test. Values of  $p < 0.05$  were considered statistically significant. Different superscript letters indicate statistical difference. *ND* means not determined.

**Table 4: Roraima's envenoming clinical evaluation according to animal's group.**

Venomous Animal	Mild	Moderate	Severe	ND	Total
Snake	576	510*	157*	99	1342
Spider	76	14	2	10	102
Scorpion	272	26	2	25	325
Caterpillar	29	2	-	6	37
Bee	205	27*	-	15	247
Others	110	16*	2	14	144
Unknown	12	2	1	29	44
<b>Total</b>	<b>1,280</b>	<b>597</b>	<b>164</b>	<b>198</b>	<b>2,239</b>


Data was analyzed by Chi-square test.

\*Indicates statistical difference ( $p < 0.05$ ) in comparison to the same venomous animal in the mild group. *ND* means not determined.

Clinical severity of envenoming (mild, moderate, or severe) was classified according to the Brazilian Health Ministry guidelines.

This peculiar distribution might be better observed once the cases are described as percentages, in which mild accidents represent 42.9% of the ophidian accidents. In addition, besides snakes present the highest numbers of moderate envenomings, they are also responsible to the most severe ones. For instance, the evaluation regarding type of accident and envenoming severity demonstrated that only ophidian accidents present a distinctive pattern of clinical evolution comparing to all analyzed peers. Meanwhile the specific statistical study demonstrated that, in exception to snakes, the other envenomings are prone to affect Roraima victims under the same clinical evolution (mild) since mild envenomings prevailed in spiders (74.5%), scorpions (83.6%), caterpillars (78.3%), and bees (82.9%).

Finally, the study of envenoming evolution, that is cure or death, reveals that 84.7% of victims evolved to cure, although 14.7% did not have their evolution registered (Figure 5A). Despite the high number of accidents in the state of Roraima, evolution to deaths is rare, and in the analyzed period there were only 9 deaths reported (2013-2, 2014-2, 2015-1, and 2016-4 deaths). Remarkably, 8 of the registered deaths were caused by snakebites (6 caused by *Bothrops*, 1 by *Crotalus*, and 1 by *Micrurus*) and one by another venomous animal genus (not mentioned in the database) (Figure 5B). Therefore, the presented panel suggests that, despite the accidents severity, time intervention of treatment is capable of preventing death in most of the envenomings in Roraima, leading to a very low lethality rate (0.004%).



**Figure 5:** (A) Roraima envenoming evolution (cure versus deaths) (2013-2016); (B) Snakebite lethality distribution according to snake genus (2013-2016). *ND* means not determined. This figure is in color in the electronic version.

## Discussion

Epidemiological data on envenomings within venomous animals are critical to reflect the real magnitude of these accidents. In Brazil, databases such as SINAN and National Poisoning Information System (*SINITOX, Sistema Nacional de Informações Tóxico-Farmacológicas*) are considered fundamental for the development of public health in the country, based on their analyzes it is possible to identify and quantify the antivenoms to be distributed into federated units. In other words, epidemiological studies can improve and strengthen production, supply, and distribution of life-saving antivenoms and other commodities in the country. Furthermore, based on the collected

data, government can determine strategic areas to be supervised, monitor and improve victims assistance in health units, as well as develop strategies to control envenoming accidents such as by prevention campaigns [17-20].

The present study analyzed epidemiological information on envenomings caused by venomous animals in Roraima state, the northernmost state of Brazil, with most of its area comprised of exuberant Amazonian rainforest sheltering ample biodiversity in almost all of its territory, similar to other Brazilian states of the North region [12,21]. Therefore, these factors express a great risk for the occurrence of envenomings, which possibly justifies the Amazon region having the highest incidence of ophidian accidents [14].

The number of envenomings caused by venomous animals in Roraima during the period 2013-2016 is low (less than 0.33%) compared to the numbers registered in Brazil (2013-159-597, 2014-168-399, 2015-172-014, and 2016-172-412) [22]. It is interesting to mention that in 2015, in the state of Roraima, there was a decrease in reported accidents. This occurrence could be a reflection of national public strategies to control these accidents such as:

- i. Campaign to aware the population to avoid certain environments during rainy season, a season where snakes and other venomous animals usually leave their nests and invade urban areas [23-25];
- ii. Propagation of the use of PPE (Personal Protection Equipment) to workers in specific areas, such as forests, open field, civil construction, and many others [26,27];
- iii. Underreporting, incomplete registration, or omissions in data collection [16,18].

The distribution of accidents among municipalities reveals the highest number of notifications in Boa Vista, which was expected, since the capital presents most of the population of Roraima (63.5%), as well as concentrates the main health reference centers such as the General Hospital of Roraima (HGR, *Hospital Geral de Roraima Rubens de Souza Bento*) and the Children's Emergency Service (HCSA, *Hospital da Criança Santo Antonio*), besides the capital presenting the most health professionals trained to effectively report envenomings accidents compared to smaller municipalities. However, other municipalities such as Amajari, Mucajaí, Uiramutã, and Rorainópolis also present relevant contribution. These municipalities are located in forest boundaries, which could explain the high incidence of envenoming within venomous animals [21,28,29].

Regarding gender, it was reported highest number of envenomings with men in all age groups, especially males between 20 and 39 years old. This immediately suggests that accidents occur in economically active individuals and outside the home environment, probably due to working conditions that expose them to such risk (such as farming, hunting, etc) [30], which was previously observed [31-33]. Furthermore, this finding is in agreement with the national data [18,34].

Looking into the animal genera responsible for envenomings in Roraima, snakes predominated with nearly 60% of all accidents, which were especially caused by *Bothrops* (popular name *jararaca*) and *Crotalus* (rattlesnake, popular name *cascavél*) genera, which results can be supported by other studies [35,36]. The snakebite predominance can be justified by the fact that snakes have considerable preference for environments that are prone to rodent proliferation (*e.g* storehouses, barns, and rubble), located inside forests or surrounding forests, a very common places in the municipalities of Roraima

[21,37-40]. Regarding rattlesnakes envenomings in the state, it is important to emphasize the *Crotalus durissus ruruima* species, which in Brazil can be found just in Roraima (although they habit southern Venezuela) [41]. Besides presenting particular components and high individual variation [42], the main problem is that *C.d. ruruima* venom is not used to produce anticrotalic antivenom in Brazil (horses are immunized with *C.d. terrificus* and *C.d. collilineatus* venoms, 50% of each) [43,44]. Indeed, antivenomic studies demonstrated that *C.d. ruruima* venom is not well recognized and neutralized by the Brazilian rattlesnake antivenom, recommending the use of botrocrotalic antivenom for therapeutics of victims [41].

The scorpion envenomings reach the second position, being responsible for 14.51% of the total number of accidents in Roraima. Our findings differ from the epidemiological data from most of Brazilian states, such as São Paulo and Minas Gerais, where scorpionism leads [17,45,46]. One possible justification is the absence of *Tityus serrulatus* scorpion in Roraima, which is the species mainly responsible for the high scorpionism incidence in the country. In addition, Roraima presents low urbanized areas (*i.e* less populated state of Brazil), which results to lowland areas favorable to scorpion proliferation due to the low accumulation of debris, garbage, and insects [47].

During 2013 to 2016, 50.95% of the victims were treated in less than 6 hours after the accident, which suggest a positive scenario since only 7.42% of the envenomings were considered severe. Clinical severity of envenoming (mild, moderate, or severe) was classified according to the Brazilian Health Ministry guidelines [48]. Other study also analyzed the time interval in receiving assistance in Rondônia (other Amazon state), in which the time between the snakebite and arrival at the hospital was higher than 6 hours in 72.7% of patients [49]. According to studies conducted in the last 100 years in Brazil, the main challenge of treating envenoming patients is the delayed care, which will result in severe envenoming and difficulties to control venom effects into patient efficiently [18,50,51]. Thus, for the best prognosis and reduction of venom induced-sequelae, it is recommended that the victims are treated in the first 6 hours after the accident [50], which emphasizes that in Roraima state the envenomings are, in most of cases, being treated with less time delay. Notably, scorpions and caterpillars envenomings resulted in the earliest care (less than 1 hour), possibly because most of them occurred in urban areas, where access to treatment can be established earlier [28,52,53].

Although most of snakebites data were registered in Boa Vista, several happened in peripheric and/or rural areas, nearby cities, or even in Indian communities. Thus, a delay in the treatment is expected based on the distance from origin of the accident and the hospital. Moreover, Roraima habitants keep some traditional culture to treat envenomings and, in many cases, victims only look for properly medical care after using traditional medicines instead [54]. Indeed, a recent study demonstrated that only few native Indians were registered as having received treatment in Brazilian Amazon (6.88%) [33].

Despite the considerable number of accidents in Roraima state, evolution to deaths is rare, and in the analyzed period there were only 9 deaths (2013-2, 2014-2, 2015-1, and 2016-4 deaths). Out of the nine deaths, eight were caused by snakes, 75% affected by *Bothrops*, 12.5% by *Crotalus*, and 12.5% by *Micrurus* genera. Nevertheless, deaths from snakebites are known to be under-reported in the surveillance systems, which are estimated to be around 30% [55].

Beyond Brazil, other underdeveloped countries have also demonstrated high envenomings incidence within different numbers of mortality [56]. In Kenya, snakebites are considered a silent public health problem, presenting individual incidence between 2.7 to 6.7 per 100,000 inhabitants per year and 2.27% of mortality [57]. Although the country lacks a surveillance system to measure the incidence of snakebites, the Northern region of Cameroon presents 200 snakebites per 100,000 inhabitants [58]. In Costa Rica, there are reported 2 to 19 snakebite cases per 100,000 population per year [59]. On the other hand, Nigeria presents even higher snakebite incidence reaching 497 cases per 100,000 population per year and 12% of mortality. The high mortality in Nigeria can be explained by the low medical assistance, since only 8.5 % of snakebite victims attend hospitals in Nigeria [60]. Unlike underdeveloped countries, snakebite envenomings in developed countries are lower, as can be evidenced by Switzerland [61], France and Germany [62], Czech Republic [63], and USA [64-66]. Thus, a more detailed analysis of envenomings in underdeveloped countries is required, in order to design interventions that could reduce the impact of the public health problem in these countries.

## Conclusion

The envenomings registered in Roraima consist a huge and largely neglected health problem in Brazil. The challenge of the region is to specify the requirements for antivenom production for the local necessities. In order to do this, it is essential to organize the collection of epidemiological data. Thus, this study outlined the Roraima epidemiological profile of envenomings, which demonstrated to be caused most by snakes (*Bothrops* and *Crotalus* genera), affecting mainly men aged 20 to 39 years. Furthermore, Roraima envenomings mostly presented mild evolution, with the majority of victims being treated in less of 6 hours, and rare deaths. Our results contribute to the knowledge of local reality, which is needed to advocate measures that aim to reduce the accidents by venomous animal attacks and to promote therapeutic measures, such as supplying of specific antivenoms in places where they are most required.

## Acknowledgement

This work was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, The National Council for Scientific and Technological Development, scholarship to MBP grant number 307155/2017-0) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, São Paulo Research Foundation, scholarship to FAC grant number 2017/14035-1 and 2018/14158-9, and scholarship to ISO grant number 2017/03580-9 and 2018/21233-7). Also, the authors declare that they have no competing interests.

## References

- Gutiérrez JM, Theakston RD, Warrell DA (2006) Confronting the neglected problem of snake bite envenoming: The need for a global partnership. *PLoS Med* 3: 150.
- de Souza W (2010) *Doenças negligenciadas*. Rio de Janeiro.
- Molyneux D (2013) Neglected tropical diseases. *Community eye health* 26: 21-24.
- WHO (2019) Snakebite envenoming-A strategy for prevention and control.
- Chippaux JP (2015) Epidemiology of envenomations by terrestrial venomous animals in Brazil based on case reporting: From obvious facts to contingencies. *J Venom Anim Toxins incl Trop Dis* 21.
- Pucca MB, Oliveira F, Schwartz EF, Arantes EC, Lira-da-Silva RM (2014) Scorpionism and dangerous species of Brazil. *Toxinology* 1-24.
- Bochner R, Fiszon JT, Machado C (2014) A profile of snake bites in Brazil, 2001 to 2012. *J Clin Toxicol* 4: 194.
- Pimenta RJG, Brandão-Dias PFP, Leal HG, Carmo AOD, Oliveira-Mendes BBR, et al. (2019) Selected to survive and kill: *Tityus serrulatus*, the Brazilian yellow scorpion. *PLoS One* 14: 0214075.
- Chippaux JP (2009) [Impact of the environment on envenomation incidence and severity]. *Med Sci (Paris)* 25: 858-862.
- Machado C (2016) Um panorama dos acidentes por animais peçonhentos no Brasil/An overview of accidents involving venomous animals in Brazil/Un panorama de los accidentes por animales venenosos en Brasil. *Journal Health NPEPS* 1.
- Cardoso JLC (1993) *Acidentes por Animais Peçonhentos na Coordenação de Zoonoses e Animais Peçonhentos-Comentários e Sugestões*. Brasília: Ministério da Saúde.
- Leobas GF, Seibert CS, Feitosa SB (2016) *Acidentes por animais peçonhentos no Estado do Tocantins: Aspectos clínico-epidemiológicos*. *DESAFIOS* 2: 269-282.
- Chapin FS, Sala OE, Huber-Sannwald E (2013) *Global Biodiversity in a Changing Environment: Scenarios for the 21st Century*. Springer Science & Business Media, Berlin, Germany.
- Bernarde PS, Gomes JDO (2012) Serpentes peçonhentas e ofidismo em Cruzeiro do Sul, Alto Juruá, Estado do Acre, Brasil. *Acta Amazonica* 42: 65-72.
- Needleman RK, Neylan IP, Erickson T (2018) Potential environmental and ecological effects of global climate change on venomous terrestrial species in the wilderness. *Wilderness Environ Med* 29: 226-238.
- Fiszon JT, Bochner R (2008) Subnotificação de acidentes por animais peçonhentos registrados pelo SINAN no Estado do Rio de Janeiro no período de 2001 a 2005. *Revista Brasileira de Epidemiologia* 11: 114-127.
- Brasil MS (2015) *Análise dos dados epidemiológicos de 2010-2014*. Ministério da Saúde-Portal da Saúde.
- Bochner R, Struchiner CJ (2003) *Epidemiologia dos acidentes ofídicos nos últimos 100 anos no Brasil: uma revisão*. *Cad Saúde Pública* 19.
- Fetherston T (2015) The importance of critical incident reporting-and how to do it. *Community Eye Health* 28: 26-27.
- Moloo A (2019) WHO launches global strategy for prevention and control of snakebite envenoming.
- Nascimento SP (2000) [Epidemiological characteristics of snakebites in the state of Roraima, Brazil, 1992-1998]. *Cad Saude Publica* 16: 271-276.
- SINAN (2019) *Accident by Venomous Animals*.
- Henderson RW, Hoeyers LG (1977) The seasonal incidence of snakes at a locality in northern Belize. *Copeia* 1977: 349-355.
- Punde DP (2005) Management of snake-bite in rural Maharashtra: A 10-year experience. *Natl Med J India* 18: 71-75.
- Fry BG (2018) Snakebite: When the human touch becomes a bad touch. *Toxins (Basel)* 10: 170.
- Sawada SI, Kuklane K, Wakatsuki K, Morikawa H (2017) New development of research on Personal Protective Equipment (PPE) for occupational safety and health. *Ind Health* 55: 471-472.

27. Torp S, Grøgaard JB, Moen BE, Bråtveit M (2005) The impact of social and organizational factors on workers' use of personal protective equipment: A multilevel approach. *J Occup Environ Med* 47: 829-837.
28. dos Santos MC (1995) Serpentes de interesse médico da Amazônia: biologia, venenos e tratamento de acidentes. Universidade Do Amazonas, Amazonas, Brazil.
29. Barraviera B, Júnior RSF (2015) Acidentes ofídicos. In: Atheneu (ed.). Veronesi: Tratado de infectologia. Pg no: 2350-2367.
30. Leynaud GC, Reati GJ (2009) Identifying areas of high risk for ophidism in Cordoba, Argentina, using SIGEpi software. *Rev Panam Salud Publica* 26: 64-69.
31. Feitosa ES, Sampaio V, Sachett J, Castro DB, Noronha Md, et al. (2015) Snakebites as a largely neglected problem in the Brazilian Amazon: Highlights of the epidemiological trends in the State of Amazonas. *Rev Soc Bras Med Trop* 1: 34-41.
32. Abreu CB, Medeiros JM, Souza WMP, Cerni FA, Pereira UZ, et al. (2017) Epidemiological profile of accidents caused by venomous animals in Amazonas state. *Health and Diversity* 1: 96-103.
33. Magalhães SFV, Peixoto HM, Moura N, Monteiro WM, de Oliveira MRF (2019) Snakebite envenomation in the Brazilian Amazon: A descriptive study. *Trans R Soc Trop Med Hyg* 113: 143-151.
34. Malaque CMS (2014) Acidentes por animais peçonhentos. Hospital Vital Brazil Instituto Butantan, São Paulo, Brazil.
35. Loren Rebouças Santos H, Diego de Brito Sousa J, Arthur Alcântara J, de Almeida Gonçalves Sachett J, Soares Villas Boas T, et al. (2019) Rattlesnakes bites in the Brazilian Amazon: Clinical epidemiology, spatial distribution and ecological determinants. *Acta Trop* 191: 69-76.
36. Waldez F, Vogt RC (2009) Aspectos ecológicos e epidemiológicos de acidentes ofídicos em comunidades ribeirinhas do baixo rio Purus, Amazonas, Brasil. *Acta Amazonica* 39: 681-692.
37. Veronesi R, Focaccia R (2016) Tratado De Infectologia-2 Volumes. Editora Atheneu, São Paulo, Brazil.
38. Bernarde PS (2014) Serpentes Peçonhentas: E Acidentes Ofídicos No Brasil. Anolis Books Editora, São Paulo, Brazil.
39. Coser A, Wolff I, de Quadros RM, Marques SMT (2015) Acidentes ofídicos nas mesorregiões do estado de Santa Catarina. *Revista de Ciências Médicas e Biológicas* 14: 14-21.
40. Yañez-Arenas C, Díaz-Gamboa L, Patrón-Rivero C, López-Reyes K, Chiappa-Carrara X (2018) Estimating geographic patterns of ophidism risk in Ecuador. *Neotropical Biodiversity* 4: 55-61.
41. Dos Santos MC, Ferreira LCL, Da Silva WD, Maria de Fátima DF (1993) Caracterización de las actividades biológicas de los venenos 'amarillo' y 'blanco' de *Crotalus durissus ruruima* comparados con el veneno de *Crotalus durissus terrificus*. Poder neutralizante de los antivenenos frente a los venenos de *Crotalus durissus ruruima*. *Toxicon* 31: 1459-1469.
42. Dos-Santos MC, Assis EB, Moreira TD, Pinheiro J, Fortes-Dias CL (2005) Individual venom variability in *Crotalus durissus ruruima* snakes, a subspecies of *Crotalus durissus* from the Amazonian region. *Toxicon* 46: 958-961.
43. Wen FH, Monteiro WM, Silva AMM, Tambourgi DV, da Silva IM, et al. (2015) Snakebites and scorpion stings in the Brazilian Amazon: Identifying research priorities for a largely neglected problem. *PLoS Negl Trop Dis* 9: 0003701.
44. Oliveira IS, Pucca MB, Sampaio SV, Arantes EC (2018) Antivenomic approach of different *Crotalus durissus collilineatus* venoms. *J Venom Anim Toxins Incl Trop Dis* 24: 34.
45. Reckziegel GC, Junior VLP (2014) Análise do escorpionismo no Brasil no período de 2000 a 2010. *Rev Pan-Amaz Saude*.
46. Pardal PpDo, Gadelha MAdC, Menezes MMGO, Malheiros RS, Ishikawa EAY, et al. (2014) Envenenamento grave pelo escorpião *Tityus obscurus* Gervais, 1843. *Revista Pan-Amazônica de Saúde* 5: 65-70.
47. Pucca MB, Oliveira FN, Schwartz EF, Lira-da-Silva RM (2015) Scorpionism and Dangerous Species of Brazil Scorpion Venoms Gopalakrishnakone. Springer Reference 2015: 299-324
48. Acidentes por Animais Peçonhentos e Venenosos.
49. Roriz KRPS, Zaqueo KD, Setubal SS, Katsuragawa TH, Silva RRD, et al. (2018) Epidemiological study of snakebite cases in Brazilian Western Amazonia. *Rev Soc Bras Med Trop* 51: 338-346.
50. FUNASA (2001) Manual de diagnóstico e tratamento de acidentes por animais peçonhentos. Ministério da Saúde, Fundação Nacional de Saúde.
51. Sheikhtaheri A (2014) Near misses and their importance for improving patient safety. *Iran J Public Health* 43: 853-854.
52. Sharma SK, Bovier P, Jha N, Alirol E, Loutan L, et al. (2013) Effectiveness of rapid transport of victims and community health education on snake bite fatalities in rural Nepal. *Am J Trop Med Hyg* 89: 145-150.
53. Rechel B, Džakula A, Duran A, Fattore G, Edwards N, et al. (2016) Hospitals in rural or remote areas: An exploratory review of policies in 8 high-income countries. *Health Policy* 120: 758-769.
54. Chippaux JP (2017) Snakebite envenomation turns again into a neglected tropical disease! *J Venom Anim Toxins Incl Trop Dis* 23: 38.
55. da Silva Souza A, de Almeida Gonçalves Sachett J, Alcântara JA, Freire M, Alecrim MDGC, et al. (2018) Snakebites as cause of deaths in the Western Brazilian Amazon: Why and who dies? Deaths from snakebites in the Amazon. *Toxicon* 145: 15-24.
56. Kaushik M, Banyal V, Sondhi S, Thakur MK, Kaushik A (2018) Clinical profile of snake bite patients in tertiary care hospital in Himachal Pradesh: A prospective study. *International Journal of Research in Medical Sciences* 6: 2444-2448.
57. Ochola FO, Okumu MO, Muchemi GM, Mbaria JM, Gikunju JK (2018) Epidemiology of snake bites in selected areas of Kenya. *Pan Afr Med J* 29: 217.
58. Chippaux JP, Rage-Andrieux V, Le Mener-Delore V, Charronnière M, Sagot P, et al. (2002) Epidémiologie des envenimations ophidiennes dans le nord du Cameroun. *Bull Soc Pathol Exot* 95: 184-187.
59. Sasa M, Vazquez S (2003) Snakebite envenomation in Costa Rica: A revision of incidence in the decade 1990-2000. *Toxicon* 41: 19-22.
60. Habib AG, Gebi UI, Onyemelukwe GC (2001) Snake bite in Nigeria. *Afr J Med Med Sci* 30: 171-178.
61. Plate A, Kupferschmidt H, Schneemann M (2016) [Bites of venomous snakes in Switzerland]. *Praxis (Bern 1994)* 105: 679-685.
62. Schaper A, Desel H, Ebbecke M, De Haro L, Deters M, et al. (2009) Bites and stings by exotic pets in Europe: An 11 year analysis of 404 cases from Northeastern Germany and Southeastern France. *Clin Toxicol (Phila)* 47: 39-43.
63. Valenta J, Stach Z, Michalek P (2014) Exotic snake bites in the Czech Republic--Epidemiological and clinical aspects during 15-year period (1999-2013). *Clin Toxicol (Phila)* 52: 258-264.
64. Parrish HM (1966) Incidence of treated snakebites in the United States. *Public Health Rep* 81: 269-276.
65. O'Neil ME, Mack KA, Gilchrist J, Wozniak EJ (2007) Snakebite injuries treated in United States emergency departments, 2001-2004. *Wilderness Environ Med* 18: 281-287.
66. Othong R, Sheikh S, Alruwaili N, Gorodetsky R, Morgan BW, et al. (2012) Exotic venomous snakebite drill. *Clin Toxicol (Phila)* 50: 490-496.



- Journal of Anesthesia & Clinical Care  
Journal of Addiction & Addictive Disorders  
Advances in Microbiology Research  
Advances in Industrial Biotechnology  
Journal of Agronomy & Agricultural Science  
Journal of AIDS Clinical Research & STDs  
Journal of Alcoholism, Drug Abuse & Substance Dependence  
Journal of Allergy Disorders & Therapy  
Journal of Alternative, Complementary & Integrative Medicine  
Journal of Alzheimer's & Neurodegenerative Diseases  
Journal of Angiology & Vascular Surgery  
Journal of Animal Research & Veterinary Science  
Archives of Zoological Studies  
Archives of Urology  
Journal of Atmospheric & Earth-Sciences  
Journal of Aquaculture & Fisheries  
Journal of Biotech Research & Biochemistry  
Journal of Brain & Neuroscience Research  
Journal of Cancer Biology & Treatment  
Journal of Cardiology: Study & Research  
Journal of Cell Biology & Cell Metabolism  
Journal of Clinical Dermatology & Therapy  
Journal of Clinical Immunology & Immunotherapy  
Journal of Clinical Studies & Medical Case Reports  
Journal of Community Medicine & Public Health Care  
Current Trends: Medical & Biological Engineering  
Journal of Cytology & Tissue Biology  
Journal of Dentistry: Oral Health & Cosmesis  
Journal of Diabetes & Metabolic Disorders  
Journal of Dairy Research & Technology  
Journal of Emergency Medicine Trauma & Surgical Care  
Journal of Environmental Science: Current Research  
Journal of Food Science & Nutrition  
Journal of Forensic, Legal & Investigative Sciences  
Journal of Gastroenterology & Hepatology Research  
Journal of Gerontology & Geriatric Medicine  
Journal of Genetics & Genomic Sciences  
Journal of Hematology, Blood Transfusion & Disorders  
Journal of Human Endocrinology  
Journal of Hospice & Palliative Medical Care  
Journal of Internal Medicine & Primary Healthcare  
Journal of Infectious & Non Infectious Diseases  
Journal of Light & Laser: Current Trends  
Journal of Modern Chemical Sciences  
Journal of Medicine: Study & Research  
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology  
Journal of Neonatology & Clinical Pediatrics  
Journal of Nephrology & Renal Therapy  
Journal of Non Invasive Vascular Investigation  
Journal of Nuclear Medicine, Radiology & Radiation Therapy  
Journal of Obesity & Weight Loss  
Journal of Orthopedic Research & Physiotherapy  
Journal of Otolaryngology, Head & Neck Surgery  
Journal of Protein Research & Bioinformatics  
Journal of Pathology Clinical & Medical Research  
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance  
Journal of Physical Medicine, Rehabilitation & Disabilities  
Journal of Plant Science: Current Research  
Journal of Psychiatry, Depression & Anxiety  
Journal of Pulmonary Medicine & Respiratory Research  
Journal of Practical & Professional Nursing  
Journal of Reproductive Medicine, Gynaecology & Obstetrics  
Journal of Stem Cells Research, Development & Therapy  
Journal of Surgery: Current Trends & Innovations  
Journal of Toxicology: Current Research  
Journal of Translational Science and Research  
Trends in Anatomy & Physiology  
Journal of Vaccines Research & Vaccination  
Journal of Virology & Antivirals  
Archives of Surgery and Surgical Education  
Sports Medicine and Injury Care Journal  
International Journal of Case Reports and Therapeutic Studies

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>