

## Research Article

### Insecticidal Activity of Ethanolic Leaf Extracts against the Maize Weevil, *Sitophilus zeamais* Motsch (Coleoptera: Curculionidae)

\*Suleiman M and Sani I

Department of Biology, Umaru Musa Yar'adua University, Katsina, Nigeria

#### Abstract

Laboratory studies were conducted in order to evaluate the toxicity of ethanolic leaf extracts of *Euphorbia balsamifera* Aiton, *Lawsonia inermis* L., *Mitracarpus hirtus* (L.) DC and *Senna obtusifolia* L. against *Sitophilus zeamais* Motsch. Ten 7 day old *S. zeamais* were released in 20g of disinfested sorghum grains treated with ethanolic extracts of the botanicals in plastic bottles at different concentrations of 2.5, 5.0 and 10.0x10<sup>4</sup> ppm at 30°C and 70% RH. Adult mortalities and lethal concentrations that killed 50% (LC<sub>50</sub>) of the weevils' population at 24 Hours After Treatment (HAT) were determined. Concentration of body protein as well as inhibition rate of Acetylcholinesterase (AChE) activity in *S. zeamais* exposed to sorghum treated with the botanical extracts were also investigated. Results showed that the adult mortality ranged from 82.50 ± 2.50 to 100.00 ± 0.00% within 3 Days After Treatment (DAT). *L. inermis* had the highest LC<sub>50</sub> value (1.72 x 10<sup>4</sup> ppm), while *E. balsamifera* had the lowest (1.12 x 10<sup>4</sup> ppm). Biochemical tests showed that the botanicals reduced body protein and inhibited the activity of Acetylcholinesterase (AChE) in *S. zeamais*. The botanicals were found to be toxic to adult *S. zeamais*

\*Corresponding author: Suleiman M, Department of Biology, Umaru Musa Yar'adua University, Katsina, Nigeria, Tel: +234 8067195818; E-mail: mohammed.suleiman@umyu.edu.ng

**Citation:** Suleiman M, Sani I (2018) Insecticidal Activity of Ethanolic Leaf Extracts against the Maize Weevil, *Sitophilus zeamais* Motsch (Coleoptera: Curculionidae). Arch Zool Stud 1: 004.

**Received:** September 01, 2018; **Accepted:** October 26, 2018; **Published:** November 09, 2018

**Copyright:** © 2018 Suleiman M and Sani I. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and therefore could be utilized to reduce the weevils' infestations in sorghum grains.

**Keywords:** Acetylcholinesterase; Body protein; Botanicals; LC<sub>50</sub>; Mortality; *Sitophilus zeamais*

#### Introduction

Synthetic insecticides have been commonly used for controlling pests in stored products [1-4]. However, this has resulted in problems such as pest resurgence and increase in costs of application arising from the development of resistance to insecticides [5,6]. This has led to the search of alternative, eco-friendly methods of controlling insect pests of storage such as the use of botanicals [7,8]. In this regard the toxicity of many plant materials has been evaluated against *S. zeamais* in stored grains with recorded success [7,9-12].

One of the ways used to evaluate the toxicity of insecticides is by measuring their mortality effects on insects. Ethanolic, methanolic and acetic extracts of various plant materials have been reported to cause high mortality of *S. zeamais* [9,12-14]. It was reported that some medicinal plants such as *olax subscorpioidea* Oliver (Santalales: Olacaceae), *afmomum melegueta k. schum* (Zingiberales: Zingiberaceae), and *zingiber officinale* Rosce (Zingiberales: Zingiberaceae) caused adult mortality of *S. zeamais* which ranged from 18.35 to 88.35% after 72 hours of exposure [15]. Tilahun and Daniel reported the toxicity effects of seed powder of *Azadirachta indica a. Juss* (Sapindales: Meliaceae) against *S. zeamais* and recorded 23.00 to 28.67% adult mortality after 45 days of treatment [16].

Another way of assessing toxicity of insecticides is by evaluating their effects on biochemical parameters of insects [17]. Kumar and Michael noted that the carbohydrates, proteins and lipids play an important role in the biochemical process underlying growth and development of insects [18]. Fox, et al., reported that the reproductive capacity of adult female insects is affected largely by their body protein content [19]. Piri, et al., noted that proteins are important for individual-level fitness-associated traits such as body size, growth rate and fecundity [20]. At high levels of organization, proteins have been linked to population dynamics, life cycles and even biological diversification [20]. Some of the biochemical composition of *Acanthoscelides obtectus* Say, *S. oryzae*, *S. granarius* and *T. confusum* have been investigated [17-19,21,22]. However, little is known with regards to toxicity of botanicals against body protein of *S. zeamais*.

In addition to body protein, enzymes such as Acetylcholinesterase (AChE) play a vital role in insect survival. AChE is a key enzyme of the cholinergic system regulating the level of acetylcholine and terminates nerve impulses by catalyzing the hydrolysis of acetylcholine [23]. Some botanicals were reported to inhibit AChE activity in some insect pests of storage such as *S. oryzae*, *S. granarius* and *T. castaneum*, but little is known on effects of botanicals in inhibiting AChE activity in *S. zeamais* [24-28]. This study was carried out to investigate the ability of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* to cause adult mortality, reduce concentration of body protein and inhibit AChE activity in *S. zeamais*.

## Materials and Methods

### Mass rearing of *S. zeamais*

One hundred adults of *S. zeamais* were obtained from infested sorghum grains at Katsina Central Market and then introduced into 500ml rearing bottles containing 250g of the disinfested sorghum grains. The bottles were covered with muslin cloth and secured with rubber bands. They were then kept in an incubator for oviposition at 30°C and 70% RH for seven days, after which the parents were removed. The bottles were maintained in the incubator under the same condition for adult emergence. Progeny of 7 days old were sieved and used for the laboratory experiments.

### Preparation of the botanicals

Fresh leaves of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* were shade-dried and ground into powder using a laboratory blender. One hundred grams of each plant powder was dissolved in 400ml of ethanol in conical flasks. Mouths of the conical flasks were properly corked and kept in a refrigerator for 48 hours. The extract was separated using muslin cloth and filtered with Whatman No.1 filter papers as described by Khaliq, et al., [29]. The filtrate was separately concentrated by evaporating excess solvents using rotary evaporator with rotary speed of 3 to 6 rpm for 8 hours. The resulting extract was air-dried to remove traces of the solvent and stored in refrigerator at 4°C [30].

### Adult mortality tests of *S. zeamais* exposed to sorghum grains treated with ethanolic leaf extracts

The method of de Oliveira, et al., was adopted to assess adult mortality of *S. zeamais* in sorghum grains treated with ethanolic leaf extracts of the test plants [31]. Crude extracts were diluted with ethanol to make different concentrations of 2.5, 5.0 and 10.0 x 10<sup>4</sup> ppm. Twenty grams of sorghum grains were weighed and put into 250 ml plastic bottles. The grains were impregnated with 2ml of the ethanolic extracts at three concentrations, while the control contained the grains. Ten weevils were released into each of the bottles after complete evaporation of the solvent. The grain mass was mixed thoroughly with the aid of glass rod and air-dried until complete solvent evaporation. All treatments were arranged in a Completely Randomized Design (CRD) in the incubator and replicated four times. The set-ups were inspected daily and dead weevils in each treatment were removed and recorded daily for three days.

Adult Mortality was assessed as;

$$\% \text{Mortality} = \left( \frac{\text{Number of Dead Weevils}}{\text{Total Number of Weevils Introduced}} \right)$$

### Determination of Lethal Concentrations (LC<sub>50</sub>) of ethanolic leaf extracts

To evaluate LC<sub>50</sub> of the ethanolic extracts of the botanicals against *S. zeamais*, methods of Ebadollahi and Mahboubi were employed [32]. The number of dead weevils at the end of 24 hour exposure to the extracts was used to determine the LC<sub>50</sub> of the botanicals by using probit analysis with SPSS (version 16.0) software package [33].

### Determination of body protein in *S. zeamais* treated with ethanolic leaf extracts

Samples for protein assays were prepared following Askar, et al.,

[17]. Adult weevils exposed to LC<sub>50</sub> of ethanolic extracts of the botanicals for 7 days were freeze killed at -20°C. A sample of 0.5g of the insects was taken from each bottle and homogenized in 10 times (5mL) volumes (w/v) of phosphate buffer (pH 7.2) using a glass homogenizer under ice. The total homogenates were centrifuged at 4000 rpm for 10min at 4°C using a centrifuge. The supernatant was transferred to new eppendorfs tubes and preserved at -20°C for spectrophotometry.

The body protein was estimated following the method of Lowry, et al., in triplicates with Bovine Serum Albumin (BSA) as the standard [34]. The absorbance was measured spectrophotometrically after 30mins at 750nm against the blank for both test samples and the standard solutions. The protein concentration was calculated and expressed as µg/ml wet tissue using the formula:

$$\text{Concentration of Body Protein } x = \frac{y}{m}$$

Where:

x = The concentration

y = Absorbance of the solution

m = Slope of the standard curve

### Determination of AChE activity inhibition in *S. zeamais* exposed to ethanolic leaf extracts

*S. zeamais* adults were fumigated with LC<sub>50</sub> of ethanolic extracts of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* as fumigant toxicity assay. After seven days of fumigation, the weevils were homogenized in phosphate buffer saline (50mM, pH 8) using a Teflon glass tissue homogenizer. The homogenate was centrifuged at 10,000 rpm for 20min at 4°C. The supernatant was filtered and used as the AChE preparation. Methods of Ellman, et al., and Chaubey were adopted for the assay and AChE activity was determined spectrophotometrically [28,35]. Absorbance in the treated samples was evaluated at 412nm for 6min and compared to that of the control. All measurements were made in triplicate [25,26,28].

AChE activity inhibition was calculated as follows [26]:

$$\% \text{ AChE Activity Inhibition} = \frac{A_{\text{Control}} - A_{\text{Treatment}}}{A_{\text{Control}}} \times 100$$

Where:

A<sub>Control</sub> = The absorbance of the untreated control

A<sub>Treatment</sub> = The absorbance of the treated samples

### Data analysis

Data generated were tested for normality using Shapiro-Wilk and Jacques-Bera normality tests [36-38]. Data from adult mortality tests were subjected to two-way ANOVA using Graph Pad Prism (version 7.03). Probit analysis was employed to calculate LC<sub>50</sub> of the extracts with the significant differences determined by X<sup>2</sup> goodness of fit using SPSS (version 16.0). One-way ANOVA was used to test if there were significant differences in the body protein and AChE activity inhibition in adults of *S. zeamais* exposed to LC<sub>50</sub> of the extracts by using PAST (version 2.17). Significantly different means were separated by Bonferroni's and Tukey's multiple comparisons tests [38]. All analyses were performed as statistically significant p<0.05.

## Results

### Adult mortality of *S. zeamais* in sorghum grains treated with ethanolic leaf extracts

Ethanolic leaf extracts of the selected botanicals applied at different concentrations of 2.5, 5.0 and 10.0x10<sup>4</sup> ppm exhibited varying percentage mortalities of *S. zeamais* within 1, 2 and 3 days after treatment (Table 1).

Treatments	Conc.(x 10 <sup>4</sup> ppm)	Mean Adult Mortality (% ± S. E.)		
		Days After Treatment (DAT)		
		1	2	3
<i>E. balsamifera</i>	2.5	65.00 ± 2.89 <sup>bc</sup>	90.00 ± 4.08 <sup>ab</sup>	100.00 ± 0.00 <sup>a</sup>
	5	75.00 ± 2.89 <sup>ab</sup>	95.00 ± 2.89 <sup>a</sup>	100.00 ± 0.00 <sup>a</sup>
	10	85.00 ± 2.89 <sup>a</sup>	100.00 ± 0.00 <sup>a</sup>	100.00 ± 0.00 <sup>a</sup>
<i>L. inermis</i>	2.5	55.00 ± 2.89 <sup>c</sup>	77.50 ± 4.79 <sup>b</sup>	97.50 ± 2.50 <sup>a</sup>
	5	72.50 ± 2.50 <sup>b</sup>	92.50 ± 2.50 <sup>ab</sup>	100.00 ± 0.00 <sup>a</sup>
	10	77.50 ± 2.50 <sup>ab</sup>	95.00 ± 2.89 <sup>a</sup>	100.00 ± 0.00 <sup>a</sup>
<i>M. hirtus</i>	2.5	67.50 ± 4.89 <sup>bc</sup>	80.00 ± 4.08 <sup>ab</sup>	95.00 ± 2.89 <sup>a</sup>
	5	70.00 ± 4.08 <sup>bc</sup>	92.50 ± 4.89 <sup>ab</sup>	100.00 ± 0.00 <sup>a</sup>
	10	75.00 ± 2.89 <sup>ab</sup>	95.00 ± 2.89 <sup>a</sup>	100.00 ± 0.00 <sup>a</sup>
<i>S. obtusifolia</i>	2.5	55.00 ± 2.89 <sup>c</sup>	72.50 ± 2.50 <sup>b</sup>	82.50 ± 2.50 <sup>b</sup>
	5	67.50 ± 4.79 <sup>bc</sup>	85.00 ± 2.89 <sup>ab</sup>	95.00 ± 2.89 <sup>a</sup>
	10	70.00 ± 4.08 <sup>bc</sup>	90.00 ± 4.08 <sup>ab</sup>	100.00 ± 0.00 <sup>a</sup>
Control	0	0.00 ± 0.00 <sup>d</sup>	0.00 ± 0.00 <sup>c</sup>	0.00 ± 0.00 <sup>c</sup>

**Table 1:** Mortality of adult *S. zeamais* exposed to 2.5, 5.0 and 10.0 x 10<sup>4</sup> ppm of ethanolic botanical extracts in three days after treatment.

Conc = Concentration

Means in the same column followed by different letter superscript are significantly different at p<0.05 by the Bonferroni's Multiple Comparisons Test.

Analysis of variance indicated that the difference in adult mortalities of *S. zeamais* was highly significant among the botanicals (F (4, 45) = 2947.00, p<0.0001) as well as within the concentrations (F (2, 45) = 18.00, p<0.0001). Bonferroni's multiple comparisons test revealed that at 2.5 x 10<sup>4</sup> ppm, *E. balsamifera*, *L. inermis* and *M. hirtus* caused statistically similar mortality which was higher than that in *S. obtusifolia* at 3 DAT. At 5.0 and 10.0 x 10<sup>4</sup> ppm, all the botanicals had the same effect on the weevils' mortality within 3 DAT.

### Lethal Concentration (LC<sub>50</sub>) of ethanolic leaf extracts against *S. zeamais*

Table 2 shows that *L. inermis* had the highest LC<sub>50</sub> value as 1.72 x 10<sup>4</sup> ppm, while *E. balsamifera* had the lowest as 1.12 x 10<sup>4</sup> ppm at 24 HAT. The probit equations were y = 1.074x - 0.053, y = 1.065x - 0.251, y = 0.365x + 0.296 and y = 0.669x - 0.100 for *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia*, respectively (Table 2). The slopes varied between 0.365 ± 0.723 and 1.074 ± 0.739. X<sup>2</sup> goodness of fit (0.005 - 0.092) reveals that the LC<sub>50</sub> of ethanolic extracts against the weevils was not significantly different (p>0.05) among concentrations of each of the botanicals.

### Concentration of body protein in *S. zeamais* exposed to ethanolic leaf extracts


The protein in *S. zeamais* exposed to ethanolic extracts of the

botanicals appeared to follow the order: *E. balsamifera*<*L. inermis*<*M. hirtus*<*S. obtusifolia*<control (Figure 1). Significant difference (F (4, 10) = 2861.00, p < 0.0001) existed in protein content among the weevils exposed to sorghum grains treated with LC<sub>50</sub> of ethanolic extracts of different test botanicals. Further, Tukey's multiple comparisons test revealed that all treatments differed from each other with insects exposed to *E. balsamifera* having the least body protein content.

Botanicals	LC <sub>50</sub> (x 10 <sup>4</sup> ppm)	95% CI	Slope ± S.E.	X <sup>2</sup>	p value
<i>E. balsamifera</i>	1.12	0.97 - 3.11	1.074 ± 0.739	0.005	0.946
<i>L. inermis</i>	1.72	0.88 - 3.01	1.065 ± 0.712	0.092	0.762
<i>M. hirtus</i>	1.15	1.07 - 2.30	0.365 ± 0.723	0.006	0.938
<i>S. obtusifolia</i>	1.41	1.21 - 2.55	0.669 ± 0.701	0.065	0.798

**Table 2:** LC<sub>50</sub> of ethanolic leaf extracts against adult *S. zeamais* at 24 HAT.


HAT = Hours After Treatment; CI = Confidence Interval; X<sup>2</sup> = Chi-square.



**Figure 1:** Concentration of body protein in *S. zeamais* exposed to LC<sub>50</sub> of ethanolic leaf extracts.

### Inhibition of AChE activity in *S. zeamais* by ethanolic leaf extracts

The inhibition of AChE activity in *S. zeamais* by ethanolic extracts was 79.61, 77.65, 74.90 and 70.98 for *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia*, while none was recorded in weevils exposed to the untreated grains (Figure 2). One-way ANOVA revealed that inhibition of the AChE activity was significantly different (F (4, 10) = 4190.00, p<0.0001) among the botanical treatments. Tukey's pairwise comparisons showed that the inhibition activity was statistically similar in weevils exposed to *E. balsamifera* and *L. inermis* but higher than the rest.



**Figure 2:** AChE activity inhibition in *S. zeamais* exposed to LC<sub>50</sub> of ethanolic leaf extracts.

## Discussion

### Effect of ethanolic extracts on adult mortality of *S. zeamais*

Ethanolic extracts of the selected botanicals were found to be highly effective by resulting in total mortality of adults of *S. zeamais* in sorghum grains at all concentrations of *E. balsamifera* and higher rates of *L. inermis*, *M. hirtus* and *S. obtusifolia*. The efficacy of the test botanicals is in line with earlier findings in which ethanolic extracts of some plant species caused adult mortality of maize weevils [10,13]. Ajayi reported that ethanolic extract of *D. regia* seeds applied at 2.0mg/g maize grains resulted in 12.50% adult mortality of *S. zeamais* after five days of exposure [10]. Similarly, findings of Ibrahim, et al., showed that application of ethanolic leaf extract of *C. odorata* at 10ml / 50g maize caused 14.00% adult mortality of *S. zeamais* at 7 DAT [13]. The observed increase in mortality of *S. zeamais* caused by ethanolic extracts with increase in post treatment period in this study agrees with previous findings [10,12,13].

Ethanolic leaf extracts of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* were generally found toxic to adult *S. zeamais* and suppressed their survival in sorghum grains by causing considerably high mortality. All the selected botanicals were effective against the weevil even at lower concentrations. This was possible because plant species contain secondary metabolites which are vast storehouse of compounds such as the steroids, phenolic compounds and tannins with wide range of biological activity reported to have great impact on insecticidal activities [39,40]. Other bioactive compounds such as terpenoids, flavonoids, alkaloids, saponins and glycosides were found in the leaf extracts of *E. balsamifera*, *L. inermis* and *S. obtusifolia* by others and reported to be pesticidal in nature [40-42]. Ileke, et al., reported the presence of alkaloids, flavonoids, saponins and tannins, in the powders and methanolic extracts of *M. fragrans* and *A. melegueta* and concluded their insecticidal activity against *S. zeamais* [14]. The characteristic smell of the study botanicals might have also contributed to their insecticidal activity by repelling the insects away from the grains as suggested by Ileke, et al., [14].

### Lethal Concentration (LC<sub>50</sub>) of ethanolic leaf extracts against *S. zeamais*

The present study investigated LC<sub>50</sub> of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* against *S. zeamais* in sorghum grains due to lack of information on this aspect in previous studies. However, lethal concentrations of other botanicals used to control insect pests of storage were determined [43-45].

LC<sub>50</sub> of the plants showed that the ethanolic extracts had great efficiency by causing 50% adult mortality of *S. zeamais* in sorghum even at a concentration below the lowest amount applied within 24 HAE. *E. balsamifera* was more effective than the other botanicals followed by *M. hirtus*. The effectiveness the selected botanicals in killing adult weevils at lower concentration within short period is in conformity with Biswas, et al., who reported high efficiency of *L. inermis* against *T. castaneum* [39].

Low LC<sub>50</sub> values of ethanolic extracts of the test botanicals concur with Rani and Devanand, who reported that among methanolic extracts of the plants tested *Momordica charantia* L [43]. was the most effective with LC<sub>50</sub> of 2.82mg / 20g maize grains against *S. zeamais*.

### Effect of ethanolic leaf extracts on body protein of *S. zeamais*

There was a decrease in body protein of the maize weevils exposed to lethal concentrations of all the botanicals compared to the control. The reduction in body protein agrees with Askar et al., who reported that both clove oil and diatomaceous earth reduced body protein of *S. zeamais* and *S. oryzae* from 0.10mg/ml in the control to 0.09 and 0.06mg/ml, respectively [17]. It was earlier reported by Piri, et al., that sub-lethal concentrations of spinosad resulted in a significant decrease in protein content of *Glyphodes pyloalis* Walker [20]. This decrease in body protein could be attributed to low feeding efficiency of the insects due to antifeedant effect of many other insecticides and hence a decrease in protein concentration [40,46]. Another reason might be as a result of protein break down into amino acids which help in energy supply to the insect [47].

It was found that application of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* in the form of powders and extracts significantly affected body protein of *S. zeamais* which could probably affect the aforementioned biological aspects of the weevil. The reduction in concentration of the body protein might have contributed to decline in the insect's population and hence decreased infestations in the sorghum grains. This could also be attributed to decrease in vitellogenin, a precursor of yolk component, present in the body protein of vitellogenic females, which its amount reflects the insect reproductive performance [48].

### Effect of ethanolic leaf extracts on the activity of AChE in *S. zeamais*

AChE is a hydrolytic enzyme in insect body that influences the nervous system [27]. In the present study the LC<sub>50</sub> of ethanolic extracts of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* significantly inhibited the activity of AChE in *S. zeamais* agreeing with Li, et al., who reported that methyl alcohol, ethyl acetate and petroleum ether fruit extracts of *Illicium verum hook. F* [27], inhibited the activity of AChE in *S. zeamais*. Similarly, the LC<sub>50</sub> of essential oils of *Pimpinella anisum* and *Ocimum basilicum* have demonstrated their interference with AChE activity in *S. granarius* [25]. Recently, methanolic extract of *Urginea maritime* at a concentration of 1000µg/ml was found to result in 73.37% inhibition of the activity of AChE in *S. oryzae* [26]. Also, another finding has confirmed the inhibitory effect of botanicals on the activity of AChE where Chaubey reported that LC<sub>50</sub> of *Cuminum cyminum* essential oil reduced AChE activity in *S. zeamais* to 66.90% compared to 31.59% of the control [28].

The inhibitory effect on AChE activity in *S. zeamais* exposed to lethal concentrations of all the botanicals has shown their possible potentiality as biopesticides. This is because it has been reported that inhibition of AChE in cholinergic synapses of the nervous system is the primary mechanism of acute toxicity of insecticides [25]. Therefore, the botanicals probably have neurotoxic effects on *S. zeamais* by interfering with the passage of impulses in the insect nervous system leading to inability of AChE to hydrolyze acetylcholine which builds up in the synapse and results in excessive neuro excitation [24]. The mortality effects of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* against *S. zeamais* recorded in this study may also be attributed to their ability to inhibit the activity of AChE as suggested by Rajashekar, et al., [24]. Thus, the neurotransmitting system of the weevils represents a target for insect control.

## Conclusion

It was found that application of ethanolic extracts of the selected botanicals resulted in high adult mortality of *S. zeamais* even at the lower concentration of  $2.5 \times 10^4$  ppm after three days of treatment with *E. balsamifera* as the most effective followed by *L. inermis* and therefore could be used to reduce *S. zeamais* infestations in sorghum grains.

The  $LC_{50}$  of the botanicals were found to be either lower than or within the range of recommended dose/concentration (1-5%) of botanicals in stored commodity. Findings of this study have revealed that application of *E. balsamifera*, *L. inermis*, *M. hirtus* and *S. obtusifolia* at the lowest concentration of  $2.5 \times 10^4$  ppm could cause the death of 50% *S. zeamais* within one to three days. Lethal concentrations of all the botanicals reduced body protein of *S. zeamais* and have the ability to interfere with neurotransmitting system of the weevil by inhibiting the activity of AChE.

The toxicity of the test botanicals against adult *S. zeamais* infers that they could be utilized to reduce the weevils' infestations in stored sorghum. However, further investigations on their toxicity on mammals and other insect pests are hereby recommended.

## References

1. Nguemtchouin MGM, Ngassoum MB, Chaliel P, Kamga R, Ngamo LST, et al. (2013) *Ocimum gratissimum* essential oil and modified montmorillonite clay, a means of controlling insect pests of stored products. *J Stored Prod Res* 52: 57-62.
2. Baributsa D, Baoua IB, Bakoye ON, Amadou L, Murdock LL (2017) PICS bags safely store unshelled and shelled groundnuts in Niger. *J Stored Prod Res* 72: 54-58.
3. Nayak MK, Daglish GJ (2017) Base-line susceptibility of field populations of *Rhyzopertha dominica* (F.) to spinosad in Australia. *J Stored Prod Res* 70: 1-6.
4. Paudyal S, Opit GP, Osekre EA, Arthur FH, Bingham GV, et al. (2017) Field evaluation of the long-lasting treated storage bag, deltamethrin incorporated, (ZeroFly® Storage Bag) as a barrier to insect pest infestation. *J Stored Prod Res* 70: 44-52.
5. Liu ZL, Zhao NN, Liu CM, Zhou L, Du SS (2012) Identification of insecticidal constituents of the essential oil of *Curcuma wenyujin* rhizomes active against *Liposcelis bostrychophila* badonnel. *Molecules* 17: 12049-12060.
6. Cosmas P, John CT, Agathar K, Ronald M, Kufa M, et al. (2018) Use of botanical pesticides in controlling *Sitophilus zeamais* (Maize Weevil) on stored *Zea mays* (Maize) grain. *Modern Concepts and Developments in Agronomy* 1: 1-4.
7. Suleiman M (2014) Efficacy of some spices as sorghum grain protectants against *Sitophilus zeamais* Motschulsky [Coleoptera: Curculionidae]. *African Journal of Agricultural Research* 9: 841-845.
8. Suleiman M, Rugumamu CP (2017) Management of insect pests of stored sorghum using botanicals in Nigerian traditional stores. *Journal of Stored Products and Postharvest Research* 8: 93-102.
9. Adedire CO, Akinkulore RO, Obembe OM (2012). Efficacy of cashew kernel extracts in the control of the maize weevil, *Sitophilus zeamais* (Coleoptera: Curculionidae). *Archives of Phytopathology and Plant Protection* 45: 831-839.
10. Ajayi OE (2013) Insecticidal activity of powder and extracts of *Delonix regia* seed against maize weevil, *Sitophilus zeamais* (Coleoptera: Curculionidae). *FUTA Journal of Research in Sciences* 1: 54-62.
11. Wini GJ, Nukenine EC, Ndjonka D, Suh C, Adler C, et al. (2015) Efficacy of diatomaceous earth and wood ash for the control of *Sitophilus zeamais* in stored maize. *Journal of Entomology and Zoology Studies* 3: 390-397.
12. Suleiman M, Rugumamu CP, Ibrahim ND (2018) Insecticidal toxicity of some botanicals against *Sitophilus zeamais* Motsch. (Coleoptera: Curculionidae) in stored sorghum grains in Nigeria. *Journal of Entomology and Zoology Studies* 6: 1280-1287.
13. Ibrahim YJ, Abugri AD, Afun JVK (2016) Efficacy of ethanolic leaf extract of *Chromolaena odorata* in controlling *Sitophilus zeamais* in stored maize. *Journal of Experimental Agriculture International* 14: 1-10.
14. Ileke KD, Adesina JM, Obajulaye EO (2016) Synergic effects of two botanical entomocides as pest-protectants in maize grains. *Journal of Biological Research* 89: 34-39.
15. Oni MO, Ogungbite OC (2015) Entomotoxicant potential of powders and oil extracts of three medicinal plants in the control of *Sitophilus zeamais* infesting stored maize. *Journal of Plant and Pest Science* 2: 8-17.
16. Tilahun FE, Daniel DH (2016) Effect of neem leaf and seed powders against adult maize weevil (*Sitophilus zeamais* Motschulsky) mortality. *International Journal of Agricultural Research* 11: 90-94.
17. Askar SI, Al-Assaal MS, Nassar AMK (2016) Efficiency of some essential oils and insecticides in the control of some *Sitophilus* insects (Coleoptera: Curculionidae). *Egypt J Plant Pro Res* 4: 39-55.
18. Kumar MDA, Michael AS (2012) Changes in the activities of proteins and carbohydrates in haemolymph of silkworm, *bombyx mori* treated with greenleaf. *International Journal of Environmental Sciences* 3: 119-125.
19. Fox CW, Stillwell RC, Wallin WG, Hitchcock LJ (2006) Temperature and host species affect nuptial gift size in a seed-feeding beetle. *Funct Ecol* 20: 1003-1011.
20. Piri F, Sahragard A, Ghadamyari M (2014) Sublethal Effects of Spinosad on Some Biochemical and Biological Parameters of *Glyphodes pyralis* Walker (Lepidoptera: Pyralidae). *Plant Protect Sci* 50: 135-144.
21. Ali NS, Ali SS, Shakoori A (2011) Effects of sublethal doses of talstar on biochemical components of melathion-resistant and -susceptible adults of *Rhyzopertha dominica*. *Pakistan J Zool* 43: 879-887.
22. Lingampally V, Solanki VR, Kaur A, Raja SS (2013) Effect of andrographolide on the protein content of *Tribolium Confusum* (Duvall). *International Journal of Pure and Applied Zoology* 1: 70-79.
23. Praveena A, Sanjayan KP (2011) Inhibition of acetylcholinesterase in three insects of economic importance by linalool, a monoterpene phytochemical. *Insect Pest Management, A Current Scenario* 340-345.
24. Rajashekar Y, Raghavendra A, Bakthavatsalam N (2014) Acetylcholinesterase inhibition by biofumigant (Coumaran) from leaves of *Lantana camara* in stored grain and household insect pests. *Biomed Res Int* 2014: 1-6.
25. Abo El Makarem H, El Kholy SE, Abdel-Latif A, Seif AI (2015) Physiological and biochemical effects of some essential oils on the granary weevil, *sitophilus granarius* (L.) (coleoptera: curculionidae). *Egypt J Exp Biol (Zoo.)* 11: 117-123.
26. Maazoun AM, Hlel TB, Hamdi SH, Belhadj F, Jemâa JMB, et al. (2017) Screening for insecticidal potential and acetylcholinesterase activity inhibition of *urginea maritima* bulbs extract for the control of *Sitophilus oryzae* (L.). *J Asia-Pac Entomol* 20: 752-760.
27. Li J, Guo Q, Han S, Jiang L, Liang G (2013) Types, morphologies and distribution of antennal sensilla of *Quadrastichus erythrinae* (Hymenoptera: Eulophidae). *Fla Entomol* 96: 1288-1297.
28. Chaubey MK (2017) Evaluation of insecticidal properties of *Cuminum cyminum* and *Piper nigrum* essential oils against *Sitophilus zeamais*. *Journal of Entomology* 14:148-154.

29. Khaliq A, Nawas A, Ahmad MH, Sagheer M (2014) Assessment of insecticidal potential of medicinal plant extracts for control of maize weevil, *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae). *Basic Research Journal of Agricultural Science and Review* 3: 100-104.
30. Abou-Elnaga, Zeinab Sh (2015) Efficacy of extracts of some Egyptian plants against economically important stored grains pest *Sitophilus oryzae* L. *Journal of Entomology and Zoology Studies* 3: 87-91.
31. de Oliveira TA, Ronche-Teles B, da Fonseca CRV, da Silva SLR, Santos PA, et al. (2012) Insecticidal activity of *Vitex cymosa* (Lamiaceae) and *Eschwehlera pedicellata* (Lecythidaceae) extracts against *Sitophilus zeamais* adults (Curculionidae). *Emir J Food Agr* 24: 49-56.
32. Ebadollahi A, Mahboubi M (2011) Insecticidal activity of the essential oil isolated from *Azilia eryngioides* (PAU) hedge et lamond against beetle pests. *Chil J Agr Res* 71: 406-411.
33. SAS Institute (1999) SAS/STAT User's Guide, Version 8, Volume 2. SAS Institute, North Carolina, USA.
34. Lowry OH, Passonneau JV, Rock MK (1961). The stability of pyridine nucleotides. *J Biol Chem* 236: 2756-2759.
35. Ellman GL, Courtney KD, Andres V Jr., Featherstone RM (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol* 7: 88-95.
36. Sokal RR, Rohlf FJ (1995) *Biometry* (3rd edn): A Series of books in biology. WH Freeman and Company, New York, USA. Pg no: 887.
37. Jolicoeur P (1999) *Introduction to Biometry* (1st edn). Springer+Business Media LLC, New York, USA. Pg no: 516.
38. Le CT (2003) *Introductory Biostatistics*. John Wiley & Sons Inc., New Jersey, USA, Pg no: 553.
39. Biswas KK, Sharmin N, Rabbi MA (2016) Evaluation of the insecticidal activity of *Lawsonia inermis* Linn. against the red flour beetle, *Tribolium castaneum* (Herbst). *Natural Products* 12: 8-11.
40. Hikal WM, Baeshen RS, Said-Al Ahl HAH (2017) Botanical insecticides as simple extractives for pest control. *Cogent Biology* 3: 1-16.
41. Doughary JH, El-mahmood AM, Tyoyina I (2008) Antimicrobial activity of leaf extracts of *Senna obtusifolia* (L.). *Afr J Pharm Pharmacol* 2: 007-013.
42. Wagini NH, Soliman AS, Abbas MS, Hanafy YA, Badawy EM (2014) Phytochemical analysis of Nigerian and Egyptian henna (*Lawsonia inermis* L.) leaves using TLC, FTIR and GCMS. *Plant* 2: 27-32.
43. Rani PU, Devanand P (2011) Efficiency of different plant foliar extracts on grain protection and seed germination in maize. *Research Journal of Seed Science* 4: 1-14.
44. Naeem M, Anjum SI, Ismail M, Khan J, Bibi S, et al. (2015) Laboratory assessment of different botanical extracts and cypermethrin against insect pests. *Journal of Entomology and Zoology Studies* 3: 84-88.
45. Jayakumar M, Arivoli S, Raveen R, Tennyson S (2017) Repellent activity and fumigant toxicity of a few plant oils against the adult rice weevil *Sitophilus oryzae* Linnaeus 1763 (Coleoptera: Curculionidae). *Journal of Entomology and Zoology Studies* 5: 324-335.
46. Etebari K, Mirhoseini SZ, Matindoost L (2005) A study on interspecific diversity of eight groups of silkworm (*Bombyx mori*) by biochemical markers. *Insect Sci* 12: 87-94.
47. Nath BS, Suresh A, Varma BM, Kumar RP (1997) Changes in protein metabolism in hemolymph and fat body of the silkworm, *Bombyx mori* (Lepidoptera: Bombycidae) in response to organophosphorus insecticides toxicity. *Ecotoxicol Environ Saf* 36: 169-173.
48. Rugumamu CP (2006) Varietal role in the management of the larger grain borer, *Prostephanus truncatus* (horn) in stored maize. *Tanzania Journal of Science* 32: 13-22.



- Journal of Anesthesia & Clinical Care  
Journal of Addiction & Addictive Disorders  
Advances in Microbiology Research  
Advances in Industrial Biotechnology  
Journal of Agronomy & Agricultural Science  
Journal of AIDS Clinical Research & STDs  
Journal of Alcoholism, Drug Abuse & Substance Dependence  
Journal of Allergy Disorders & Therapy  
Journal of Alternative, Complementary & Integrative Medicine  
Journal of Alzheimer's & Neurodegenerative Diseases  
Journal of Angiology & Vascular Surgery  
Journal of Animal Research & Veterinary Science  
Archives of Zoological Studies  
Archives of Urology  
Journal of Atmospheric & Earth-Sciences  
Journal of Aquaculture & Fisheries  
Journal of Biotech Research & Biochemistry  
Journal of Brain & Neuroscience Research  
Journal of Cancer Biology & Treatment  
Journal of Cardiology & Neurocardiovascular Diseases  
Journal of Cell Biology & Cell Metabolism  
Journal of Clinical Dermatology & Therapy  
Journal of Clinical Immunology & Immunotherapy  
Journal of Clinical Studies & Medical Case Reports  
Journal of Community Medicine & Public Health Care  
Current Trends: Medical & Biological Engineering  
Journal of Cytology & Tissue Biology  
Journal of Dentistry: Oral Health & Cosmesis  
Journal of Diabetes & Metabolic Disorders  
Journal of Dairy Research & Technology  
Journal of Emergency Medicine Trauma & Surgical Care  
Journal of Environmental Science: Current Research  
Journal of Food Science & Nutrition  
Journal of Forensic, Legal & Investigative Sciences  
Journal of Gastroenterology & Hepatology Research  
Journal of Gerontology & Geriatric Medicine  
Journal of Genetics & Genomic Sciences  
Journal of Hematology, Blood Transfusion & Disorders  
Journal of Human Endocrinology  
Journal of Hospice & Palliative Medical Care  
Journal of Internal Medicine & Primary Healthcare  
Journal of Infectious & Non Infectious Diseases  
Journal of Light & Laser: Current Trends  
Journal of Modern Chemical Sciences  
Journal of Medicine: Study & Research  
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology  
Journal of Neonatology & Clinical Pediatrics  
Journal of Nephrology & Renal Therapy  
Journal of Non Invasive Vascular Investigation  
Journal of Nuclear Medicine, Radiology & Radiation Therapy  
Journal of Obesity & Weight Loss  
Journal of Orthopedic Research & Physiotherapy  
Journal of Otolaryngology, Head & Neck Surgery  
Journal of Protein Research & Bioinformatics  
Journal of Pathology Clinical & Medical Research  
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance  
Journal of Physical Medicine, Rehabilitation & Disabilities  
Journal of Plant Science: Current Research  
Journal of Psychiatry, Depression & Anxiety  
Journal of Pulmonary Medicine & Respiratory Research  
Journal of Practical & Professional Nursing  
Journal of Reproductive Medicine, Gynaecology & Obstetrics  
Journal of Stem Cells Research, Development & Therapy  
Journal of Surgery: Current Trends & Innovations  
Journal of Toxicology: Current Research  
Journal of Translational Science and Research  
Trends in Anatomy & Physiology  
Journal of Vaccines Research & Vaccination  
Journal of Virology & Antivirals

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>