

## Research Article

# Contribution of Climate Change to Soil Carbon and Nitrogen Storages in Three Parent Materials in Akwa Ibom State, Nigeria

I D Edem\* and Uduak C Udoinyang

Department of Soil Science and Land Resources Management, University of Uyo, Uyo, Nigeria

### Abstract

Most agricultural soils contain only a small proportion of carbon and nitrogen, but this small amount plays a very large role in soil quality. However, the estimation of carbon and nitrogen storages in rain forest area of Nigeria has not been adequately documented. A study was conducted to examine the effects of climate change on the Carbon and Nitrogen storages of soils derived from Coastal Plain Sands (CPS), Beach Ridge Sands (BRS) and Sand Stone/Shale (SSS) parent materials in Akwa Ibom State. Secondary soil data of organic carbon, and bulk density as well as climatic data of 2004 to 2014 were used. Organic C stock and SON were calculated as product of soil organic carbon (or total N) contents, bulk density and soil depth. Data were analysed using descriptive statistics, regression analysis and ANOVA at  $\alpha_{0.05}$ . Organic C stocks on the surface (0-15 cm) soil depth were  $44.13 \pm 15.91$ ,  $43.89 \pm 17.39$  and  $52.40 \pm 18.02$  Mg C ha<sup>-1</sup> for CPS, BRS and SSS, respectively. The corresponding values for subsurface depth (15-30 cm) were  $28.30 \pm 9.58$ ,  $24.83 \pm 8.93$  and  $26.37 \pm 5.74$  Mg C ha<sup>-1</sup>. However, N stocks at 0-15 cm soil depth were  $3.37 \pm 0.90$ ,  $3.19 \pm 0.94$  and  $3.56 \pm 1.00$  Mg N ha<sup>-1</sup> for CPS, BRS and SSS, respectively. Whereas in the 15-30 cm soil depth, SON were  $2.07 \pm 0.55$ ,  $2.00 \pm 0.58$  and  $2.04 \pm 0.56$  Mg N ha<sup>-1</sup> for the respective parent materials. Highly significant SOC stock of  $78.90$  Mg C ha<sup>-1</sup> occurred in 2005 in SSS, while the least stock of  $23.71$  Mg C ha<sup>-1</sup> occurred in 2007 CPS. Across the parent materials, organic C stocks increased as rainfall decreases. Whereas, N stocks increased with decrease in relative humidity. Change in climate significantly affected soil organic carbon and nitrogen storages in Uyo, Nigeria.

**Keywords:** Carbon and nitrogen stocks, Climate change, Parent materials, Rainfall

\*Corresponding author: I D Edem, Department of Soil Science and Land Resources Management, University of Uyo, Uyo, Nigeria, Tel: +234 8027031426; E-mail: dennis.edem@gmail.com

**Citation:** Edem ID (2019) Contribution of Climate change to Soil Carbon and Nitrogen Storages in Three Parent Materials in Akwa Ibom State, Nigeria. J Atmos Earth Sci 3: 008.

**Received:** February 02, 2019; **Accepted:** February 20, 2019; **Published:** March 06, 2019

**Copyright:** © 2019 Edem ID, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Introduction

Climate change refers to a long-term shift in weather conditions. It is measured by changes in a variety of climatic indicators such as temperature, precipitation, and wind. Climate change can be the result of natural processes and/or human activity [1,2]. Like all soils the world over, ultisol found in Akwa Ibom State, Nigeria are strongly affected by climatic gradients in temperature and moisture which in turn affects the storage of Soil Organic Carbon (SOC) and Soil Organic Nitrogen (SON) content. Therefore, to maintain stable Carbon (C) and Nitrogen (N) stocks within the soils of this vast region, it requires a balance between plant primary productivity and decomposition. This balance is affected by Carbon dioxide (CO<sub>2</sub>), temperature and soil moisture, which is further modified by soil texture [3]. Vegetation drives maximum C input to the soil and varies across ecosystems primarily with moisture gradients [4,5]. Consequently, SOC and SON stocks are also highly related to temperature and precipitation gradients [6,7]. Temperature and moisture control Soil Organic Matter (SOM) decomposition and CO<sub>2</sub> respiration driving both global and regional patterns of SOC.

Soils with greater silt and clay content typically have greater SOC stocks due to physical protection of organic minerals [8,9]. Climate change scenarios predict a 2.2°C to 3.6°C (4°F to 6.5°F) increase in temperature and increased variability in precipitation across most of the United States by 2040 to 2059 according to Marschner et al., , will likely alter the distribution and composition of plant biomass. Initial plant responses to changes in temperature and CO<sub>2</sub> observed by Hinzman may not reflect long-term SOC dynamics since increased plant C inputs can lead to increased decomposition [10,11].

In Nigeria, it is obvious that climate change is equally resulting in variations in temperature and precipitations with its attendant effects on the vegetation [12]. However, evidence of the effects of changes in SOC and SON stocks documentation such as those ascribed to climate change in Ultisol is scanty. This study therefore, attempts to examine the effects of climate change on the soil organic carbon and nitrogen storage of soils on three parent materials in Akwa Ibom State, Nigeria over a ten-year period.

### Materials and Methods

#### Site information

The study was conducted in Akwa Ibom State, Southeastern Nigeria. The State is located between latitudes 4°30' and 5°30' N and longitudes 7°27' and 8°27'E and is characterised by a uniformly hot wet and humid tropical climate [13]. The climate is characterised by two seasons; the wet and dry seasons. The wet or rain season lasts between the months of April and October, during which the rains are heavy and of high intensity. The pattern of rainfall is bimodal with peaks in July and September. The mean annual rainfall varies from 2500 - 4000 mm. The dry season lasts from November to March. The average daily temperature varies from 26°C to 28°C. Solar radiation falls within 6 to 15 mm/day. Relative humidity ranges from 75 to 95%, while evapotranspiration ranges from 4.11 to 4.95 mm, partly because of the high values of insolation and temperature.

## Vegetation and land use

The hot humid climate favours the luxuriant tropical rainforest [14], which has however been almost completely replaced by secondary forest of predominantly wild palm trees, woody shrubs and various grasses undergrowth. Poor management and improper land use, coupled with human population pressure has led to forest depletion with attendant soil degradation and low productivity [15,16]. Mangrove ecosystem extends into the estuaries and flood plains of the Imo River at Ikot Abasi, the Qua River at Eket and the Cross River at Mbo and Oron. Large tracts of Riverine Swamp and flood plain environments with wetland characteristics flank the Qua Iboe River valley through Etinan and Abak L.G.A. Farming is very restricted on the beach ridge sands, while agriculture and road network are adversely affected on the narrow-crested sandstone ridges with steep-sided hills and valley which cover much of Itu, Ibiono Ibom, Ikono and Ini L.G.A. The predominant land use is the bush fallow cropping system operated with low productive farming tools. This cropping system leaves more than 80% of the soil surface bare. And this exposed the soil to high intensity rainfall erosivity [16] that is common to this area.

## Data collection

Secondary soil data spanning from 2004 to 2014 were used. Data were collated from three major parent materials in the area viz: Coastal Plain Sands (CPS), Sand Stone Shale (SSS) and Beach Ridge Sands (BRS). Soil data for each of the three parent materials since 2004 were considered for the study and soil samples of 0 - 15 and 15 - 30 cm depths were used. The soil data were pulled together and were considered as representative of each parent materials for the years under study.

## Procedure for determining Soil Organic Carbon (SOC) stock

Calculating SOC stocks for a given depth consists of summing SOC Stocks by layer determined as a product of bulk density ( $D_b$ ) [17], Organic Carbon (OC) concentration, and layer thickness for an individual profile with  $n$  layers [18]. Organic carbon stock was determined by the following equation [19].

$$SOC = \sum_{i=1}^n D_b * C_i * D_i * 1000 * 0.0001$$

Where, SOC = soil carbon ( $Mg\ ha^{-1}$ );  $D_b$  = Bulk density ( $Mg\ m^{-3}$ );  $C_i$  = Organic carbon content of depth  $i$  ( $g\ kg^{-1}$ );  $D_i$  = soil depth  $i$

## Procedure for determining Soil Nitrogen (SON) Stock

Soil Nitrogen carbon stock was determined by the following equation:

$$SON = \sum_{i=1}^n D_b * N_i * D_i * 1000 * 0.0001$$

Where, SON = soil nitrogen ( $Mg\ ha^{-1}$ );  $D_b$  = Bulk density ( $Mg\ m^{-3}$ );  $N_i$  = soil total N of depth  $i$  ( $g\ kg^{-1}$ );  $D_i$  = soil depth  $i$

## Statistical analysis

Data were analysed using descriptive statistics and ANOVA at  $\alpha_{0.05}$ . The regression analysis was used to determine the association of the climatic variables (rainfall, temperature and relative humidity) with SOC and SON over the periods.

## Results and Discussion

Results of the soil properties affected by change in climatic factors are summarised in Tables 1-3. Mean bulk density (BD) on the surface soil of CPS, BRS and SSS were  $1.47 \pm 0.05$ ,  $1.46 \pm 0.07$  and  $1.56 \pm 0.05\ Mg\ m^{-3}$ , respectively. The corresponding values in the sub surface were  $1.59 \pm 0.06$ ,  $1.58 \pm 0.05$  and  $1.66 \pm 0.04\ Mg\ m^{-3}$ , (Table 1). The trend of soil compaction on both soil surfaces among the three parent materials was  $SSS > CPS > BRS$ . The shale which associated with sandstone consistently characterised with high bulk density. Relatively, bulk density was moderately high on the surface soil compared with subsurface depth. This according to Agbede and Hamarshid is attributed to the organic matter content on the surface soil that promotes good soil aggregation, thus lowering bulk density of the soil [20-23].

Year	Bulk Density ( $Mg\ m^{-3}$ )					
	0 - 15 cm			15 - 30 cm		
	CPS	BRS	SSS	CPS	BRS	SSS
2004	1.43	1.48	1.53	1.58	1.57	1.59
2005	1.43	1.53	1.58	1.69	1.62	1.62
2006	1.51	1.46	1.57	1.56	1.66	1.66
2007	1.55	1.38	1.58	1.68	1.55	1.63
2008	1.48	1.45	1.49	1.55	1.52	1.71
2009	1.54	1.45	1.47	1.61	1.56	1.64
2010	1.42	1.36	1.56	1.52	1.49	1.69
2011	1.46	1.57	1.57	1.59	1.62	1.66
2012	1.45	1.53	1.58	1.57	1.59	1.64
2013	1.38	1.48	1.61	1.54	1.56	1.71
2014	1.52	1.33	1.66	1.63	1.65	1.7
Mean	1.47	1.46	1.56	1.59	1.58	1.66
SD	0.05	0.07	0.05	0.06	0.05	0.04
CV	3.7	5.1	3.4	3.5	3.4	2.4

**Table 1:** Bulk density of soils of three parent materials in Akwa Ibom State (2004 to 2014).

Organic C content of  $20.13 \pm 7.50$ ,  $19.99 \pm 7.59$  and  $22.29 \pm 7.48\ g\ kg^{-1}$  on the surface soil of CPS, BRS and SSS were significantly higher compared to  $11.89 \pm 4.23$ ,  $10.48 \pm 3.76$  and  $10.57 \pm 2.19\ g\ kg^{-1}$  in the subsurface soil, respectively. The corresponding values for soil nitrogen content were  $1.53 \pm 0.40$ ,  $1.45 \pm 0.36$  and  $1.51 \pm 0.41\ g\ kg^{-1}$  on the surface, while soil nitrogen content in the subsurface were  $0.86 \pm 0.22$ ,  $0.84 \pm 0.22$  and  $0.82 \pm 0.23\ g\ kg^{-1}$ , respectively. However, SSS parent material recorded significantly high SOC than the other parent materials on the surface, whereas high SOC in the subsurface was noticed in CPS parent material. This result is consistent with the earlier report of Bai et al., on soil organic carbon stock changes at varying depths [24]. Organic carbon was highly varied (>35%) in both the 0 - 15 and 15 - 30 cm soil depths of CPS and BRS, but moderately varied in both depth of SSS. Harris et al., [25] noted this high variable distribution of SOC at the field scale both horizontally and vertically. Accordingly, level of SOC variability was attributed to differences in vegetative cover, geomorphic soil position and thermal diffusivity of the soil.

Years	Organic Carbon (g kg <sup>-1</sup> )					
	0 - 15 cm			15 - 30 cm		
	CPS	BRS	SSS	CPS	BRS	SSS
2004	13.04	15.23	23.57	9.46	7.45	10.38
2005	24.67	20.56	33.29	14.57	15.83	7.13
2006	15.65	16.23	32.44	12.25	7.45	12.55
2007	10.2	34.32	20.45	9.21	16.98	11.34
2008	28.22	18.48	18.55	18.2	11.67	14.13
2009	15.21	12.98	14.87	7.43	5.78	9.33
2010	27.49	15.34	31.39	20.6	10.33	10.64
2011	22.9	32.13	16.74	11.5	14.33	13.12
2012	33.23	26.34	10.75	8.76	7.49	7.33
2013	16.9	13.67	18.89	9.89	8.83	10.34
2014	13.87	14.56	24.23	8.95	9.12	10.03
Mean	20.13	19.99	22.29	11.89	10.48	10.57
SD	7.5	7.57	7.48	4.23	3.76	2.19
CV%	37.3	37.9	33.9	35.5	35.9	20.7

**Table 2:** Organic carbon concentration in soils of three parent materials in Akwa Ibom State (2004 to 2014).

Year	Total Nitrogen (g kg <sup>-1</sup> )					
	0 - 15 cm			15 - 30 cm		
	CPS	BRS	SSS	CPS	BRS	SSS
2004	1.38	1.43	1.89	0.91	0.67	0.82
2005	1.21	2.22	1.05	0.84	0.83	0.59
2006	1.1	1.26	1.43	0.69	1.34	0.66
2007	2.32	1.06	2.33	1.21	0.88	1.43
2008	1.33	1.38	1.08	0.88	0.76	0.62
2009	1.32	1.37	1.26	0.68	0.79	0.83
2010	1.36	1.22	1.45	0.66	0.69	0.69
2011	2.01	2.02	1.52	1.32	1.17	0.89
2012	1.55	1.54	1.19	0.78	0.77	0.71
2013	2.01	1.23	2.04	0.83	0.66	0.89
2014	1.25	1.18	1.42	0.71	0.68	0.88
Mean	1.53	1.45	1.51	0.86	0.84	0.82
SD	0.4	0.36	0.41	0.22	0.22	0.23
CV	26	24.9	27	25	26.2	28.2


**Table 3:** Total nitrogen concentration in soils of three parent materials in Akwa Ibom State (2004 to 2014).

### Annual Trends of Climatic Elements in Akwa Ibom State from


The respective yearly rainfall and temperature data are presented in Figures 1 and 2. Significantly high annual rainfall of 319.2 mm in the study area was recorded in 2012 while the least annual rainfall of 153.4 mm was recorded in 2008. Figure 1 revealed fluctuation in rainfall pattern with stupendous rising in 2012 and significantly drop in 2013 and 2014. Although, the rainfall pattern in 2013 and 2014 were significantly higher compared to the periods preceding 2012.

On the other hand, high annual temperature above 26.5 °C was constantly recorded between 2004 and 2008. Conversely, the high temperature regime dropped to 21.8 °C in 2009 and subsequently increased above 26 °C in 2010 and climaxed at 27.5 °C in 2011.

From the forgoing, year 2011 was the hottest period and the peak period for water demand for decomposition of organic materials.



**Figure 1:** Monthly rainfall pattern in Akwa Ibom State (2004 - 2014).



**Figure 2:** Monthly temperature variations in Akwa Ibom State (2004 - 2014).

Organic C stocks of 44.13±15.91, 43.89±17.39 and 52.40±18.02 Mg C ha<sup>-1</sup> on the surface soil depth of CPS, BRS and SSS, respectively, were significantly higher than 28.30±9.58, 24.83±8.93 and 26.37±5.74 Mg C ha<sup>-1</sup> recorded in the subsurface soil depth (Table 4). On the surface soil, organic C stocks decreased in the order of SSS > CPS > BRS. Whereas, in the subsurface depth, it decreased in the order of CPS > SSS > BRS. Consistently, BRS stored the lowest C content in the soil. According to Hashamoto et al., differences in biomass production, cropping systems, land use types and soil types could greatly contribute to the variation in SOC stock at an annual time scale [26].

Similarly, total N stocks of 3.37±0.90, 3.19±0.94 and 3.56±1.00 Mg N ha<sup>-1</sup> on the surface soil depth of CPS, BRS and SSS, respectively, were significantly higher than 2.07±0.55, 2.00±0.58 and 2.04±0.56 Mg N ha<sup>-1</sup> recorded in the subsurface soil depth of the corresponding parent material (Table 5). Total N stocks in the 0 - 15 cm depth decreased in the order of SSS > CPS > BRS while in the 0 - 30 cm depth, it decreased from CPS > SSS > BRS. The highest total N stocks of 5.22 Mg N ha<sup>-1</sup> recorded on the surface soil of SSS was in 2007, while the least storage of 2.19 Mg N ha<sup>-1</sup> was recorded in CPS of the same year (Table 5).

### Effect of Climate on Carbon and Nitrogen Storages

The effects of climate on C and N storage among CPS, BRS and SSS parent materials were assessed using rainfall and soil temperature. In CPS, BRS and SSS, organic C stock increased with reduction in the amount of rainfall by 30.1, 12.6 and 54.2%, respectively. Organic C stock was highest in SSS parent material and least in the

BRS. Hector et al., equally found an increased SOC to the total SOC pool in low rainfall soils [16]. On the other hand, Lal et al., reported that soil organic C levels increased with a corresponding increase in mean annual precipitation [27]. This is true for soils with coarse textures soil such as BRS. The significant increase in soil C stock with decreasing rainfalls especially in SSS soil, is therefore, linked to moderation of the biomass loss effects with high decomposition rates in fine textured soil. However, the relationships of organic C stocks and rainfall for CPS (0.301) and BRS (0.126) were not significant. The rainfall effects on N stock in CPS and SSS were significant and directly related, whereas the effect in BRS was inversely related. Across the parent materials, 200 mm of monthly rainfall was found to be the peak rainfall amount that contributed to the storage of large quantity of N (4.5 Mg N ha<sup>-1</sup>) in the soil. However, temperature with highest N stock in CPS, BRS and SSS were 26 (4.5 Mg N ha<sup>-1</sup>), 27 °C (4.0 Mg N ha<sup>-1</sup>) and 28 °C (4.7 Mg N ha<sup>-1</sup>), respectively.

## Conclusion

It is evident from the study that organic C stock in the surface soil decreased significantly among the parent materials in the order of sand stone/shale > coastal plain sand > beach ridge sand. However, soil nitrogen stock did not follow any definite trend as the organic C stocks obtained in the surface depth. Comparing the three parent materials, organic C stocks in coastal plain sand and sand stone/shale only increased with decrease in rainfall amount as the temperature increases, whereas in beach ridge sand parent material, N stock increased as rainfall amount decreases. Among the years examined, highest total N stock was obtained in 2007 from sand stone/shale parent material and the least from coastal plain sand. High temperature rate reduced the level of organic carbon and nitrogen storages. Therefore, climatic factors contribute significantly to the variability of soil carbon and nitrogen storages among the three parent materials in Akwa Ibom State, Nigeria.

## References

- Cleland EE, Chuine I, Menzel A, Mooney HA, Schwartz MD (2007) Shifting Plant Phenology in Response to Global change. *Trends Ecol Evol* 22: 357-365.
- IPCC (International Panel on Climate Change) (2001) Impacts, Adaptation and vulnerability, Summary for Policy Makers. Cambridge University Press, Cambridge, UK.
- Emmett BA, Beier C, Estiarte M, Tietema A, Kristensen HL, et al. (2004) The Response of Soil Processes to Climate Change: Results from Manipulation Studies of Shrublands across an Environmental Gradient. *Ecosystems* 7: 625-637.
- Doret SH, Shkesby RA, Walsh RP (2009) Soil Water Repellency. Its Causes, Characteristics and Hydrogeomorphological Significance. *Earth-Science Reviews* 51: 33-65.
- Edem ID, Edem SO (2008) Insitu Erosion Variability Management under Vetiver Hedges in Alfisol Assessment of Management Control on Loss and Runoff in the Field with Multi-slot Device. LAMBERT Academic Publishing GmbH and Co., KG. Saarbrücken, Germany.
- Brady NC, Weil R (2002) The Nature and Properties of Soils (13<sup>th</sup> edn). Prentice Hall. Upper Saddle River, New Jersey, USA, Pg no: 960.
- MacDonald NW, Zak DR, Pregitzer KS (1995) Temperature Effects on Kinetics of Microbial Respiration and Net Nitrogen and Sulfur Mineralization. *Soil Sci Soc Am J* 59: 233-240.
- Cookson WR, Osman M, Marschner P, Abaye DA, Clark I, et al. (2007) Controls on soil Nitrogen Cycling and Microbial Community Composition across Land use and Incubation Temperature. *Soil Biology and Biochemistry* 39: 744-756.
- Frey SD, Lee J, Melillo JM, Six J (2013) The Temperature Response of Soil Microbial Efficiency and its Feedback to Climate. *Nature of Climate Change* 3: 395-398.
- Marschner B, Bredow A (2002) Temperature effects on release and ecologically relevant properties of dissolved organic carbon in sterilized and biologically active soil samples. *Soil Biology and Biochemistry* 34: 459-466.
- Hinzman LD (2005) Evidence and Implications of Recent Climate Change in Northern Alaska and other Arctic Regions. *Climatic Change* 72: 251-298.
- Watter PT (1990) Seasonal Rainfall Characteristics and daily Rainfall. *Meteorology Society* 88: 90-95.

Year	Organic Carbon Storage (Mg C ha <sup>-1</sup> )					
	0 - 15 cm			15 - 30 cm		
	CPS	BRS	SSS	CPS	BRS	SSS
2004	27.98	33.81	54.09	22.42	17.54	24.75
2005	52.92	47.19	78.9	36.95	38.47	17.33
2006	35.45	35.54	76.4	28.66	18.55	31.25
2007	23.72	71.04	48.47	23.21	39.48	27.73
2008	62.66	40.19	41.46	42.32	26.61	36.25
2009	35.14	28.23	32.79	17.94	13.53	22.95
2010	58.55	31.29	73.45	46.97	23.09	26.98
2011	50.16	75.67	39.42	27.43	34.82	32.67
2012	72.27	60.45	25.48	20.64	17.86	18.03
2013	34.98	30.35	45.62	22.85	20.66	26.52
2014	31.62	29.05	60.33	21.88	22.57	25.58
Mean	44.13	43.89	52.4	28.3	24.83	26.37
SD	15.91	17.39	18.02	9.58	8.93	5.74
CV	36.05	39.62	34.38	33.86	35.97	21.77

Table 4: Organic carbon stock in soil from different parent materials.

Year	Total Nitrogen Storage (Mg N ha <sup>-1</sup> )					
	0 - 15 cm			15 - 30 cm		
	CPS	BRS	SSS	CPS	BRS	SSS
2004	2.96	3.17	4.34	2.16	1.58	1.96
2005	2.6	5.09	2.49	2.13	2.02	1.43
2006	2.49	2.76	3.37	1.61	3.34	1.64
2007	5.39	2.19	5.52	3.05	2.05	3.5
2008	2.95	3	2.41	2.05	1.73	1.59
2009	3.05	2.98	2.78	1.64	1.85	2.04
2010	2.9	2.49	3.39	1.5	1.54	1.75
2011	4.4	4.76	3.58	3.15	2.84	2.22
2012	3.37	3.53	2.82	1.84	1.84	1.75
2013	4.16	2.73	4.93	1.92	1.54	2.28
2014	2.85	2.35	3.54	1.74	1.68	2.24
Mean	3.37	3.19	3.56	2.07	2	2.04
SD	0.9	0.94	1	0.55	0.58	0.56
CV	26.66	29.53	28.1	26.59	28.79	27.59

Table 5: Mean total N stocks of soils from different parent materials.

13. Ogban PI, Ekerette (2001) Physical and Chemical Properties of the Coastal Plain Sand Soils of South-eastern Nigeria. *Nigerian Journal of Soil Resources* 2: 6-14.
14. Hills RC, Morgan JH (1981) An Interactive Approach to the Analysis of Rainfall records for Agricultural Purposes. *Experimental Agriculture* 17: 1-16.
15. Puthen W (1994) *Agriculture, Environment, Climate and Health, Sustainable Development in 21<sup>st</sup> Century*. University of Minnesota Press, Minneapolis, USA.
16. Hector FC, Aimee TC, Emily EA, Richard JN, Christopher WS (2009) Soil microbial community responses to multiple experimental climate change drivers. *Journal of Applied and Environmental Microbiology* 76: 999-1007.
17. Grossman RB, Reinsch TG (2002) Bulk Density and linear Extensibility Core Method. In: Dane JH, Topp GC, (eds.). *Method of Soil Analysis Part 4, Physical Methods*, Madison Wisconsin, USA. Soil Science Society of America Journal 208-228.
18. Bray RH, Kurtz LT (1945) Determination of Total Organic and Available forms of Phosphorus in Soils. *Soil Science* 59:39-45.
19. Brahim N, Bernoux M, Blavet D, Gallali T (2010) Tunisian Soil Organic Carbon Stocks. *International Journal of Soil Science* 5: 34-40.
20. Agbede OO (2009) *Understanding the Soil and Plant Nutrition*. Petra Digital Press, Nigeria.
21. Hamarshid NH (2010) Effect of Soil Texture on Chemical Composition, Microbial Population and Carbon Mineralization in Soil. *Egypt J Exp Biol* 6: 59-64.
22. Balsler TC, Kinzig AP, Firestone MK (2005) Linking Soil Microbial Communities and Ecosystem Functioning. In: Kinzig AP, Pacala SW, Tilman D (eds.). *The Functional Consequences of Biodiversity: Empirical Process and Theoretical Extensions*. Princeton University Press, New Jersey, USA.
23. Mbah CN, Anikwe MAN, Onweremadu EU, Mbagwu JSC (2007) Soil organic matter and carbohydrate contents of a dystric leptosol under organic waste management and their role in structural stability of soil aggregates. *Int J Soil Sci* 2: 268-277.
24. Bai ZG, Dent DL, Olsson L, Schaepman ME (2008) Proxy global assessment of land degradation. *Soil Use Manage* 24: 223-234.
25. Harris D, Pacovsky RS, Paul EA (1985) Carbon Economy of Soybean-rhizobium-glomus Association. *New Phytologist* 101: 427-440.
26. Hashimoto S, Suzuki M (2004) The impact of forest clear-cutting on soil temperature: A comparison between before and after cutting and between clear-cut and control sites. *Journal of Forest Research* 9: 125-132.
27. Kimble JM, Follett RF, Cole CV (1998) The potential of USA cropland to sequester carbon and mitigate the greenhouse effect. *Ann Arbor Press, Chelsea, MI, USA*, Page no: 128.



- Journal of Anesthesia & Clinical Care
- Journal of Addiction & Addictive Disorders
- Advances in Microbiology Research
- Advances in Industrial Biotechnology
- Journal of Agronomy & Agricultural Science
- Journal of AIDS Clinical Research & STDs
- Journal of Alcoholism, Drug Abuse & Substance Dependence
- Journal of Allergy Disorders & Therapy
- Journal of Alternative, Complementary & Integrative Medicine
- Journal of Alzheimer's & Neurodegenerative Diseases
- Journal of Angiology & Vascular Surgery
- Journal of Animal Research & Veterinary Science
- Archives of Zoological Studies
- Archives of Urology
- Journal of Atmospheric & Earth-Sciences
- Journal of Aquaculture & Fisheries
- Journal of Biotech Research & Biochemistry
- Journal of Brain & Neuroscience Research
- Journal of Cancer Biology & Treatment
- Journal of Cardiology & Neurocardiovascular Diseases
- Journal of Cell Biology & Cell Metabolism
- Journal of Clinical Dermatology & Therapy
- Journal of Clinical Immunology & Immunotherapy
- Journal of Clinical Studies & Medical Case Reports
- Journal of Community Medicine & Public Health Care
- Current Trends: Medical & Biological Engineering
- Journal of Cytology & Tissue Biology
- Journal of Dentistry: Oral Health & Cosmesis
- Journal of Diabetes & Metabolic Disorders
- Journal of Dairy Research & Technology
- Journal of Emergency Medicine Trauma & Surgical Care
- Journal of Environmental Science: Current Research
- Journal of Food Science & Nutrition
- Journal of Forensic, Legal & Investigative Sciences
- Journal of Gastroenterology & Hepatology Research
- Journal of Gerontology & Geriatric Medicine
- Journal of Genetics & Genomic Sciences
- Journal of Hematology, Blood Transfusion & Disorders
- Journal of Human Endocrinology
- Journal of Hospice & Palliative Medical Care
- Journal of Internal Medicine & Primary Healthcare
- Journal of Infectious & Non Infectious Diseases
- Journal of Light & Laser: Current Trends
- Journal of Modern Chemical Sciences
- Journal of Medicine: Study & Research
- Journal of Nanotechnology: Nanomedicine & Nanobiotechnology
- Journal of Neonatology & Clinical Pediatrics
- Journal of Nephrology & Renal Therapy
- Journal of Non Invasive Vascular Investigation
- Journal of Nuclear Medicine, Radiology & Radiation Therapy
- Journal of Obesity & Weight Loss
- Journal of Orthopedic Research & Physiotherapy
- Journal of Otolaryngology, Head & Neck Surgery
- Journal of Protein Research & Bioinformatics
- Journal of Pathology Clinical & Medical Research
- Journal of Pharmacology, Pharmaceutics & Pharmacovigilance
- Journal of Physical Medicine, Rehabilitation & Disabilities
- Journal of Plant Science: Current Research
- Journal of Psychiatry, Depression & Anxiety
- Journal of Pulmonary Medicine & Respiratory Research
- Journal of Practical & Professional Nursing
- Journal of Reproductive Medicine, Gynaecology & Obstetrics
- Journal of Stem Cells Research, Development & Therapy
- Journal of Surgery: Current Trends & Innovations
- Journal of Toxicology: Current Research
- Journal of Translational Science and Research
- Trends in Anatomy & Physiology
- Journal of Vaccines Research & Vaccination
- Journal of Virology & Antivirals

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>