

## Research Article

# First Trials on Hatchery and Larval Development of the Sea Cucumber, *Holothuria scabra* (Jaeger 1883) In Oman

Khalfan M Al-Rashdi<sup>1\*</sup>, Michel R Claereboudt<sup>2</sup> and Igor Eeckhaut<sup>3</sup>

<sup>1</sup>Aquaculture Center, Ministry of Agriculture and Fisheries Wealth, Muscat, Oman

<sup>2</sup>College of Agricultural and Marine Sciences, Sultan Qaboos University, Al-Khoudh, Oman

<sup>3</sup>Biology of Marine Organisms and Biomimetism, University of Mons-Hainaut, Mons, Belgium

### Abstract

Although sea cucumber *Holothuria scabra* has always been part of the traditional exploitation of the benthos in Mahout Bay of Arabian Sea, the foreign demand for the product and its high price have put increased pressure on the resource leading rapidly to overfishing. Aquaculture of this species has developed as a response to the overfishing problem but has not been yet studied in Oman. As a first step to evaluate its potential for aquaculture in Oman it was thus necessary to conduct hatchery trials. Four hatchery runs were conducted to evaluate the quality of the local broodstock, the response and efficiency of *in vitro* maturation and fertilization and the success of larval development and rearing. Collected animals of 200-600g were transported by road for 5 hours with zero evisceration to the Hatchery station. *In vitro* maturation and fertilization success of more than 90% were achieved using maturation inducing fractions method leading to the development of mature eggs and normal embryos larvae. *Auricularia* larval stages were completed within 15 days and fed normally on microalgae *Phaeodactylum tricorutum*, *Chaetoceros* sp. and *Nanochloropsis* sp. Settlement rate was reached 9% for pentactula larvae. However, a high mortality of 70% was observed during early larval development caused by ciliated protozoans and copepods attack. The culture of *Holothuria scabra* population in Oman is thus promising but further research is needed to ensure

\*Corresponding author: Khalfan M Al-Rashdi, Aquaculture Center, Ministry of Agriculture and Fisheries Wealth, Muscat, Oman, Tel: +968 24736618; E-Mail: khalfanrashdi@gmail.com

**Citation:** Al-Rashdi KM, Claereboudt MR, Eeckhaut I (2019) First Trials on Hatchery and Larval Development of the Sea Cucumber, *Holothuria scabra* (Jaeger 1883) In Oman. J Aquac Fisheries 3: 014

**Received:** January 14, 2019; **Accepted:** February 12, 2019; **Published:** February 26, 2019

**Copyright:** © 2019 Al-Rashdi KM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

higher survival rates, particularly in the early larval stages where adequate water filtration and sterilization is essential.

**Keywords:** Hatchery; *Holothuria scabra*; Larval development; Oman; Sea cucumber

### Introduction

Holothurians commonly known as sea cucumbers are marine invertebrates of significant importance for marine ecosystems and for a limited number of species as a valuable fishery generally for export. From the early 1990s, the populations of various species of sea cucumbers declined due to an increasing demand of Asian markets, particularly those of China [1]. Among these species, *Holothuria scabra* is considered as one of the most valuable sea cucumber for bêche-de-mer production and have been widely fished and overfished in the tropics [2]. It is currently valued at more than 1000 US\$ per kg dry weight [3], and the demand from the Asian market remains strong [4]. This increasing demand from Asian markets driving high price, an often inadequate management of the fishery and the biological characteristics of the species such as slow growth and easy to access shallow water habitat resulted globally in severe overfishing [5]. To avoid overexploitation of natural populations while maintaining a sustainable level of exports, aquaculture is considered as one of the potential solutions for *H. scabra* production [1,6]. Several countries have progressively developed a viable aquaculture production and in some cases have initiated restocking or sea ranching programs [1,7,8]. Although the sea cucumber *Holothuria scabra* has always been a small part of the traditional exploitation of the benthos in Mahout Bay of Arabian Sea [9], the foreign demand for the product overseas and its high price have put increased pressure on this resource locally leading rapidly to overfishing [10].

The natural spawning of this species in Oman has been recently documented and showed, as in many other locations, an annual reproductive cycle [11]. Natural maturation occurred in February and March followed by spawning in April and May, which is correlated to high sea surface temperature and precipitation as both may serve as exogenous synchronizing cues for gamete maturation and spawning [11]. Alarming signs of the of the Omani population of *H. scabra* overfishing have been observed with stock densities decreasing to less than 1 individual per ha [10], a regular decrease in the size of animal processed and a progressive switch to less valuable species in only a few years of exploitation. Therefore, restocking and stock enhancement in addition to fisheries management policies are needed to ensure the sustainability of this resource. This article documents the first attempts made to develop hatchery techniques for the Omani sea cucumber *Holothuria scabra* population as initial steps towards aquaculture development and stock enhancement programs.

### Materials and Methods


#### Culture site and system

The Aquaculture Research Unit of Sultan Qaboos University located in Al-Hail some 40 km north of the capital of the Sultanate,

Muscat, was used to conduct *Holothuria scabra* hatchery trials. Land based outdoors tanks were used to keep the broodstock of *H. scabra*. The tanks were continuously supplied with seawater extracted from a marine well at a salinity of 37 PSU and a temperature of 29°C. For the spawning and larval rearing experiments, the water was filtered using sand filter and 5µm particle filter and then passed through a UV sterilization 66 unit [12].

### Broodstock collection and transportation

In total 80 healthy animals weighing 200-600g were collected at 4 occasions from Mahout Bay of Al-Wusta Province by local fishermen and through SCUBA diving starting between September 2009 and November 2010 (Figure 1).



**Figure 1:** Location of the *Holothuria scabra* population used for the spawning experiments. Individuals were collected from various locations within Mahout Bay.

Collected animals were carefully handled to avoid evisceration, cleaned by gently washing the body with seawater and packed at low density of 6-8 individuals per bags in transparent plastic bags filled with 1-2L of sea water and inflated with pure oxygen. The bags were placed in cool-boxes containing dry ice to maintain a low temperature in the boxes during the 5 hours trip to Muscat. Upon arrival at the hatchery, the animals were acclimatized while still in the transport bags with ambient seawater and later transferred to clean tanks with running oxygenated sea water until experimentation.

### Induced maturation and fertilization

For this study, *In vitro* maturation and fertilization were conducted using an artificial maturation substance extracted from sea urchin oocytes [13]. It is referred as Maturation Inducing Fractions (MIF) and based on hatchery work in Madagascar [13,14]. Genitors were dissected through a longitudinal incision on the anterior part of the trivium to extract male and female gonads for the *in vitro* fertilization.

Body fluid was drained and the gonads of males and females were gently removed using forceps and placed in separate beakers with filtered sea water. Gonads were then cut separately into small pieces using scissors to release oocytes and sperm cells from gonadal tubules. The oocytes were then sieved on 60µm mesh with filtered sea water and placed in petri dishes at concentrations of 100 oocytes/ml. Maturation of the oocytes was induced with the MIF solution for 2 hours, and then rinsed with filtered sea water to remove MIF residuals. Fertilization was achieved by adding few drops of sperm suspension to the mature oocytes. The fertilized eggs were kept in a shallow basin without aeration until hatching of gastrulated embryos. Observations and photography of embryonic development stages were carried out on random samples every 30-60 minutes using a light microscope at 40X magnification.

### Larval development and rearing

Newly-hatched gastrula larvae were transferred into 100L tanks filled with 85L of 2-µm filtered and UV-treated seawater and provided with aeration through air diffusers. Feeding and regular water exchange commenced with development of *Auricularia* stage larvae from day-2 Post-Fertilization (PF). The water was changed by siphoning and the larvae collected onto a 60µm wet sieve. Tanks were refilled with 2-µm filtered and UV-treated seawater. This method of water exchange was continued daily for 20 days until larval settlement. The newly settled juveniles were maintained in a flow-thru system with 2-µm filtered and UV-treated seawater.

*H. scabra* larvae were fed daily from the second day PF with different types of micro-algae until settlement. Three mixed algae were cultured under laboratory conditions with artificial light and aeration. A temperate species *Phaeodactylum tricornutum* was cultured at low temperature from 19 to 24°C in the laboratory. *Chaetoceros* sp. and *Nanochloropsis* sp. were cultured both under laboratory and outdoor conditions (29-30°C) and fertilized with Guillard's F/2 medium. Initially, the developing larvae were fed twice daily with *P. tricornutum* at a final concentration of approximately 2500 cells per liter. On day 8 PF, the final concentration of *P. tricornutum* was increased to 5000 cells per liter. During the Doliolaria stage (a non-feeding stage), the feeding was temporarily suspended and resumed after metamorphosis (typically on day 20 PF). At this stage both microalgae (*Chaetoceros*, *Nanochloropsis* and *Phaeodactylum*) and particulate diet (dried and powdered *Sargassum* sp.) were added to favor settlement and feed the newly settled juveniles.

## Results

### Maturation induction and fertilization

All collected animals survived through 5 hours road transportation and were successfully transferred to holding tanks. For the artificial maturation experiments, all female genitors positively responded to the MIF technique and produced more than 90% normal mature oocytes after 2 hours of induction. A fertilization rate of 90% was also achieved on these artificially matured oocytes. A total of more than 400,000 larvae were produced over 4 independent *in vitro* fertilization trials and developed successfully into normal embryos and larvae (Table 1). Although 40-70% mortality was observed during larval development 9% of the produced larvae settled successfully as pentactula and started feeding.

Fertilization number	Dates	Broodstock size (g)		Fertilized oocytes	Produced larvae
		Male	Female		
F1	September 12th, 2009	287 (1)	287&346 (2)	160000	139200
F2	April 15th, 2010	245 (1)	457&375 (2)	140000	96000
F3	July 13th, 2010	319 (1)	320&425 (2)	85000	59,766
F4	November 17th, 2010	200&260 (2)	600&375 (2)	120000	105,400
Total				505000	400366

**Table 1:** Summary results for the 4 hatchery runs in 2009 and 2010. The number between brackets is the number of genitors used in the trials.

## Embryonic and larval development


Larval development was normal and follows the documented stages for *H. scabra* described in the literature (Figure 2). The mean diameter of fertilized eggs was  $150 \pm 3.0 \mu\text{m}$  and underwent cell division within 1.5 hours. They developed into blastula and gastrula around 12 and 16 hours Post Fertilization (PF), respectively. Early hatched *Auricularia* stage was observed around 36 hours PF, and continued to develop to mid-*Auricularia* larvae at day 6 PF. Late *Auricularia* larvae was observed by day-13 PF and transformed to non-feeding doliolaria larvae after completing 15 days PF. There was a mass mortality reached 70% during the first 3 days of rearing. Primary microscopic observation of the larvae, revealed an abundance of ciliate protozoan and copepods in the larval tanks. However, from the 4<sup>th</sup> day of development, survival rate remained almost stable. After 13 days, the larvae reached late *Auricularia* stage characterized by the appearance of lipid spheres at the edge of the cilia band. The first settlements were observed on day 20 PF and small juveniles began to appear on most trials on the bottom of the rearing tank 3 to 4 weeks PF. They appeared transparent and only visible using a light torch. Their size was estimated at 0.5 mm.

There was a high mortality during the first few days PF during all four trials. With the exception of trial 1 for which there were no microalgae to feed the larvae, from day 4 to day 15 mortality remained quite low. A second mortality was observed between day 15 and day 21 in all three remaining trials (Figure 3). During the daily microscopic observations of the larvae, abundant ciliate protozoans and copepods were observed in the larval tanks. However, after the 4<sup>th</sup> day PF, survival was stable.


## Discussion

Overfishing of *Holothuria scabra* went beyond just an ecological issue, it became a socio-economic problem as livelihoods and income of many coastal dwellers rely on its fishery.

There are several national management measures applied on this fishery, including total ban as the current case here in Oman; however, these seem to be insufficient for a sustainable use of the resources [6]. A sustainable and efficient solution so far to enhance production as well as depleted natural stock of this species is through developing aquaculture [5]. To ensure *H. scabra* aquaculture sustainability and affordability, the production of fertilizable eggs and larvae in large quantities and at any time is a key element [14]. Several methods have been described to induce *H. scabra* spawning, mainly thermal shock [15-17] alone or combination with water pressure and drying treatments [18]. Spawning can also be induced using chemical such as spirulina.



**Figure 2:** A. Immature oocyte; B. Mature oocyte; C. Fertilized egg; D. Two cell stage (1 h 30 mn); E. Four-cell stage (2 h); F. Multi-cells (3 h); G. Blastula (12 h); H. Gastrula (16 h); I. Newly hatched *Auricularia* (36 h); J. Early *Auricularia* (2.5 d); K. Mid *Auricularia* (6 d); L. Late *Auricularia* (13 d); M. Doliolaria (15); N. Early Pentactula with tentacles (20).



**Figure 3:** Survival rates of *H. scabra* over 4 hatchery runs and the mean of the 3 runs during 21 days of embryonic and larval development and rearing.

However, these methods may not be reliable to produce mature fertilizable oocytes outside the natural reproductive season of the organism [14,19]. Such critical point has been addressed through a new methodology based on *in vitro* maturation using artificial inducers [13,14] which recently have been identified [20]. The use of such maturation inducers allows holothurians farmers to obtain clean, fertilized eggs year-round. This method was tested in Oman *H. scabra* population and achieved more than 90% oocyte maturation and fertilization, which is consistent with Léonet et al. [13] and Eeckhaut et al. [14], results on Madagascar *H. scabra* populations. The method requires the sacrifice of several large adult specimens which may be difficult to justify for a locally endangered species, although these few specimens will produce 1000s of juveniles [21]. Despite larval developmental timing variations among various *H. scabra* populations, the embryos underwent normal development processes up to the pentactula stage over 21 days PF which is consistent with published values for larval development in *H. scabra* as it is reviewed by Junus et al. [3].

The high mortality observed during the first few days PF took place before feeding and is thus unlikely the result of a contamination by the feeding solutions. However, the abundant copepods and ciliates may be responsible for this temporary mortality [22]. Copepods attack larvae either directly or by repeated collisions causing bodily damage to the larvae [15]. Asha & Muthia [23] observed 65% mortality due to ciliates on the 9<sup>th</sup> day of rearing *H. spinifera* larvae. Since the water used for larval rearing was filtered and UV-treated, it is unlikely the source of the copepods and ciliates, but some of the equipment used to siphon the water, transfer the larvae from one container to another, the container themselves may have been contaminated by a concomitant fish hatchery and emphasize the need for high levels of training in sterilization and decontamination and a sepsis. Unfortunately, separating copepods from sea cucumber larvae is difficult as their sizes are similar [24]. To control copepod population, Pitt & Duy [15] recommended the application of trichlorofon (Dipterex), an insecticide which may be effective against some arthropod infestation. Alternative solutions using either naturally occurring substances or better management practices are needed [25]. In addition, optimization of the diet using locally available species is also necessary [26].

## Conclusion

The result presented here is a first insight on the artificial production of sea cucumbers in Oman. The potential to develop sea cucumber aquaculture in Oman is high because (i) *H. scabra*, a high commercial value, is still present in Oman particularly in shallow waters (ii) There are some natural locations ideally suited for sea ranching such as the Bay of Mahout, (iii) The human and the basic technological potential are already in place. For example, less than 1% of the surface of the Bay of Mahout Bay could support the sea ranching of 4 M of sea cucumbers *H. scabra* if it is supplied by an efficient farm having 50,000 m<sup>2</sup> of ponds and an efficient hatchery with 600,000 l of adequate tanks.

## Acknowledgment

This study was funded with thanks by the Agricultural and Fisheries Development Fund through research project; code No. 1/3/17, titled: Diversity, abundance and Aquaculture Potential of the Sea cucumber, *Holothuria scabra*, in the Sultanate of Oman. We give

grateful thanks to Mohamed Al-Wahaibi, Adel Al-Bulushi, Ali Al-Ajmi and Mohamed Al-hikmani for contributing to hatchery activities. We would like to acknowledge Indian Ocean Trepang (IOT) Company for the technical support and using their patent.

## References

1. Purcell SW, Hair CA, Mills DJ (2012) Sea cucumber culture, farming and sea ranching in the tropics: Progress, problems and opportunities. *Aquaculture* 368-369: 68-81.
2. Conand C (2004) Present status of world sea cucumber resources and utilisation: An international overview. *Advances in sea cucumber aquaculture and management* Pg no: 14-18.
3. Junus S, Kwong JP, Khoo G (2018) A review on the recent advances in the biology and aquaculture technology of *Holothuria scabra*. *Journal of Survey in Fisheries Sciences* 4: 5-25.
4. Purcell SW, Williamson DH, Ngaluafé P (2018) Chinese market prices of beche-de-mer: Implications for fisheries and aquaculture. *Marine Policy* 91: 58-65.
5. Hamel JF, Conand C, Pawson DL, Mercier A (2001) The sea cucumber *Holothuria scabra* (Holothuroidea: Echinodermata): Its biology and exploitation as Beche-de-mer. *Advances in Marine Biology* 41: 129-223.
6. Lovatelli A, Conand C, Purcell S, Uthicke S, Hamel JF, et al. (2004) Advances in sea cucumber aquaculture and management, FAO Fisheries Technical Paper 463, Rome, Italy.
7. Eriksson H, de la Torre-Castro M, Purcell S, Olsson P (2014) Lessons for resource conservation from two contrasting small-scale fisheries. *Ambio* 44: 204-213.
8. Watanabe S, Sumbing JG, Lebata-Ramos MJ (2014) Growth pattern of the tropical sea cucumber, *Holothuria scabra*, under captivity. *JARQ* 48: 457-464.
9. Al-Rashdi KM, Al-Busaidi SS, Al-Rassadi IH (2007) Status of the sea cucumber fishery in the Sultanate of Oman. *SPC Beche-de-mer Information Bulletin* 25: 17-21.
10. Al-Rashdi KM, Claereboudt MR (2010) Evidence of rapid overfishing of sea cucumbers in the Sultanate of Oman. *SPC Beche-de-mer Information Bulletin* 30: 10-13.
11. Al-Rashdi KM, Claereboudt MR (2018) Reproductive biology of the sea cucumber *Holothuria scabra* (Jaeger 1883) in Mahout Bay, Arabian Sea, Oman. *International Journal of Fisheries and Aquatic Studies* 6: 100-108.
12. Al-Rashdi KM (2012) Project final report on diversity, stock and aquaculture potential of sea cucumber in Oman. Ministry of Agriculture and Fisheries Wealth, Aquaculture Centre, Muscat, Sultanate of Oman. Pg no: 150.
13. Léonet A, Rasolofonirina R, Wattiez R, Jangoux M, Eeckhaut I (2009) A new method to induce oocyte maturation in holothuroids (Echinodermata). *Invertebrate Reproduction and Development* 53: 13-21.
14. Hair CA, Pickering TD, Mills DJ (2012) Asia-Pacific tropical sea cucumber aquaculture: Proceedings of an international symposium held in Noumea, New Caledonia, 15-17 February, 2011. *ACIAR Proceedings* 136: 1-209.
15. Pitt R, Nguyen DQD (2004) Breeding and rearing of the sea cucumber *Holothuria scabra* in Vietnam. *FAO Fisheries Tech Pap* 463: 333-346.
16. Duy NDQ (2010) Seed production of sandfish (*Holothuria scabra*) in Vietnam. *Aquaculture Department, Southeast Asian Fisheries Development Center, Iloilo, Philippines*.
17. Hair C, Kaure T, Southgate PC, Pickering T (2011) Potential breakthrough in hatchery culture of Sandfish *Holothuria scabra* by using algal concentrate as food. *SPC Bêche-de-mer Inf. Bull* 31: 60-61.

18. Agudo N (2007) Sandfish Hatchery Techniques. Australian Center for International Agricultural Research (ACIAR), Thynne St, Australia. Pg no: 42.
19. Battaglione SC, Seymour JE, Ramofafia C, Lane I (2002) Spawning induction of three tropical sea cucumbers, *Holothuria scabra*, *H. fuscogilva* and *Actinopyga mauritiana*. *Aquaculture* 207: 29-47.
20. Léonet et al. (2019) in press.
21. Purcell SW, Ngaluafé P, Foale SJ, Cocks N, Cullis BR, et al. (2016) Multiple factors affect socioeconomics and wellbeing of artisanal sea cucumber fishers. *PLoS One* 11: 0165633.
22. James BD (2004) Captive breeding of the sea cucumber, *Holothuria scabra*, from India. *FAO Fisheries Technical Paper* 463: 385-398.
23. Asha PS, Muthia P (2002) Spawning and larval rearing of sea cucumber *Holothuria (Theelothuria) spinifera* Theel. *SPC Beche-de-mer Information Bulletin* 16: 11-15.
24. James BD, Rajapandian ME, Basker BK, Gopinathan CP (1988) Successful induced spawning and rearing of the holothurian *Holothuria (Metriatyla) scabra* Jaeger at Tuticorin. *Marine Fisheries Infor Ser* 87: 30-33.
25. McLean E, Ibrahim FS, Al-Rashdi KM (2011) Better management practices for Omani aquaculture. Ministry of Agriculture & Fisheries Wealth, Aquaculture Centre, Muscat, Sultanate of Oman. Pg no: 80.
26. Militz AT, Leini E, Duy ND, Paul GS (2018) Successful large-scale hatchery culture of sandfish (*Holothuria scabra*) using micro-algae concentrates as a larval food source. *Aquaculture Reports* 9 :25-30.



Journal of Anesthesia & Clinical Care  
Journal of Addiction & Addictive Disorders  
Advances in Microbiology Research  
Advances in Industrial Biotechnology  
Journal of Agronomy & Agricultural Science  
Journal of AIDS Clinical Research & STDs  
Journal of Alcoholism, Drug Abuse & Substance Dependence  
Journal of Allergy Disorders & Therapy  
Journal of Alternative, Complementary & Integrative Medicine  
Journal of Alzheimer's & Neurodegenerative Diseases  
Journal of Angiology & Vascular Surgery  
Journal of Animal Research & Veterinary Science  
Archives of Zoological Studies  
Archives of Urology  
Journal of Atmospheric & Earth-Sciences  
Journal of Aquaculture & Fisheries  
Journal of Biotech Research & Biochemistry  
Journal of Brain & Neuroscience Research  
Journal of Cancer Biology & Treatment  
Journal of Cardiology: Study & Research  
Journal of Cell Biology & Cell Metabolism  
Journal of Clinical Dermatology & Therapy  
Journal of Clinical Immunology & Immunotherapy  
Journal of Clinical Studies & Medical Case Reports  
Journal of Community Medicine & Public Health Care  
Current Trends: Medical & Biological Engineering  
Journal of Cytology & Tissue Biology  
Journal of Dentistry: Oral Health & Cosmesis  
Journal of Diabetes & Metabolic Disorders  
Journal of Dairy Research & Technology  
Journal of Emergency Medicine Trauma & Surgical Care  
Journal of Environmental Science: Current Research  
Journal of Food Science & Nutrition  
Journal of Forensic, Legal & Investigative Sciences  
Journal of Gastroenterology & Hepatology Research  
Journal of Gerontology & Geriatric Medicine  
Journal of Genetics & Genomic Sciences  
Journal of Hematology, Blood Transfusion & Disorders  
Journal of Human Endocrinology  
Journal of Hospice & Palliative Medical Care  
Journal of Internal Medicine & Primary Healthcare  
Journal of Infectious & Non Infectious Diseases  
Journal of Light & Laser: Current Trends  
Journal of Modern Chemical Sciences  
Journal of Medicine: Study & Research  
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology  
Journal of Neonatology & Clinical Pediatrics  
Journal of Nephrology & Renal Therapy  
Journal of Non Invasive Vascular Investigation  
Journal of Nuclear Medicine, Radiology & Radiation Therapy  
Journal of Obesity & Weight Loss  
Journal of Orthopedic Research & Physiotherapy  
Journal of Otolaryngology, Head & Neck Surgery  
Journal of Protein Research & Bioinformatics  
Journal of Pathology Clinical & Medical Research  
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance  
Journal of Physical Medicine, Rehabilitation & Disabilities  
Journal of Plant Science: Current Research  
Journal of Psychiatry, Depression & Anxiety  
Journal of Pulmonary Medicine & Respiratory Research  
Journal of Practical & Professional Nursing  
Journal of Reproductive Medicine, Gynaecology & Obstetrics  
Journal of Stem Cells Research, Development & Therapy  
Journal of Surgery: Current Trends & Innovations  
Journal of Toxicology: Current Research  
Journal of Translational Science and Research  
Trends in Anatomy & Physiology  
Journal of Vaccines Research & Vaccination  
Journal of Virology & Antivirals  
Archives of Surgery and Surgical Education  
Sports Medicine and Injury Care Journal  
International Journal of Case Reports and Therapeutic Studies

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>