



Short Review

Whole Body Vibration Effectiveness in Diabetes Mellitus, Osteoporosis and Parkinson's Disease

Maria das Graças Rodrigues de Araujo^{1,3} and Marcelo Renato Guerino^{2,3*}

¹Kinesiotherapy and Manual Therapeutic Resources Laboratory - LACIRTEM, UFPE, Recife-PE, Brazil

²Electro-Thermo-Phototherapy Laboratory - LETER, UFPE, Recife-PE, Brazil

³Department of Physical Therapy of Federal University of Pernambuco, Recife-PE, Brazil

Abstract

Whole Body Vibration (WBV) intervention has been used to further improve muscle activity and the performance of various disorders. Some studies show that the complete activation of the muscle by WBV can lead to fatigue of the motor unit and, consequently, an increase in muscle strength. It is known that physical exercise promotes glycemic control in diabetic individuals and the beneficial effect is likely to be multifactorial, including increased energy expenditure. In addition, resistance and/or strength exercises can improve gait speed, balance and muscle strength in patients with type 2 diabetes and the elderly with osteoporosis. However, some of these patients, due to clinical conditions and/or possible complications, are unable or unwilling to perform the exercises at the intensities necessary to achieve satisfactory results. In this sense, the training of whole body vibration leads us to develop protocols complementary to a traditional exercise program.

WBV has the potential to stimulate the main factors associated with improved muscle performance such as amplitude and frequency, which in patients with Parkinson's disease can be direct protective effects or, in the case of higher intensities of WBV, they

can result in micro-injuries. Therefore, the types of vibration and its method of application, training intensity and exercise protocol must adapt to the specific characteristics of the dysfunctions of each group of participants.

Introduction

Vibration is a mechanical oscillation, periodic alteration of force, acceleration and displacement over time [1]. The notion that vibration can be beneficial is relatively new, therefore is important to quote that researcher's study of vibration effects in the neuronal sciences [2]. Sander [3] was the first author to recognize the therapeutic importance of the vibratory stimuli with Whedon [4], with intervention of the cardiovascular and musculoskeletal disorders. Nazarov and Spivak [5] were the first to apply vibration as a training modality for athletes and were first used therapeutically in the Soviet Union to prevent hypotonia in cosmonauts. Initially used in segmental form, it evolved to the vibration of the whole body with the use of machines, with overall effect [6,7].

We use the Whole Body Vibration (WBV) in the rehabilitation of various populations. The elderly are therefore possible beneficiaries of WBV because they have poor muscular performance and low bone quality as well as functional and structural modifications in the organism [8]. The physiological decay of the elderly adult may lead to a reduction in the ability to balance and an increased risk of falls becoming an important issue among the elderly [9]. Balance control requires adaptations in the individual support base, which changes with age; therefore, alternatives will be directed at reactions to external disturbances that were not anticipated during gait [10].

In our clinical practice as physiotherapist, we found elderly people with diabetes mellitus and severe changes in balance as well as functional repercussions in the lives of these individuals. However, there are studies that seek treatment options for this population through various types of exercises. Some studies show that WBV is considered an effective strategy to improve balance in patients with type 2 Diabetes (DM2). A study of fifty-five elderly patients with diabetic peripheral neuropathy concluded that short-term therapy with WBV was beneficial and significant in improving balance [11]. In the work of Del Pozo-Cruz et al., [12] fifteen patients with DM2 participated in the WBV program followed by other types of exercises and the results showed that WBV is a safe and well-tolerated strategy to improve balance. Yoosefinejad et al., [13] investigated the effects of one session of WBV on muscle strength and the balance of ten diabetic patients with peripheral neuropathy; in other study they worked with twenty diabetic patients, using WBV twice a week, for 6 weeks, concluding that this intervention increased strength and muscle balance in these patients [14]. A recent systematic review study with meta-analysis conducted by Gomes-Neto et al., [15] concluded that WBV can be a useful strategy in the management of symptoms and disabilities associated with type 2 diabetes. Work carried out in our Research Laboratory (LACIRTEM), inferred that the domain of training with WBV

*Corresponding author: Marcelo Renato Guerino, Department of Physical Therapy of Federal University of Pernambuco, Recife-PE, Brazil, Tel: +55 02181997714177; E-mail: marceloguerino@hotmail.com

Citation: de Araujo MGR, Guerino MR(2020) Whole Body Vibration Effectiveness in Diabetes Mellitus, Osteoporosis and Parkinson's Disease. J Gerontol Geriatr Med 6: 059.

Received: June 17, 2020; Accepted: June 24, 2020; Published: July 01, 2020

Copyright: © 2020 de Araujo MGR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

can help in choosing the training program for elderly people with type 2 diabetes [16].

Evidence on the effects of WBV intervention on DM comprises several types of studies and methodologies Robinson et al., [17]. In another study using an exercise program for upper and lower limbs for 12 weeks, a reduction in the time taken to perform the 6-Minutes' Walk Test (6MWT) was observed, but there was no improvement in the Timed Up and Go Test (TUG) [18]. In another work [19], after 8 weeks using WBV, it had a significant effect in reducing patients' fat mass, however no significant effects were found on blood pressure, mobility, and quality of life of diabetics.

In a study of diabetic foot with 54 patients in 12 weeks, aimed to determine whether a WBV program would favour parameters (glycated-HbA1c hemoglobin, sensitivity, transcutaneous-TcPO₂ oximetry) related to associated complications of diabetic foot syndrome [20]. However, there were no significant changes in HbA1c or sensitivity, but TcPO₂ registered a slight increase. In a review studying the significant effects of WBV in patients with diabetic peripheral neuropathy, it was found that WBV provided a slight improvement in the glycemic profile, plantar tactile sensitivity, pain and balance, but the quality of the evidence is low [21]. Recent work from our research laboratory checked the immediate effect of Whole Body Vibration (WBV) on quadriceps muscle strength, functional mobility and balance in elderly patients with Osteopenia and/or Osteoporosis. The sample was composed of 34 elderly and concluded that a single WBV training session was able to increase quadriceps muscle strength and the elderly balance [22].

Important studies support the effectiveness and viability of using WBV to improve muscle performance and balance in individuals with osteoporosis. One hundred and fifty-one postmenopausal women over 65 years of age performed training sessions and after 6 months, the falls were significantly reduced by the training protocol including WBV exposure [23]. In another review article have found that there was a significant increase in articles on this topic and that the majority of studies have shown an improvement of the Bone Mineral Density (BMD) or other parameters regarding muscle, bone and functional outcomes, like reduction of falls [24]. Moreira-Marcioni et al., [25] found in the literal relevance of WBV exercises in decreasing the number of fractures in women with osteoporosis, even though little research exists on the topic, they consider WBV to be an effective strategy to improve balance. After 6 months of the WBV exercises with fifty two osteoporotic women the indices for flexibility, body balance, and walking velocity were significantly improved [26]. Moreira et al., [27] after review concluded that WBV body balance and can be a useful tool as part of the prevention and treatment protocol for osteoporosis. Similar findings were reported by Weber-Rajek et al., [28] and corroborated by Bemben et al., [29].

Slatkovsla et al., [30] in order to verify the beneficial effects of WBV on bones of postmenopausal women, conducted a study of daily training of low magnitude WBV (0.3 g) at 2 frequencies (90 and 30 Hz) versus without WBV and verified that WBV at 0.3g and 90 or 30Hz for 12 months did not alter BMD in women who received calcium and vitamin D supplementation. According a systematic review, the related the effects of the vibrating platform on the bone tissue of osteoporotic women considering 11 eligible articles, however with several methodological differences, they concluded that the evidence

proving the increase in bone mass resulting from the vibration seems to be inconclusive, since authors demonstrate divergent and conflicting results [31]. Another study infers that is premature to recommend WBV as a safe and effective training for preventing osteoporosis due to differences in the type of used vibration (lateral alternating vs. oscillation), frequency, intensity, cumulative dose, body positioning and study methodology [32].

Benedetti et al., [33] reported that training with WBV promotes increased muscle strength, improves balance, and reduces the risk of falling in osteoporosis patients. However, the results are controversial about the improvement in bone mineral density, a fact, in accordance with other studies literature [34-36]. Individuals with Parkinson's Disease (PD) are affected by the second most common neurodegenerative disorder that mainly promotes motor and cognitive disability [37,38], providing an increase in the potential for falls in the elderly, impacting the quality of life and costs for the person and the family. There is weak evidence of a positive short-term effect of WBV on lower limb spasticity, mobility, balance and postural control [39], as its benefits are long-term, where WBV training improves mobility while minimizing neurological disorders since the findings of Charcot [40].

There is no clear evidence of a reducing effect on motor symptoms, balance, gait and mobility [41]. Only a few studies have found significant differences between groups regarding mobility and motor symptoms [37,42]. Aerobic exercise on the treadmill performs a dynamic movement compared to the WBV, which is isometric, where the treadmill training improves blood flow which facilitates the recovery phases (free fatty acid and amino acid feedback), in contrast to the WBV, where sustained isometric contractions produce compartmental and intramuscular pressures that can limit blood flow [37]. In another study, when WBV was compared with conventional therapy, parameters like mobility ("Up & Go" Test, a 10-m walk test, and a stand-walk-sit test) haven't had any significant treatment effect after multiple sessions of WBV. In patients diagnosed with idiopathic Parkinson's disease, using de protocol of 5 weeks, 2-3 times a week with static position, 5 × 60 s/60 s, knees slightly bent, 6 Hz, 3 mm. No significant difference was observed between the WBV group and the placebo group for the TUG test (SMD = -0.37; 95% CI: -1.34, 0.59), gait velocity (SMD = -0.21; 95% CI: -1.17, 0.74) and step length (SMD = 0.14; 95% CI: -0.81, 1.09) [43]. Research results should be analyzed with caution due to the lack of homogeneity in interventions and methodologies in some studies [44]. In our clinical practice, the established vibration frequency was 24 Hz. The 4-mm displacement peak was maintained throughout the program and was determined by the width of the volunteers' position on the platform. The intervention time was 45 s, with eight series being performed and, for each, the recovery time was maintained at 30 s for all intervals, totaling 600 s of intervention [16].

As physiotherapists, we found that the patient with Parkinson's disease, most of the time has severe physical problems such as tremor and cognition, which leads to anxiety and difficulties in social life. This led us to look for therapeutic options in addition to conventional treatment. Making the WBV exercises accessible, with an emphasis on complementing exercise programs, is essential for elderly populations resistant to certain types of treatments with physical activity, regardless of the functional level of the patients. A study investigating non-drug treatment for Parkinson's disease reports that training

with WBV has an insufficient level of evidence even in improving the time-rhythmic correction of walking in patients with Parkinson's disease [45]. Analyzing four systematic reviews we can highlight that WBV improves strength, postural control and mobility in patients with neurological disease such as Parkinson's disease, stroke and cerebral palsy [43,46-48]. They also found that balance and mobility seem to respond better to the association of WBV with therapeutic exercises, particularly in patients with low function, compared to only WBV, so care is needed when interpreting the findings. Although there is some evidence for a general effect of WBV on selected balance and mobility measures, the impact remains inconclusive. The great difficulty is in the different models of training programs, since some studies use single sessions while other studies use multiple sessions and longer periods of time, therefore, the differences between short and long term interventions need to be better examined.

When this type of therapy is compared with aerobic training, according to our clinical practice, WBV does not seem to require much recovery time as well as reduction in the feeling of fatigue after training. In fact, the feeling of fatigue of some types of stimulus to patients with Parkinson's seems to clearly minimize its effects due to fatigue, which is an important fact to be interpreted by the physiotherapist in his clinical practice, because vibratory stimulation has provided an improvement in trunk oscillation, gait in space-time parameters and cadence, which facilitates the patient's postural control [49-51], as well as the characteristics of responses in the initial and final stages of the disease [52,53].

Thus, in our view, the use of whole body vibration is an important tool for elderly people with Parkinson's disease, from medium to high functional levels. We need to continue researching our ability to improve neurological control, developing training programs, in an accessible and easy to use manner, whenever necessary.

Acknowledgement

The authors would like to thank Lucca Renato Guerino for reviewing this article.

References

1. Rittweger J (2010) Vibration as an Exercise Modality: How It May Work, and What Its Potential Might Be. *Eur J Appl Physiol* 108: 877-904.
2. Hagbarth KE, Eklund G (1969) The Muscle Vibrator--A Useful Tool in Neurological Therapeutic Work. *Scand J Rehabil Med* 1: 26-34.
3. Sanders CE (1936) Cardiovascular and peripheral vascular diseases: Treatment by a motorized oscillating bed. *JAMA* 106: 916-918.
4. Whedon GD, Deitrick JE, Shorr E (1949) Modification of the Effects of Immobilization Upon Metabolic and Physiologic Functions of Normal Men by the Use of an Oscillating Bed. *Am J Med* 6: 684-711.
5. Nazarov V, Spivak G (1985) Development of athlete's strength abilities by means of biomechanical stimulation method. *Theory PracPhys Culture* 12: 445-450.
6. Lora MH, Granados SR, Corrales BS, Páez LC (2009) Efecto de unaseñón con vibraciones mecánicas sobre la capacidad de salto. *Rev Int Med Cienc Act FisDeporte* 9: 366-378.
7. Pessoa MF, Brandão DC, de Sá RB, de Souza HCM, Fuzari HKB, et al. (2017) Effects of Whole Body Vibration on Muscle Strength and Quality of Life in Health Elderly: A Meta-Analysis *Fisioter Mov* 30: 171-182.
8. Lee WS, Cheung W-H, Qin L, Tang N, Leung K-S (2006) Age-associated Decrease of Type IIA/B Human Skeletal Muscle Fibers. *ClinOrthopRelat Res* 450: 231-237.
9. Thomas E, Battaglia G, Patti A, Brusa J, Leonardi V, et al. (2019) Physical Activity Programs for Balance and Fall Prevention in Elderly: A Systematic Review. *Medicine (Baltimore)* 98: 16218.
10. Okubo Y, Schoene D, Lord SR (2017) Step Training Improves Reaction Time, Gait and Balance and Reduces Falls in Older People: A Systematic Review and Meta-Analysis. *Br J Sports Med* 51: 586-593.
11. Lee K, Lee S, Song C (2013) Whole-body Vibration Training Improves Balance, Muscle Strength and Glycosylated Hemoglobin in Elderly Patients With Diabetic Neuropathy. *Tohoku J Exp Med* 231: 305-314.
12. Pozo-Cruz JD, Alfonso-Rosa RM, Ugia JL, Mcveigh JG, Pozo-Cruz BD, et al. (2013) A Primary Care-Based Randomized Controlled Trial of 12-week Whole-Body Vibration for Balance Improvement in Type 2 Diabetes Mellitus. *Arch Phys Med Rehabil* 94: 2112-2118.
13. Yoosefinejad AK, Shadmehr A, Olyaei G, Talebian S, Bagheri H (2014) The Effectiveness of a Single Session of Whole-Body Vibration in Improving the Balance and the Strength in Type 2 Diabetic Patients With Mild to Moderate Degree of Peripheral Neuropathy: A Pilot Study. *J Bodyw Mov Ther* 18: 82-86.
14. Yoosefinejad AK, Shadmehr A, Olyaei G, Talebian S, Bagheri H, et al. (2015) Short-term Effects of the Whole-Body Vibration on the Balance and Muscle Strength of Type 2 Diabetic Patients With Peripheral Neuropathy: A Quasi-Randomized-Controlled Trial Study. *J Diabetes Metab Disord* 14: 45.
15. Gomes-Neto M, de Sá-Caputo DC, Paineiras-Domingos LL, Brandão AA, Neves MF, et al. (2019) Effects of Whole-Body Vibration in Older Adult Patients With Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis. *Can J Diabetes* 43: 524-529.
16. Alves KFP, Ferreira APL, Parente LCO, Rodrigues FTM, Marques TV, et al. (2020) Immediate Effect of Whole-Body Vibration on Skin Temperature and Lower-Limb Blood Flow in Older Adults with Type 2 Diabetes: Pilot Study. *Appl Sci* 10: 690.
17. Robinson CC, Barreto RPG, Sbruzzi G, Plentz RDM (2016) The Effects of Whole Body Vibration in Patients With Type 2 Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Braz J Phys Ther* 20: 4-14.
18. del Pozo-Cruz B, Alfonso-Rosa RM, Pozo-Cruz JD, Sañudo B, Rogers ME (2014) Effects of a 12-wk Whole-Body Vibration Based Intervention to Improve Type 2 Diabetes. *Maturitas* 77: 52-58.
19. Domínguez-Muñoz FJ, Villafaina S, García-Gordillo MA, Hernández-Mocholi MA, Collado-Mateo D, et al. (2020) Effects of 8-Week Whole-Body Vibration Training on the HbA1c, Quality of Life, Physical Fitness, Body Composition and Foot Health Status in People With T2DM: A Double-Blinded Randomized Controlled Trial. *Int J Environ Res Public Health* 17: 1317.
20. Reyes GR, Carrera LN, Monteiro AA, Vivanco AS, Uriostegui IQ, et al. (2017) Effect of Mechanical Vibration on Transcutaneous Oxygen Levels in the Feet of Type 2 Diabetes Mellitus Patients. *Med Clin (Barc)* 148: 16-19.
21. Robinson CC, Barreto RPG, Plentz RDM (2018) Effects of Whole Body Vibration in Individuals With Diabetic Peripheral Neuropathy: A Systematic Review. *J Musculoskelet Neuronal Interact* 18: 382-388.
22. Ramos LAX, Rodrigues FTM, Shirahige L, Barros MFA, de Carvalho AGC, et al. (2019) Ferreira APL, Guerra RO, Araújo MGR. A Single Whole Body Vibration Session Influences Quadriceps Muscle Strength, Functional Mobility and Balance of Elderly With Osteopenia and/or Osteoporosis? Pragmatic Clinical Trial. *J Diabetes Metab Disord* 18: 73-80.

23. von Stengel S, Kemmler W, Mayer S, Engelke K, Kalender WA (2009) [Effect of Whole Body Vibration Exercise on Osteoporotic Risk Factors]. *Dtsch Med Wochenschr* 134: 1511-1516.
24. Dionello CF, Sá-Caputo D, Pereira HVFS, Sousa-Gonçalves CR, Maiworm AI, et al. (2016) Effects of Whole Body Vibration Exercises on Bone Mineral Density of Women With Postmenopausal Osteoporosis Without Medications: Novel Findings and Literature Review. *J Musculoskelet Neuronal Interact* 16: 193-203.
25. Moreira-Marconi E, Dionello CF, Morel DS, Sá-Caputo DC, Souza-Gonçalves CR, et al. (2016) Could whole body vibration exercises influence the risk factors for fractures in women with osteoporosis? *Osteoporosis and Sarcopenia* 2: 214-220.
26. Iwamoto J, Sato Y, Takeda T, Matsumoto H (2012) Whole Body Vibration Exercise Improves Body Balance and Walking Velocity in Postmenopausal Osteoporotic Women Treated With Alendronate: Galileo and Alendronate Intervention Trial (GAIT). *J Musculoskelet Neuronal Interact* 12: 136-143.
27. Moreira LDF, de Oliveira ML, Lirani-Galvão AP, Marin-Mio RV, de Santos RN, et al. (2014) Physical Exercise and Osteoporosis: Effects of Different Types of Exercises on Bone and Physical Function of Postmenopausal Women. *Arq Bras Endocrinol Metabol* 58: 514-522.
28. Weber-Rajek M, Mieszkowski J, Nnieszpodziński B, Ciechanowska K (2015) Whole-body vibration exercise in postmenopausal osteoporosis. *PrzMenopauzalny* 14: 41-47.
29. Bembem D, Stark C, Taiar R, Bernardo-Filho M (2018) Relevance of Whole-Body Vibration Exercises on Muscle Strength/Power and Bone of Elderly Individuals. *Dose Response* 16: 1559325818813066.
30. Slatkova L, Alibhai SMH, Beyene J, Hu H, Demaras A, et al. (2011) Effect of 12 Months of Whole-Body Vibration Therapy on Bone Density and Structure in Postmenopausal Women: A Randomized Trial. *Ann Intern Med* 155: 668-679.
31. Jung MF, Preis C, Neto LB (2015) Effects of vibration in bone tissue of postmenopausal women: a systematic review. *Geriatr Gerontol Aging* 9: 112-121.
32. Daly RM, Via JD, Duckham RL, Fraser SF, Helge EW (2019) Exercise for the Prevention of Osteoporosis in Postmenopausal Women: An Evidence-Based Guide to the Optimal Prescription. *Braz J Phys Ther* 23:170-180.
33. Benedetti MG, Furlini G, Zati A, Mauro GL (2018) The Effectiveness of Physical Exercise on Bone Density in Osteoporotic Patients. *Biomed Res Int* 4840531: 10.
34. Lau RWK, Liao L-R, Yu F, Teo T, Chung RCK, et al. (2011) The Effects of Whole Body Vibration Therapy on Bone Mineral Density and Leg Muscle Strength in Older Adults: A Systematic Review and Meta-Analysis. *Clin Rehabil* 25: 975-988.
35. Cheung AM, Giangregorio L (2012) Mechanical Stimuli and Bone Health: What Is the Evidence? *Current Curr Opin Rheumatol* 24: 561-566.
36. Jepsen DB, Tomsen K, Hansen S, Jørgensen NR, Masud T, et al. (2017) Effect of Whole-Body Vibration Exercise in Preventing Falls and Fractures: A Systematic Review and Meta-Analysis. *BMJ Open* 7: 018342.
37. Corbianco S, Cavallini G, Baldereschi G, Carboncini MC, Fiamingo FL, et al. (2018) Whole Body Vibration and Treadmill Training in Parkinson's Disease Rehabilitation: Effects on Energy Cost and Recovery Phases. *Neurol Sci* 39: 2159-2168.
38. Henry M, Baudry S (2019) Age-related changes in leg proprioception: implications for postural control. *J Neurophysiol* 122: 525-538.
39. Alashram AR, Padua E, Annino G (2019) Effects of Whole-Body Vibration on Motor Impairments in Patients With Neurological Disorders: A Systematic Review. *Am J Phys Med Rehabil* 98: 1084-1098.
40. Charcot JM (2011) Vibratory therapeutics. The Application of Rapid and Continuous Vibrations to the Treatment of Certain Diseases of the Nervous System. 1892. *J Nerv Ment Dis* 199: 821-827.
41. Dincher A, Schwarz M, Wydra G (2019) Analysis of the Effects of Whole-Body Vibration in Parkinson Disease - Systematic Review and Meta-Analysis. *PM R* 11: 640-653.
42. Rabert SM, Comas RD, Vanmeerhaeghe AF, Medina CS, Figuls MR, et al. (2012) Whole-body Vibration Training for Patients With Neurodegenerative Disease. *Cochrane Database Syst Rev* 15: CD009097.
43. Lau RWK, Teo T, Yu F, Chung RCK, Pang MYC (2011) Effects of Whole-Body Vibration on Sensorimotor Performance in People With Parkinson Disease: A Systematic Review. *Phys Ther* 91: 198-209.
44. Fischer M, Vialleron T, Laffaye G, Fourcade P, Hussein T, et al. (2019) Long-Term Effects of Whole-Body Vibration on Human Gait: A Systematic Review and Meta-Analysis. *Front Neurol* 10: 627.
45. Pokhobov DV, Abramov VG, Pokhobov DD (2016) [Possibilities of non-drug treatment for Parkinson's disease]. *ZhurnalNevrologii i PsikiatriiImeni SS Korsakova* 116: 22-29.
46. Orr R (2015) The Effect of Whole Body Vibration Exposure on Balance and Functional Mobility in Older Adults: A Systematic Review and Meta-Analysis. *Maturitas* 80: 342-358.
47. Yang X, Wang P, Liu C, He C, Reinhardt JD (2015) The Effect of Whole Body Vibration on Balance, Gait Performance and Mobility in People With Stroke: A Systematic Review and Meta-Analysis. *Clin Rehabil* 29: 627-638.
48. Huang M, Liao LR, Pang MY (2017) Effects of Whole Body Vibration on Muscle Spasticity for People With Central Nervous System Disorders: A Systematic Review. *Clin Rehabil* 31: 23-33.
49. Turbanski S, Haas CT, Schmidtbleicher D, Friedrich A, Duisber P (2005) Effects of Random Whole-Body Vibration on Postural Control in Parkinson's Disease. *Res Sports Med* 13: 243-256.
50. Giardini M, Nardone A, Godi M, Guglielmetti S, Arcolin I, et al. (2018) Instrumental or Physical-Exercise Rehabilitation of Balance Improves Both Balance and Gait in Parkinson's Disease. *Neural Plast* 2018: 5614242.
51. Serio F, Minosa C, De Luca M, Conte P, Albani G, et al. (2019) Focal Vibration Training (Equistasi®) to Improve Posture Stability. A Retrospective Study in Parkinson's Disease. *Sensors (Basel)* 19: 2101.
52. Bzdůšková D, Valkovič P, Hirjaková Z, Kimijanová J, Hlavačka F (2018) Parkinson's Disease Versus Ageing: Different Postural Responses to Soleus Muscle Vibration. *Gait Posture* 65: 169-175.
53. Gorst T, Marsden J, Freeman J (2019) Lower Limb Somatosensory Discrimination Is Impaired in People With Parkinson's Disease: Novel Assessment and Associations With Balance, Gait, and Falls. *Mov Disord Clin Pract* 6: 593-600.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>