

Review Article

Correction of Metabolic Imbalance and Formation of Amino Acid Pool in Cardiovascular Pathology (Review)

Karavay PA¹, Tabola PI¹ and Nefyodov LI^{2*}

¹Department of Angiography, Central Clinical Hospital of Ostrovets, Belarus

² Department of Biochemistry, Yanka Kupala Grodno State University, Belarus

Abstract

A review of data on the mechanisms of formation of the pool of free amino acids in cardiovascular insufficiency and methods for correcting of metabolic imbalance..

Keyword: Amino acids; Blood plasma; Cardiovascular insufficiency

Introduction

The relevance of studies on the role of amino acids in the pathogenesis, prevention and treatment of cardiovascular insufficiency (CVI) is primarily due to the significant practical results of the use of highly purified substances of this class of compounds and their compositions in the treatment of cardiovascular pathology [1-7].

At the same time, the most significant number of applied research is devoted to the search for marker amino acids or their derivatives for the diagnosis of heart and vascular diseases [8-12]. It has been convincingly demonstrated that elimination or correction of changes in intermediate metabolism can be achieved by using individual amino acids and their derivatives, or by combining them as universal natural bioregulators - compounds that directly affect the mechanisms of myocardial cell metabolism in physiological concentrations. To date, there is evidence of the importance of amino acids not only as building blocks for protein synthesis, but also regulators of gene expression at the level of mRNA translation by the mTOR-dependent

*Corresponding author: Nefyodov LI, Department of Biochemistry, Yanka Kupala Grodno State University, Belarus Email Id: l.nefyodov@mail.ru

Citation: Karavay PA, Tabola PI, Nefyodov LI (2022) Correction of Metabolic Imbalance and Formation of Amino Acid Pool in Cardiovascular Pathology (Review), J Hematol Blood Transfus Disord 9: 028.

Received: April 13, 2022; **Accepted:** April 20, 2022; **Published:** April 27, 2022

Copyright: © 2022 Nefyodov LI, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

mechanism, signaling molecules and modifiers of biological responses, as well as precursors of a wide range of bioregulators that play a key role in integration of the main metabolic flows in vascular pathology [13-18].

The human heart uses a large amount of free amino acids as regulators of both myocardial protein metabolism and as substrates for energy metabolism. The dependence of the myocardium on the amino acid fund of the heart increases in heart failure due to the high activity of anabolism in the myocardium and a lack of energy for cardiomyocytes. Anabolic reactions in the heart are dependent on the oxidation of fatty acids, amino acids and glucose. So, normally, the functional activity of the Krebs cycle (TCAC) largely depends on the concentration of amino acids. Free amino acids stimulate the energy of mitochondria under anaerobic conditions, and also contribute to the substrate supply of TCAC [19-22].

Essential to the availability of amino acids is that their uptake by the myocardium depends solely on their arterial levels. The content of branched chain amino acids (BCAA - Ile, Leu, Val) in the myocardium is the most important activator of anabolism in the heart, the level of which does not depend on insulin. A slight increase in the concentration of "arterial" amino acids leads to a significant increase in their absorption by the myocardium. In heart failure, the arterial pool of free aromatic amino acids (AAA- Tyr, Phe), which is the determining factor in the absorption of amino acids by the myocardium. Thus, in patients with CVI, arterial levels of amino acids were reduced and associated with the severity of chronic heart failure and left ventricular dysfunction. Therefore, amino acids are now becoming more and more widely used in practice as cardioprotective substances, promoting metabolism in the heart under anaerobic conditions and hypoxia [23-27].

Comparative assessment and interpretation of changes in the pool of free amino acids at different stages of cardiovascular failure and in the dynamics of its treatment are devoted to only a few works. At the same time, the question of the informativeness of the established changes in the levels of individual amino acids and their significance in comparison with other clinical and biochemical criteria remains practically unclear. The problem of the choice of individual amino acids in the used "aminosols" for the targeted correction of metabolic imbalance in cardiac and vascular pathology remains unsolved. The importance of amino acids in the regulation of the functions of pathological conditions (vasoarterogenesis, arterial thrombosis) of the cardiovascular system has been convincingly established in a number of studies. The decrease in plasma lipid levels under the action of glycine and its derivatives, the positive effect of cystine and aspartate in patients with hyperlipidemia, the hypolipidemic effect of arginine, characterized by a decrease in LDL low density lipids levels and an increase in HDL (high density lipids levels) in plasma, has been repeatedly described [28].

High concentrations of amino acids and their derivatives in platelets have been demonstrated, upon activation of which the agonist binds to a specific receptor to form a complex, thereby transmitting an

energy signal to the cell that activates phosphatase and mobilizes ionized calcium from the dense tubular system. A study of the amino acid sequences of glycoprotein receptor polypeptides that specifically bind hemocoagulation substrates has shown the ability to inhibit platelet aggregation, adhesion, and thrombus formation with synthetic and natural (snake venom) polypeptides containing arginine, glycine, valine and asparagine [29].

The role of free amino acids in the processes of tissue ischemia tolerance and post-ischemic recovery deserves special attention. The protective effect of BCAA in the myocardium is manifested in maintaining contractility, levels of macroergs (ATP, creatinine phosphate), normalization of aortic and coronary blood flow, cardiac output and cardiac input. BCAA activate the production of catabolites of the adenine system during postischemic reperfusion and the utilization of administered amino acids to high-energy substrates of TCAC and promotes the restoration of the functional capabilities of smooth muscle structures [30].

It is well known that the heart is “metabolically omnivorous” because it is able to actively oxidize fatty acids, glucose, ketone bodies, pyruvate, lactate, amino acids, and even its structural proteins (in decreasing order of preference). The energy of these substrates provides not only mechanical contraction, but also the operation of various transmembrane pumps and transporters required to maintain ion homeostasis, electrical activity, metabolism, and myocardial catabolism. Cardiac ischemia and the resulting coronary and heart failure alter both the electrical and metabolic activity of the myocardium. The effects of ischemia on metabolic preference for substrates are poorly understood, although hypoxia during ischemia significantly alters the relative selectivity of the heart in the use of different substrates. Metabolic changes in case of heart rhythm disturbances are the main component of cardiac myopathies. At the same time, the potential contribution of amino acids to the maintenance of cardiac electrical conduction and stability during ischemia is underestimated. Despite clear evidence that amino acids have a cardioprotective effect in ischemia and other cardiac disorders, their role in the metabolism of the ischemic heart has not yet been fully elucidated [30-32].

Studies on the determination of taurine and a number of amino acids prevailing in the myocardium (glutamate, aspartate, glutamine and asparagine) in coronary insufficiency showed their differences in content in the left and right ventricles in coronary insufficiency. Comparison of the levels of these amino acids in aortic stenosis and coronary heart disease in myocardial biopsies showed higher concentrations of taurine in the left ventricle in both situations [33]. With pronounced, progressive cardiosclerosis in the myocardium of rabbits, the content of phenylalanine and tyrosine increased, which was also found in patients with coronary heart disease, and the degree of increase in the level of amino acids varied depending on the clinical forms of coronary atherosclerosis (angina pectoris of various functional classes, myocardial infarction).

The antiatherogenic properties of the derivative of sulfur-containing amino acids taurine (Tau) are due to the fact that the synthesis of taurocholates promotes the absorption of lipids, lipolysis, and the absorption of fatty acids in the intestine. On the other hand, the conjugation of taurine with bile acids affects the elimination of cholesterol from the body and thereby controls cholesterologenesis in atherosclerosis [34]. It is possible that the high level of taurocholates in some mammalian species (rats) makes it difficult to model experimental atherosclerosis, because the exchange rate of bile acids is increased due to the formation of choliltaurine.

Sulfur containing amino acids (SAA) are recognized as one of the most potent lipid metabolism modulators among the amino acids. SAA has been shown to act on HDL (high density lipoprotein) cholesterol levels and reduce LDL (low density lipoprotein) lipoprotein. So, SAA have some beneficial effects in atherosclerosis and related diseases (metabolic syndrome) [35].

The relative availability of SAA, as well as their amount in dietary proteins, determine lipid metabolism. Although it is not completely clear how SAAs affect gene expression and lipid metabolism at the molecular level, it has been shown that SAAs affect metabolism through the activation of transcription and post-translational modification of a number of regulatory proteins [36].

Amino acids arginine and glycine induce a decrease, lysine and BCAA an increase in serum cholesterol levels. It has been hypothesized that the control of cholesterol by insulin and glucagon is regulated by dietary and endogenous amino acids. So the insulin / glucagon ratio has been proposed as an early metabolic index of the effect of dietary proteins on serum cholesterol levels, a risk factor and a general mechanism by which nutritional factors influence the development of atherosclerosis and cardiovascular disease [28-40]. Recently, new evidence has been obtained for the participation of amino acids in the pathogenesis of CVD. For example, glutamate and aspartate are components of the malate / aspartate shunt and their concentrations control the rate of mitochondrial oxidation of glycolytic NADH. Glutamate also controls the rate of urea synthesis, not only as a precursor of ammonia and aspartate, but as a substrate for the synthesis of N-acetylglutamate, an essential activator of carbamoyl phosphate synthase. This mechanism makes it possible to regulate the synthesis of urea at a relatively constant concentration [37].

Certain amino acids (leucine) stimulate protein synthesis and inhibit autophagic protein degradation regardless of changes in cell volume, since they stimulate mTOR and protein kinase, which is one of the components of insulin signal transduction. In the case of low energy supply to cells, stimulation of mTOR with amino acids is inhibited by activation of cAMP-dependent protein kinase. Amino acid-dependent signaling also promotes β -cell insulin production. This stimulates the anabolic action of amino acids [38].

In relation to coronary heart disease, a special role is played by disturbances in the formation of methionine, leading to the accumulation of its precursor, homocysteine, in the blood and urine. Examination and treatment of patients with homocysteinuria revealed early and active development of atherosclerosis in young patients: hyperhomocyst(e)inemia is a significant risk factor for the development of atherosclerosis and coronary heart disease. Clinical studies have revealed a significant effect of methionine on the proliferation of smooth muscle cells, followed by vascular endothelial dysfunction and the development of arterial hypertension with a high risk of thrombosis.

Lysine is involved in the formation of collagen, strengthening the vascular wall, in the formation of carnitine, promotes the utilization of fatty acids for the energy potential of cells and the preservation of the body's immune reactivity. When the walls of the arteries rupture, collagen filaments, connected to each other by lysine, separate and protrude into the lumen of the vessels, like the remains of lysine, and are washed by circulating blood. Lipoprotein A, a specific form of cholesterol present in the bloodstream, has receptors for lysine, binds to it and penetrates into the intima of the vessels, thus triggering the generation of hydrogen peroxide and superoxide radicals [39].

Arginine, a semi-essential amino acid, serves as a precursor of nitric oxide, which affects platelet aggregation and adhesion, decreasing the ability to thrombus formation and decreasing vascular reactivity of atherosclerotic arteries and promotes collagen formation in the vessel walls [24]. In the blood plasma of patients with endothelial disruption in atherosclerosis, the levels of citrate, GABA, glutamate and cysteine were significantly different in comparison with myocardial ischemia in the content of glutamate and phenylalanine. On this basis, a differential diagnosis of aortic injury with ischemic heart disease is considered possible. Arginine is widely used as an antihypertensive drug and prophylactic drug [40].

The development and progression of atherosclerosis, which ultimately leads to cardiovascular disease, is causally related to hypercholesterolemia. Mechanistically, the interaction between lipids and the immune system during the progression of atherosclerotic plaques contributes to the chronic inflammation seen in the artery wall during atherosclerosis. Localized inflammation and increased cell-cell communication can affect the polarization and proliferation of immune cells through changes in amino acid metabolism. In particular, the amino acids L-arginine (Arg), L-homoarginine (hArg), and L-tryptophan (Trp) have been extensively studied in the context of cardiovascular disease and have been shown to act as key regulators of vascular homeostasis, similar to the functions of immune cells. Cyclic effects between endothelial cells, innate and adaptive immune cells occur when the metabolism of Arg, hArg and Trp changes, which has a significant effect on the development of atherosclerosis. Thus, the metabolism and biological functions of Arg, L-homoarginine, and Trp make it possible to reasonably use them for the therapy of atherosclerosis [41].

It has been proven that free amino acids, especially BCAAs, have significant regulatory functions in the processes of protein synthesis. Thus, recent studies have shown that BCAA protect the cardiovascular system from the metabolic consequences of ischemia / reperfusion (I / R). The authors investigated the signaling pathways and functions of mitochondria, as well as the levels of BCAAs that influence the listed processes [30]. Thus, the *in vivo* I / R damage model was tested in controls, mTOR *+/+* and mTOR *+/-*. The mice received BCAA, rapamycin, or BCAA + rapamycin. In addition, isolated cardiomyocytes were subjected to modeling ischemia with a quantitative assessment of their death. The degree of mitochondrial swelling was also assessed. In mice treated with BCAAs, there was a significant reduction in the size of the infarction. In addition, BCAAs prevent mitochondrial swelling, which was controlled by the addition of rapamycin. BCAAs significantly retained cell viability. Thus, BCAAs are protective against I / R myocardial injury, in which mTOR plays an important role [30].

Summarizing, it should be noted that the use of amino acid preparations for pathology of the heart and blood vessels is rational, and the strategy for their use should be based on the elimination of the existing amino acid imbalance in this disease, correction of the pool of free sulfur-containing amino acids, including the use of taurine, arginine and lysine, angio- and cardioprotective properties of which should be considered sufficiently reasonable and promising. Our proposed methodology for the development of formulations of new multicomponent infusion solutions based on amino acids and related compounds, intended for the correction of metabolic imbalance arising in cardiovascular diseases, is based on the application of the results of studying the patterns of formation of the amino acid pool in biological fluids and human tissues with pathology of the cardiovascular system.

References

1. Aquilani R, Rovere MTL, Corbellini D, Pasini E, Verri M, et al. (2017) Plasma Amino Acid Abnormalities in Chronic Heart Failure. Mechanisms, Potential Risks and Targets in Human Myocardium *Metabolism Nutrients* 9: 1251.
2. *Amino Acids and Their Derivatives* (1996) (chemistry, biochemistry, pharmacology, medicine) ed. L.Nefyodov *Proc of Internat.Symp, Grodno*125 P.
3. Nefyodov LI, Karavay PA, Karavay NL (2014) Regulatory action of free amino acids and development on the basis of highly of substances infusion solutions with pathogenetic deterministic composition. *Laboratory diagnosis Eastern Europe* 3: 111-115.
4. Nefyodov LI (2010) The results of biochemical research and development of nitrogen-containing compounds of natural origin: methodology of exploitation of biological properties as universal natural regulators of metabolism and drugs.
5. Nefyodov L I (2020) Amino Acid Imbalance in Atherosclerosis, *Archives of Blood Transfusion and Disorders* 5: 15-23.
6. *Amino Acids* (1990) (Chemistry, Biology, Medicine) / Ed. Lubec C., Rosental J.A. - N.Y.: Escom, 1196 p.
7. Bender DA (1975) *Amino Acid Metabolism*. - N.Y.: J. Willey & Sons, 234 p.
8. Vretlind A, Sudgvan A (1990) *Clinical nutrition*. Stockholm - Moscow: Kabi-Vitrum 355p.
9. Nefyodov LI, Klimovich II, Moroz AR (1991) Statistical analysis of amino acid pool structure in donors blood plasma *Zdravoochranenie Belarusi* 11: 10-13.
10. Meijer AJ (2003) Amino Acids as Regulators and Components of Nonproteinogenic Pathways *The Journal of Nutrition*, 133: 2057-2062.
11. Drake KJ, Sidorov VY, McGuinness OP, Wasserman DH, Wikswo JP (2012) Amino Acids as Metabolic Substrates during Cardiac Ischemia *Exp Biol Med* (Maywood) 237: 1369-78.
12. Wang L, Liu S, Yang W, Yu H, Zhang L, et al. (2017) Plasma Amino Acid Profile in Patients with Aortic Dissection *Scientific Reports* 7, Article number: 40146.
13. Nefyodov LI (2010) The results of biochemical research and development of nitrogen-containing compounds of natural origin: methodology of exploitation of biological properties as universal natural regulators of metabolism and drugs.
14. *Biological activity and transport of drugs* (1999) ed. L. Nefyodov *Proc of Internat.Symp. Grodno* 189 p.
15. VI Ordinary General Assembly Society of Biochemistry of Belarus (2000) ed. L. Nefyodov *Proc of Internat. Symp, Grodno* 225 p.
16. *Amino acids and their derivatives in biology and medicine* (2001) ed. L. Nefyodov *Proc of Internat. Symp, Grodno* 124 p.
17. Nefyodov LI (2001) Target - oriented regulation of metabolic equilibrium by amino acids and strategy of their application as drugs with directional effects p.327. Meijer A (2003) Amino acids as regulators and components of nonproteinogenic pathways. 133: 2057-2062.
18. Bruhat A, Cherasse Y, Chaveroux C, Jousse C, Fafournoux P, et al. (2009) Amino acids as regulators of gene expression in mammals: molecular mechanisms 35: 249-257.
19. Karavay PA (2016) Amino acids in Metabolomics: Perspective for the Use of Regulatory effects of Free Amino Acids in the Creation on their Basis of Infusion Solutions *International Journal of Hematology & Therapy* 2: 1-2.
20. Nefyodov LI (1992) Biological activity of taurine (review) *Vesti AN BSSR* 3: 99-106.

21. Nefyodov LI (1990) Taurine metabolism in mammals (review) *Vesti AN BSSRn* 5: 99-106.
22. Patel RP, Levonen A, Crawford JH, Darley-USmar VM (2000) Mechanisms of the pro- and anti-oxidant actions of nitric oxide in atherosclerosis. *Cardiovasc Res* 47: 465-74.
23. Zalba G, Beaumont J, San Jose G, Fortuno A, Fortuno MA, et al. (2000) Vascular oxidant stress: molecular mechanisms and pathophysiological implications. *J Physiol Biochem* 56: 57-64.
24. Aziz M, Yadav KS (2016) Pathogenesis of Atherosclerosis. *Med Clin Rev* 2:22.
25. Lewis M, Littlejohns B, Lin H, Angelini GD, Saadeh M (2014) Suleiman Cardiac taurine and principal amino acids in right and left ventricles of patients with either aortic valve stenosis or coronary artery disease: the importance of diabetes and gender 3: 523.
26. Birsheim E, Quynh-Uyen TB, Tissier S, Cree MG, Rimsen O, et al. (2009) Amino acid supplementation decreases plasma and liver triglycerides in elderly *Nutrition*. 25: 281–288.
27. Oda H (2006) Functions of Sulfur-Containing Amino Acids in Lipid Metabolism *The Journal of Nutrition*, 136: 1666-1669.
28. Kothandam H, Biradugadda P, Maganti B, Keerthi T, Vegunta B, et al. (2012) "Taurine: A Key Amino Acid in the Drug Discovery" - A Review *Asian Journal of Biomed. & Pharmaceut. Sciences* 2: 21-27.
29. Sanchez A, Hubbard RW (1991) Plasma Amino Acids and the Insulin/Glucagon Ratio as an Explanation for the Dietary Protein Modulation of Atherosclerosis *Medical Hypothesis* 36: 27-32
30. Ma LL, Ji GY, Jiang ZQ (2014) Influence of Dietary Amino Acid Profile on Serum Lipids in Hypercholesterolemic Chinese Adults. *J Nutr Food Sci* 4:258.
31. Holeček M (2018) Branched-chain amino acids in health and disease: metabolism, alterations in blood plasma, and as supplements *Nutrition & Metabolism* 15:33.
32. Mann GE, Yudilevich DL, Sobrevia L (2003) Regulation of Amino Acid and Glucose Transporters in Endothelial and Smooth Muscle Cells *Physiol Rev* 83: 183-252.
33. Selhub J, Troen AM (2016) Sulfur amino acids and atherosclerosis: a role for excess dietary methionine *Ann N Y Acad Sci* 1363:18-25.
34. Nitz K, Lacy M, Atzler D (2019) Amino Acids and Their Metabolism in Atherosclerosis /Arteriosclerosis, Thrombosis, and Vascular Biology 39: 319-330.
35. Satomi S, Morio A, Miyoshi H, Nakamura R, Tsutsumi R, et al. (2020) Branched-chain amino acids-induced cardiac protection against Ischemia/reperfusion injury. *Life Sciences* 245: 117368.
36. Nefyodov LT (1999) (biochemistry, pharmacology, medical application) *Grodno RIPH* 145 p.
37. Nefyodov LI (1993) Formation of the fund of free amino acids and their derivatives in conditions of metabolic imbalance: ... dis. doct. honey. *Sciences: 03.00.04 L.I. Nefyodov; Belar.State Univ Minsk* 264 p.
38. Meijer AJ (2003) Amino Acids as Regulators and Components of Nonproteinogenic Pathways *The Journal of Nutrition*, 133: 2057-2062.
39. Drake KJ, Sidorov VY, McGuinness OP, Wasserman DH, Wikswo JP (2012) Amino Acids as Metabolic Substrates during Cardiac Ischemia *Exp Biol Med* (Maywood).
40. Lewis M, Littlejohns B, Lin H, Angelini GD, M-Saadeh S (2014) Cardiac taurine and principal amino acids in right and left ventricles of patients with either aortic valve stenosis or coronary artery disease: the importance of diabetes and gender 3:523.
41. Oda H (2006) Functions of Sulfur-Containing Amino Acids in Lipid Metabolism *The Journal of Nutrition*, 136: 1666-1669.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.herallopenaccess.us/submit-manuscript>