

## Original Article

# Determination of Red Cell Indices and Sero-prevalence of Transfusion Transmissible Infectious Diseases among Stable Paediatric Sickle Cell Patients

Esan Ayodele Jacob<sup>1\*</sup>, Oyegue Kelvin<sup>2</sup>, Ogunbusuyi Boluwatife<sup>3</sup> and Oyedele Titilayo E<sup>4</sup>

<sup>1</sup>Department of Haematology and Blood Transfusion Science, Federal Teaching Hospital, Ido-Ekiti, Nigeria

<sup>2</sup>Department of Haematology and Blood Transfusion Science, Federal Medical Centre, Owo, Nigeria

<sup>3</sup>Department of Medical Laboratory Science, Afe Babalola University, Nigeria

<sup>4</sup>Department of Medical Laboratory Science, Achievers University, Owo, Nigeria

## Abstract

**Background:** Red blood cell indices are very important tools in screening for anaemia especially in children with sickle cell disease (SCD). Blood transfusion forms an integral part of management of sickle cell disease.

**Aim:** To determine the red cell indices and sero-prevalence of transfusion-transmissible infectious disease among stable paediatric sickle cell patients.

**Methodology:** One hundred clinically stable SCD patients that were transfused in the least six months and not in crisis in the last four weeks were recruited into this study. 4ml of venous blood was collected, 2ml was dispensed into K2EDTA for immediate analysis of red cell indices using haematology analyzer, haemoglobin phenotype was also determined using cellulose acetate at alkaline solution. The remain 2ml was dispensed into plain bottles, blood was allowed to clot and centrifugation at 5000 rpm for 5 minutes. Serum

\*Corresponding author: Esan Ayodele Jacob, Department of Haematology and Blood Transfusion Science, Federal Teaching Hospital, Ido-Ekiti, Nigeria, Tel: +2348035477756; E-mail: ayodelejacob4u@gmail.com

**Citation:** Jacob EA, Kelvin O, Boluwatife O, Titilayo EO (2021) Determination of Red Cell Indices and Sero-prevalence of Transfusion Transmissible Infectious Diseases among Stable Paediatric Sickle Cell Patients. J Hematol Blood Transfus Disord 8: 027.

**Received:** July 05, 2021; **Accepted:** July 12, 2021; **Published:** July 20, 2021

**Copyright:** © 2021 Jacob EA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

was extracted into another plain tubes for serological testing of TTIs using enzyme-linked immunosorbent assay (ELISA).

**Results:** Mean±SD value of RBC, HGB and HCT among the SCD participants were significantly ( $p<0.05$ ) lower compared with control subjects. However, mean±SD value of MCV, MCH, MCHC and RDW among the SCD participants were significantly ( $p<0.05$ ) higher compared with control subjects. Sero-prevalence of TTIs among blood transfused SCD participants in this study showed zero prevalence in all the TTIs tested using standard technique.

**Conclusion:** Red blood cell indices values were lower compared with control subjects, nature has it that SCD patients in the steady state have adapted their body system to this anaemic state and live healthy. Blood transfusion does not significantly increase the sero-prevalence of transfusion transmissible infections in our clinically stable SCD participants.

**Keyword:** Red blood cell indices, Transfusion-transmissible infections, seroprevalence, steady sickle cell disease patients

## Introduction

Sickle cell disease (SCD) is majorly a red blood cell (RBCs) disorder with significant changes observed in certain haematological parameters [1]. This commonest hereditary disorder is characterized by severe haemolytic anaemia results into frequent blood transfusion [2]. Red blood cell indices are very important tools in screening for anaemia in children with and without sickle cell disease, due to chronic haemolysis and increased bone marrow activity in sickle cell disease patients, the red cell indices of children with sickle cell disease is differ from those of children without sickle cell disease [3]. It has been noticed that SCD patients in the steady state have adapted their body system to this anaemic state and live healthy despite low haemoglobin level [4]. Although, SCD patients in crisis (vaso-occlusion) has been reported to have higher haemoglobin level compared with SCD in steady state, this was related to higher blood viscosity in SCD during crisis which further increase VOC [5]. Mean cell volume (MCV), mean cell haemoglobin (MCH) and mean cell haemoglobin concentration (MCHC) are useful parameters in detecting underlying causes of anaemia [9, 6]. There is a significant high red cell distribution width (RDW) reflecting the degree of chronic haemolysis and anaemia in SCD patients [7]. Blood transfusion forms an integral part of management of sickle cell disease. Blood transfusion is also established as a route of transmission of the transfusion-transmissible infectious agents such as hepatitis B virus (HBV), human immunodeficiency virus (HIV), hepatitis C virus (HCV) and syphilis among the greatest threats to blood safety for blood transfusion recipients and pose a serious public health problem [15]. Blood transfusion in sickle cell disease patients is usually performed when haemoglobin is less than 6g/dl, patients with sickle cell disease usually have haemoglobin level as low as 8g/dl in their steady state. Sickle cell disease (SCD) affects about 2% of Nigerians at birth and it is the commonest genetic disorder in the country. [9] reported, 0.69% prevalence of sickle cell anaemia among the children in Ido-Ekiti. Sickle cell disease is an autosomal recessively inherited disorder resulting in a chronic anaemia

and frequently a life-threatening. Sickle red cells assume an abnormal, rigid shape under conditions of reduced oxygen tension and equally adhere to the endothelial walls (in addition with platelets and white cells), leading to vascular occlusion and endothelial dysfunction [10]. Anaemia in sickle cell disease often requires red cell transfusion in its management, complications of blood transfusion in individuals living with SCD are iron overload, red cell alloimmunization and transmission of transfusion-transmissible infections such hepatitis B, hepatitis C, Human Immunodeficiency Virus (HIV) and syphilis. The risk of transfusion-transmissible infections (TTIs) through blood is determined by its prevalence in the donor population, the problem of seroconversion window (where the donor is infectious but seronegative), the availability and affordability of sensitive diagnostic capabilities. The risk of posttransfusion-transmissible infections in the developed countries is presently extremely low [15]. It is therefore, important that blood given to SCD patients is routinely and properly screened for these transfusion-transmissible infections (TTIs) using standard techniques, especially those on regular blood transfusion to reduce the chances of transmitting transfusion-transmissible infections. This is even made more imperative by the observation that a larger proportion of the blood donor pool in Nigeria is commercial (paid) donors. The reason for this may be attributable to a number of factors, ranging from poor awareness on voluntary blood donation to poverty and illiteracy, particularly among the rural populations of the country [11]. Children with SCD constitute a group that requires frequent blood transfusion, this study was therefore, designed to determine the red cell indices and sero-prevalence of transfusion-transmissible infectious disease among stable paediatric sickle cell patients in the paediatric clinic of Federal Teaching Hospital, Ido-Ekiti, Nigeria.

## Materials and Methods

### Study design

This study was carried out in Federal Teaching Hospital, Ido-Ekiti among paediatric sickle cell disease patients in steady state. The study was approved by the hospital's ethics and research committee. Informed consent was obtained from the parents of the participants. Questionnaire was used to collect the information about demographic data, history of blood transfusion/hospital admission and parents' socioeconomic parameters. Inclusion criteria includes confirmed status of sickle cell patients in steady state within the age group between 1 and 15 years. Subjects that were transfused with blood in the last six months and not in crisis in the last four weeks. Exclusion criteria include sickle cell disease patients that does not within the study age group, those that decline from participating in the study and Subjects that were not haemoglobin SS. Control subjects were recruited from apparently healthy children whose haemoglobin phenotype is AA within the age group 1-15 years of both sexes for the purpose of comparing red cell indices parameters with SCD participants. Haemoglobin electrophoresis of subjects were pre-determined using cellulose acetate at alkaline solution (pH 8.6) to confirm the haemoglobin phenotype status of the participants [12]. 4ml of venous blood sample was collected from one hundred SCD participants and control subjects, 2ml was dispensed into K<sub>2</sub>EDTA for immediate analysis of red cell indices using haematology analyzer, haemoglobin phenotype was also determined using cellulose acetate at alkaline solution. The remain 2ml was dispensed into plain bottles, blood was allowed to clot and centrifugation at 5000 rpm for 5 minutes. Serum was thereafter extracted from the clotted sample into another clean plain tubes for serological testing of TTIs using enzyme-linked immunosorbent

assay (ELISA). Data obtained were analyzed using the Statistical Package for Social Sciences, version 22.0 (SPSS 22.0). Statistical significance was set at a p-value less than or equal to 0.05.

## Results

Table 1a shows the socio-demographic characteristics of stable sickle cell patients. Age group 9-12 and 1-4 had highest and lowest prevalence respectively among the study SCD participants. In gender distribution, 55 were male and 45 were female. Christians and Yoruba ethnic in religion and tribe variable respectively had highest prevalence in the study group as showed in table 1a. 55.0% of secondary education, 40.0% of family size of three, 38.0% of firstborn children in the family and 34% were transfused with two Units of blood in the last 6months as showed in Table 1b.

Variable	Participants (%) N = 100	Control (%) N = 100
<b>Age group (in years)</b>		
1 – 4	10 (10.0)	5 (5.0)
5 – 8	32 (32.0)	15 (15.0)
9 – 12	46 (46.0)	30 (30.0)
13 – 15	12 (12.0)	50 (50.0)
Mean age ± SD	10.6 ± 5.1	8.9 ± 3.9
Age range	2 – 15	4 – 15
<b>Gender</b>		
Male	55 (55.0)	60 (60.0)
Female	45 (45.0)	40 (40.0)
<b>Religion</b>		
Christianity	88 (88.0)	75 (75.0)
Islam	12 (12.0)	25 (25.0)
<b>Tribe</b>		
Yoruba	96 (96.0)	80 (80.0)
Ibo	2 (2.0)	15 (15.0)
Hausa	2 (2.0)	5 (5.0)

**Table 1a:** Socio-demographic characteristics of stable sickle cell participants and control.

Variable	Participants N (%) N = 100
<b>Current educational level</b>	
None yet	4 (4.0)
Primary	38 (38.0)
Secondary	55 (55.0)
Tertiary	3 (3.0)
<b>No of children in the family</b>	
1	12 (12.0)
2	26 (26.0)
3	40 (40.0)
4	15 (15.0)
5	5 (5.0)
6	2 (2.0)
<b>Position</b>	
1 <sup>st</sup>	38 (38.0)
2 <sup>nd</sup>	32 (32.0)

3 <sup>rd</sup>	20 (20.0)
4 <sup>th</sup>	6 (6.0)
5 <sup>th</sup>	3 (3.0)
6 <sup>th</sup>	1 (1.0)
<b>Units of blood transfused in the last 6 months</b>	
1	26 (26.0)
2	34 (34.0)
3	20 (20.0)
4	12 (12.0)
5	8 (8.0)

**Table 1b:** Socio-demographic characteristics of the stable sickle cell participants

The mean ± SD value of RBC, HGB and HCT among the SCD participants were significantly ( $p < 0.05$ ) lower compared with control subjects. However, mean ± SD value of MCV, MCH, MCHC and RDW among the SCD participants were significantly ( $p < 0.05$ ) higher compared with control subjects as showed in table 2a.

Variable	Participants N = 100	Control N = 45	Statistical test p-value
	Mean ± SD	Mean ± SD	
RBC	3.31 ± 0.72	5.27 ± 0.83	0.02
HGB	8.58 ± 1.35	13.24 ± 5.86	0.01
HCT	24.87 ± 5.08	39.51 ± 4.03	0.01
MCV	84.18 ± 9.69	81.31 ± 7.18	0.04
MCH	29.42 ± 4.35	25.60 ± 3.10	0.01
MCHC	36.23 ± 1.73	30.23 ± 4.33	0.02
RDW	22.87 ± 2.41	15.34 ± 2.52	0.01

**Table 2a:** Mean± SD of red cell indices among stable sickle cell participants and control.

KEY:

RBC – Red Blood Cell ( $10^6/\mu\text{L}$ )

HGB – Haemoglobin (g/dL)

HCT – Haematocrit (%)

MCV – Mean Cell Volume (fL)

MCH – Mean Cell Haemoglobin (pg)

MCHC – Mean Cell Haemoglobin Concentration (g/dL)

RDW – Red Cell Distribution Width (fL)

Mean± SD of red cell indices in SCD participants and control subjects among the gender was higher in male compared with female although the difference was not significant ( $p > 0.05$ ) except in RDW among the control subjects where the difference is significant as showed in table 2b.

Sero-prevalence of TTIs among blood transfused SCD Participants in this study showed zero prevalence in all the TTIs tested using standard technique as showed in table 3.

## Discussion

One of the characteristic features in the pathophysiology of SCD is chronic hemolytic anemia causing increased tendency for RBC lysis and adhesion which shortened red cell survival, there is also low erythropoietin response to anaemia [5,11]. These facts are associated with low red cell indices in SCD patients during steady state as

Variable	Participants			Control		
	Male N = 55	Female N = 45	p-value	Male N = 60	Female N = 40	p-value
RBC	3.13±0.74	3.03±0.49	0.48	5.10±0.35	4.97±1.44	0.40
HGB	8.47±1.38	8.01±1.25	0.79	13.43±8.43	12.56±4.75	0.84
HCT	25.59±5.37	24.92±4.53	0.46	39.12±5.78	38.81±4.05	0.91
MCV	85.49±6.53	84.98±7.39	0.38	81.31±8.05	80.57±6.32	0.17
MCH	30.22±3.49	29.96±5.40	0.45	25.19±3.26	24.37±3.24	0.08
MCHC	36.32±1.28	35.94±1.39	0.82	30.54±3.76	30.14±2.33	0.22
RDW	22.71±3.30	22.15±3.50	0.63	15.26±5.78	16.34±1.31	0.04

**Table 2b:** Comparison of Mean±SD of red cell indices among the genders SCD participants in steady state and control.

KEY:

RBC – Red Blood Cell ( $10^6/\mu\text{L}$ )

HGB – Haemoglobin (g/dL)

HCT – Haematocrit (%)

MCV – Mean Cell Volume (fL)

MCH – Mean Cell Haemoglobin (pg)

MCHC – Mean Cell Haemoglobin Concentration (g/dL)

RDW – Red Cell Distribution Width (fL)

Parameters	SCD Participants	
	Positive (%)	Negative (%)
HbsAg (n=100)	0.00	100
HCV (n=100)	0.00	100
HIV (n=100)	0.00	100
VDRL (n=100)	0.00	100

**Table 3:** Sero-prevalence of TTIs among blood transfused SCD Participants.

reported in this present study. The values of red cell indices obtained in this study was similar to the results of previous studies conducted in this country, findings in this study confirmed the features of chronic haemolytic disorder in SCD patients [14]. Chronic haemolysis in sickle cell disease patients stimulate haemopoiesis, haemopoietic activity thus providing a more rapid supply of young red blood cells causing mean cell volume to be high despite the anaemia, this shows that haemoglobin value is true measurement of anaemia and not packed cell volume. High MCHC and MCH reported in this study can be supported with the fact that increase in haemoglobin concentration per red cell is associated with cellular dehydration in SCD patients causing red cells of SCD patients to be less deformity in hypoxic state, making it difficult for sickle cells to traverse the microvasculature thus promoting vaso-occlusive crisis. High Red cell Distribution Width (RDW) recorded in this study confirmed the fact that red cell distribution width (RDW) is an index of the variation in the size of the red blood cells which can be used to detect subtle degrees of anisocytosis [15], since SCD is associated with marked anisocytosis morphologically due to rapid erythropoiesis, sickling of red cell and different stages of maturation. The prevalence rates of HBsAg, HCV, HIV and Syphilis in the study were 0 (0.0%), 0 (0.0%), 0 (0.0%), and 0 (0.0%) respectively. It was reported in the previous study that HIV

prevalence in SCD patients varies between 0% and 11.5% which is similar to the findings in this study [1]. Supporting the findings in this study, might due to proper screening of the donor using standard techniques. However, majority of our SCD participants sourced the blood for transfusion at our hospital facility blood bank alone where a higher standard of ensuring safe blood transfusions is ensured, another possible reason for zero prevalence of TTIs in this study may be due to family replacement donor or voluntary donors that normally come to donate especially for the SCD patients because of possible multiple transfusion [15]. However, frequent blood transfusion among the participants in this study was low, this may contribute to zero prevalence of TTIs reported in this study. All SCD participants were vaccinated with hepatitis B vaccine since it is component of the national programme on immunization for children, this might be due to possible cause of zero prevalence in HBV. Supporting the findings in this study, Bolarinware reported that overall risk of HBV infection in SCD patients was not clearly increased by blood transfusion, he further stated that the prevalence of viral markers was not significantly different in SCD participants that had received blood transfusion and those who did not [8].

## Conclusion

Based on the findings in this study, red blood cell indices values were lower compared with control subjects, nature has it that SCD patients in the steady state have adapted their body system to this anaemic state and live healthy. Blood transfusion does not significantly increase the seroprevalence of transfusion transmissible infections in our clinically stable SCD participants; this may be related to the low blood transfusion rate observed in the patients, coupled with the standard donor screening procedures applied in our facility.

## Recommendations

It is recommended that screening of donated blood for TTIs using standard techniques should be continued to prevent risk of transmitting these infections via blood transfusions. Further strengthening of timely birth-dose vaccination will be important for reducing chronic HBV infection. A booster dose should be given to those who did not mount a response to vaccination.

## References

1. Owusu DAE, Visser BJ, Nagel MI, Mens PF, Grobusch MP, Mayer KH et al (2015) The Interaction Between Sickle Cell Disease and HIV Infection: A Systematic Review, *Clinical Infectious Diseases*, 60: 612–626
2. Alagbe EE, Susu AA, Dosunmu AO (2013) Sickle cell disease Management: A Theoretical Review *International Journal of Advanced Research and Review* 16:483-500.
3. Akodu SO, Njokanma OF, Adeolu KO (2015) Erythrocyte indices in Pre-school Nigerian Children with Sickle Cell Anaemia in Steady State. *International Journal of Hematology-Oncology and Stem Cell Research*. 9: 5-9.
4. Akinbami A, Dosunmu A, Adediran A, Oshinaike O, Adebola P, Arogundade et al (2012). "Haematological values in homozygous sickle cell disease in steady state and haemoglobin phenotypes AA controls" *Biomedical Center of Research Notes*. 5: 396.
5. Antwi-Boasiako C, Ekem I, Abdul-Rahman M, Sey F, Doku A, Dzudzor B, Dankwah GB, Out KH, Ahenkorah J, Aryee R et al (2018) Hematological parameters in Ghanaian sickle cell disease patients, *Journal of blood medicine* 9: 203-209.
6. Obeagu EI, Ogbuabor BN, Ikechukwu OA, Chude CN (2014) Haematological parameters among sickle cell anaemia patients in steady state and haemoglobin genotype AA individuals *International Journal of Current Microbiology and Applied Sciences* 3:1000-1005.
7. Abbas M (2014) Haematological parameters in Sudanese children with sickle cell disease *American Journal of Research Communication* 2: 20-32.
8. Esan AJ, Omisakin CT, Okhuakhua O, Omoniyi DP, Owoseni MF et al (2012) "Frequency Distribution of Haemoglobin Variants, ABO and Rhesus Blood Groups among Children in Ido-Osi Local Government, Ekiti State, Nigeria". *Journal of Medical Laboratory Science* 21: 10-18.
9. Iheanacho OE (2015) Haematological parameters of adult and paediatric subjects with sickle cell disease in steady state *International Blood Research and Reviews* 3:171-177.
10. Bian JB, Bates I, Laffan AM, Lewis MS (2012) Cellulose Acetate Electrophoresis at Alkaline pH Dacie and Lewis practical haematology 11<sup>th</sup> edition, Churchill Livingstone 310-312.
11. Nagose V, Rathod SS (2018) Hematological profile of sickle cell anemia subjects in central India: a cross-sectional analysis. *Ann Pathol Lab Med* 5:87–A91.
12. Uche CL, Akinola NO, Durosinmi MA (2017) Haematological and Clinical Assessment of Steady State Homozygous Hb Sand Homozygous Hb A *Journal of Dental and Medical Sciences* 16:117-122.
13. Esan AJ (2016) Prevalence and Evaluation of Iron Deficiency Anemia in Anemic Hospitalized Patients in Osogbo, Nigeria *J Hematol Blood Transfus Disord* 3: 007.
14. Ubesie AC, Emodi IJ, Ikefuna AN, Ilechukwu GC, Ilechukwu GC et al (2012) Prevalence of Human Immunodeficiency Virus Transmission among Transfused Children with Sickle Cell Anemia in Enugu Nigeria, *Ann Med Health Sci Res* 2012 Jul-Dec 2:109–113.
15. Bolarinwa RA, Aneke JC, Olowookere SA, Salawu L (2015) Seroprevalence of transfusion transmissible viral markers in sickle cell disease patients and healthy controls in Ile-Ife, South-Western Nigeria: A case-control study *J Appl Hematol* 6:162-7.
16. Imoro M, Kabiru S, Shehu A, Umar A (2011) Haematological values in Nigeria Children with steady state homozygous sickle cell disease *International Journal of Academic Research* 3: 2.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>