

## Research Article

# Prevalence of Sleep Bruxism in Brazilian children and association with type of delivery, breastfeeding period and oral habits with follow-up in 3 decades

Renato Barcellos Rédua<sup>1\*</sup>, Paulo César Barbosa Rédua<sup>2</sup> and Adriana de Oliveira Lira<sup>3</sup>

<sup>1</sup>Pediatric Dentistry Department, Multivix School of Dentistry, Vitória, Espírito Santo, Brazil

<sup>2</sup>Pediatric Dentistry Department, São Leopoldo Mandic School of Dentistry, Vila Velha, Espírito Santo, Brazil

<sup>3</sup>Pediatric Dentistry Department, Cruzeiro do Sul University, São Paulo, Brazil

### Abstract

**Purpose:** to verify the prevalence of Sleep Bruxism (SB) in children in the last 3 decades and to verify the possible association of this condition with type of delivery, breastfeeding period, oral habits and gender.

**Materials and methods:** A 30-year cohort composed of the dental records of 600 children who were 12-23 months old at their first dental appointment and who were followed up to 12 years of age (200 followed since 1981, Group 1; 200 followed since 1991, Group 2; and 200 followed since 2001, Group 3) was analytically and quantitatively evaluated. Random sample calculation was performed with 95% confidence, a maximum error of 2.95% and a ratio of 50%.

**Results:** The prevalence of SB in children was 13%, 16.5% and 20.5% in the 80s, 90s and 2000s respectively. There was no association between SB and type of delivery, breastfeeding period longer than 1 year, oral habits and gender.

**Conclusion:** The prevalence of sleep bruxism in children has increased in the last three decades. Type of delivery, breastfeeding

\*Corresponding author: Renato Barcellos Rédua, Pediatric Dentistry Department, Multivix School of Dentistry, Vitória, Espírito Santo, Brazil, Tel: +55 27999432952; E-mail: renatoredua@hotmail.com

**Citation:** Rédua RB, Rédua PCB, Lira AO (2021) Prevalence of Sleep Bruxism in Brazilian children and association with type of delivery, breastfeeding period and oral habits with follow-up in 3 decades. J Neonatol Clin Pediatr 8: 086.

**Received:** August 19, 2021; **Accepted:** August 31, 2021; **Published:** September 07, 2021

**Copyright:** © 2021 Rédua RB, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

period, oral habits and gender has no association with the occurrence of SB.

**Keywords:** Breastfeeding; Childhood sleep bruxism; Delivery; Finger sucking; Obstetric; Sleep bruxism

### Introduction

Bruxism has two different circadian manifestations. Awake Bruxism (AB) is a daytime parafunctional habit, while Sleep Bruxism (SB) is considered a sleep behavior [1], that can affect oral health, depending on its severity [2]. The SB is a masticatory muscle activity during sleep that is characterized as rhythmic (phasic) or non-rhythmic (tonic), and it is not considered a movement disorder or a sleep disorder in otherwise healthy individuals [3], but a sign of a health condition in some individuals (e.g. obstructive sleep apnea, sleep disorders, gastro-esophageal reflux) [2,4-6].

Regarding the consequences of SB, they include tooth surface wear, recurrent restorative fractures and even pulp exposure [1,7], masticatory muscles pain or fatigue and temporomandibular disorders [8], parent apprehension about teeth creak in by child during sleep, implications in oral health-related quality of life related to causal factors [9] like obstructive sleep apnea, sleep disorders, snoring and gastro-esophageal reflux [2,4-6,9].

The prevalence of SB in children is very variable, ranging from 3.5 to 46% [10,11]. This variance may be attributed to fact that the diagnosis of SB in children is still challenging, once it is predominantly accessed through parental report [12]. Although polysomnography is the current reference standard for diagnosing SB, it has some disadvantages, such as high cost and technical difficulties when used in children [13]. Other validated methods such as physical examination and/or questionnaires are often used [14]. However, it is important to consider that signs of tooth wear can be related to past episodes of bruxism, as if they were scar not necessarily indicating its occurrence in the present time.

The SB has no gender prediction [10], but psychosocial factors, in particular anxiety and stress, were often associated with in children [15]. Common events in a child's life, such as the birth of a sibling, introduction to school life and school change, for example, can lead to anxiety or fear and, as a result, trigger episodes of SB [16]. Children who have a night sleep period smaller than 8 hours are more likely to develop SB, as well as chronic rhinitis, asthma, amygdala hypertrophy, adenoid hypertrophy, obstructive sleep apnea, sleep disorders and snoring [2,4-6,9]. Gastroesophageal reflux is also mentioned as an etiological factor, although the physiological mechanism of this stimulus is not yet fully understood [6,17].

Although Sari and Sonmez [18] and Behr [19] demonstrate a relationship between bruxism and occlusal factors, there is greater and strong scientific evidence that this association cannot be proven [7,10,20-23]. Regarding the association between oral habits and SB, the literature presents few and divergent studies [9,10,24], but always recognizing the relationship of the child's anxiety with the SB.

We did not find until the search period (December 2019) studies verifying the association of SB with type of delivery and breastfeeding period.

This Practice-based Research aimed to verify if there was an association between SB in children and the type of delivery, breastfeeding period, oral habits and gender. Besides checking if the prevalence of SB in children has increased in the last 3 decades.

## Materials and methods

Before data collection, this study received approval from the local Human Research Ethics Committee. A convenience sample of 600 patients was defined to compose 3 groups of 200 children: Group 1 children who had their first dentist visit in the early 1980s, Group 2 in the early 1990s, and Group 3 in the early 2000s. Inclusion criteria were: Age between 12 and 23 months of life during the first dentist visit, first dentist visit performed in the early years of the decade and child followed until 12 years old. A total of 1,323 dental records of children were analyzed. 771 records from children whose first dental visit occurred between 12 and 23 months of age were included (271 in Group 1, 258 in Group 2 and 242 in Group 3) and 552 records of children in which the first visit occurred at different ages were excluded. Exclusion criteria were: Children with neuropsychomotor alterations reported at the first anamnesis or diagnosed later were excluded, as well as incomplete or illegible records, children who abandoned follow-up before 12 years old, and children attending another pediatric dentist during the follow-up period (71 in Group 1, 58 in Group 2 and 42 in Group 3). All children were followed for 11 years by the same professional, who was a specialist and professor of pediatric dentistry, in a private practice in the city of Vitória, ES, Brazil.

The sample was verified by ratio error and the expected maximum error was 2.96%. The error was verified by calculating the sample size for proportion estimators, considering the 95% confidence level and 50% ratio, which maximizes the variability and results in the highest possible value for the sample. The correction factor for finite population was used.

Statistical analysis was done by descriptive analysis of the research data. The logistic regression test was applied to verify the association between SB and the factors investigated. The significance level was 5% with a 95% confidence interval. Statistical analyses were performed on the SPSS Statistics version IMB 21 (SPSS Inc, Chicago, USA).

In the first visit the data collected includes the type of birth of the child, breastfeeding period and presence of oral habits. Information was given about the rational use of sugar, preferably at predetermined feeding times, oral hygiene instructions and information to parents about SB expression and its consequences. During this first visit, a liability term was provided for parents ensuring that, if the child returned at every 6 months for preventive visits, in the case of carious lesions, the dental professional would perform the treatment without financial costs for the family [25].

After the first dental visit, follow-up visits were scheduled by sending letters for periodic clinical evaluations at every six months. When the child did not show up, two additional letters were sent, at seven months after the last visit and at 11 months after the last visit [25]. At each recall visit parents were asked about oral habits and about the occurrence of SB in the child. Clinically, the presence of dental wear was evaluated; however, the diagnosis of SB was made only when confirmed by parents' report about children teeth grinding at night.

## Results

The mean age of children at the first office visit was 20.4, 19 and 17.1 months in Groups 1, 2 and 3 respectively. There were 307 boys and 293 girls with 106 and 94, 103 and 97 and 98 and 102 boys and girls in Groups 1, 2 and 3 respectively. Intervals between sessions of up to 8 months or less were observed in 41 children (20.5%) in Group 1, 78 (39%) in Group 2 and 92 (46%) in Group 3. A maximum interval between sessions of approximately 9 to 12 months was observed in 123 children (61.5%) in Group 1, 93 (46.5%) in Group 2, and 74 (37%) in Group 3. A period exceeding 12 months between sessions was observed in 36 children (18%) in Group 1, 29 (14.5%) in Group 2 and 34 (17%) in Group 3.

Of the 600 children included in the study, 100 (16.6%) were diagnosed with SB, (46 boys and 54 girls). Of the 200 children in each group, 26 children (13%) were diagnosed with SB in Group 1, 33 (16.5%) in Group 2 and 41 (20.5%) in Group 3. Of the children observed with bruxism, 30 of them were diagnosed in the first to fourth year of life, 55 from the fifth to the eighth year and 15 from the ninth to the twelfth year (Table 1).

	Children with SB	Male	Female	Diagnosed from the first to fourth year age	Diagnosed from the fifth to eighth year age	Diagnosed from the ninth to twelfth year age
Group 1	26	13	13	4	12	10
Group 2	33	17	16	9	22	2
Group 3	41	16	25	17	21	3
Total	100	46	54	30	55	15

**Table 1:** Prevalence and age of SB diagnosis in 600 children.

About the type of delivery, in Group 1 91 (45.5%) children were born by caesarean section and 108 (54%) through normal delivery, and one did not obtain information (adoption), in Group 2 113 (56.5%) through cesarean section and 87 (48.5%) through normal delivery and in Group 3 171 (85.5%) through cesarean section and 29 (14.5%) through normal delivery.

The average period of breastfeeding was 6.76 months in Group 1, 7.09 months in Group 2 and 7.7 months in Group 3. Of the total sample, 18 children were not breastfed, 100 received breastfeeding from 1 to 3 months, 197 from 4 to 6 months, 219 from 7 to 12 months, 61 from 13 to 24 months and 5 children were breastfed for over 25 months.

Of the 600 children in the study, 99 had persistent oral habits (pacifier and / or finger) after 4 years of age, 29 children in Group 1, 35 in Group 2 and 35 in Group 3.

The logistic regression test with a 5% significance level showed no statistically significant association between SB occurrence and breastfeeding period longer than 1 year (Table 2). The logistic regression test demonstrated that the type of delivery was not a risk factor for the occurrence of SB (Table 3). The logistic regression test showed that the presence of oral habits has no association with the occurrence of SB (Table 4). There was no gender predisposition with the occurrence of SB (Table 5).

	p value	OR	95.0% confidence interval for OR	
			Inferior limit	Upper limit
Breastfeeding period longer than 1 year	0.312	1.022	0.980	1.066
OR. OddsRatio.				
Pseudo-R <sup>2</sup> = 0.004.				

**Table 2:** Logistic regression model to verify the association between SB occurrence and breastfeeding period.

	p value	OR	95.0% confidence interval for OR	
			Inferior limit	Upper limit
Type of delivery (Caesarean)	0.729	0.927	0.601	1.428
OR. OddsRatio.				
Pseudo-R <sup>2</sup> = 0.004.				

**Table 3:** Logistic regression model to verify the association between SB occurrence and type of delivery.

	p value	OR	95.0% confidence interval for OR	
			Inferior limit	Upper limit
Presence of oral habits	0.194	1.368	0.852	2.195
OR. Odds Ratio.				
Pseudo-R <sup>2</sup> = 0.005.				

**Table 4:** Logistic regression model to verify the association between SB occurrence and oral habits.

	p value	OR	95.0% confidence interval for OR	
			Inferior limit	Upper limit
Gender (Male)	0.239	0.776	0.509	1.184
OR. OddsRatio.				
Pseudo-R <sup>2</sup> = 0.005.				

**Table 5:** Logistic regression model to verify the association between SB occurrence and gender.

## Discussion

A search of the available literature indicates that this appears to be the first 30-year historical cohort study investigating the prevalence of SB in children and investigating the association between SB and type of delivery and breastfeeding period.

In this study, we consider the diagnosis of SB only when confirmed by the parents report. The use of electromyographic for this diagnosis was not available in the 1980s; in addition to the high costs in the following decades limit its use [13]. Even considering the follow-up of the child from the first year of life to 12 years, the diagnosis was not made exclusively by signs of tooth wear, as these may be related to other causes.

We verify in this study the increase of SB in children in the last 3 decades, this factor may be related to the increase in children's life rhythm, additional out-of-school activities and even greater demand for school performance, as suggested by Serra et al [26] and Castorflorio et al [5] who also found an association of high levels of responsibility by children. Additional factor considered is the increase in pollution in large centers, with studies relating a greater number of

respiratory problems in the population, such as amygdala hypertrophy and adenoid hypertrophy, especially in children.

The results not show prediction by gender, corroborating other published studies [10,27]. Although we have not found other studies relating the occurrence of SB with type of delivery and breastfeeding period, our results show the absence of this relationship. Results that were expected because the literature demonstrates that SB is related to central changes: Such as psychological factors related to fear and anxiety of the child [9,15] and as changes in oxygen saturation during the sleep caused by obstructive sleep apnea [28,29].

Our results do not show association between oral habits and SB corroborating with Soares-Silva [9,24] who demonstrates no relationship about SB and finger or pacifier sucking, while Gonçalves et al [10] related that onicofagy was related to SB in children but it is much more related to the child's anxiety component than to the local stimulus, as discussed in the previous paragraph.

## Conclusion

The prevalence of sleep bruxism in children was 13%, 16.5% and 20.5% in the 80s, 90s and 2000s respectively. There was no association between SB and type of delivery, breastfeeding period, oral habits and gender.

## References

- McKee JA, Banik SS, Boyer MJ, Hamad NM, Lawson JH, et al. (2003) Human arteries engineered *in vitro*. EMBO Rep 4: 633–638.
- Chang S, Song S, Lee J, Yoon J, Park J, et al. (2014) Phenotypic Modulation of Primary Vascular Smooth Muscle Cells by Short-Term Culture on Micropatterned Substrate. PLoS ONE 9: e88089.
- Buehrer BM, Cheatham B (2013) Isolation and Characterization of Human Adipose-Derived Stem Cells for Use in Tissue Engineering. Methods Mol Biol 1001: 1-11.
- Wang A, Tang Z, Li X, Jiang Y, Tsou DA, et al. (2011) Derivation of smooth muscle cells with neural crest origin from human induced pluripotent stem cells. Cells Tissues Organs 195: 5-14.
- Wang C, Yin S, Cen L, Liu Q, Liu W, et al. (2010) Differentiation of adipose-derived stem cells into contractile smooth muscle cells induced by transforming growth factor-beta1 and bone morphogenetic protein-4. Tissue Eng. Part A 16: 1201-1213.
- Kim MR, Jeon ES, Kim YM, Lee JS, Kim JH (2009) Thromboxane a(2) induces differentiation of human mesenchymal stem cells to smooth muscle-like cells. Stem Cells 27: 191-199.
- Elçin AE, Parmaksiz M, Dogan A, Seker S, Durkut S, et al. (2017) Differential gene expression profiling of human adipose stem cells differentiating into smooth muscle-like cells by TGFβ1/BMP4. Exp Cell Res Mar 352: 207-217.
- Bourin P, Bunnell BA, Casteilla L, Dominici M, Katz AJ, et al. (2013) Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT). Cytotherapy 15: 641-648.
- Zuk PA, ZhuM, Ashjian P, De Ugarte DA, Huang JI, et al. (2002) Human adipose tissue is a source of multipotent stem cells. Mol Biol Cell 13: 4279-4295.
- Houtgraaf JH, den Dekker WK, van Dalen BM, Springeling T, de Jong R, et al. (2012) First experience in humans using adipose tissue-derived regenerative cells in the treatment of patients with ST-segment elevation myocardial infarction. J Am Coll Cardiol 59: 539-540.

11. Henry TD, Pepine CJ, Lambert CR, Traverse JH, Schatz R, et al. (2017) The Athena trials: Autologous adipose-derived regenerative cells for refractory chronic myocardial ischemia with left ventricular dysfunction. *Catheter Cardiovasc Interv* 89: 169-177.
12. Travnickova M, Pajorova J, Zarubova J, Krocilova N, Molitor M, et al. (2020) The Influence of Negative Pressure and of the Harvesting Site on the Characteristics of Human Adipose Tissue-Derived Stromal Cells from Lipoaspirates. *Stem Cells Int* 2020: 1016231.
13. Gimble J, Guilak F (2003) Adipose-derived adult stem cells: isolation, characterization, and differentiation potential. *Cytherapy* 5: 362-369.
14. Fraser JK, Wulur I, Alfonso Z, Hedrick MH (2006) Fat tissue: an underappreciated source of stem cells for biotechnology. *Trends Biotechnol* 24: 150-154.
15. Silva FJ, Holt DJ, Vargas V, Yockman J, Boudina S, et al. (2014) Metabolically Active Human Brown Adipose Tissue Derived Stem Cells. *Stem Cells* 32: 572-581.
16. Si Z, Wang X, Sun C, Kang Y, Xu J, et al. (2019) Adipose-derived stem cells: Sources, potency, and implications for regenerative therapies. *Biomed Pharmacother* 114: 108765.
17. Tsekouras A, Mantas D, Tsilimigras ID, Moris D, Kontos M, et al. (2017) Comparison of the Viability and Yield of Adipose-Derived Stem Cells (ASCs) from Different Donor Areas. *In Vivo* 31: 1229-1234.
18. Owens GK, Kumar MS, Wamhoff BR (2004) Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol Rev* 84: 767-801.
19. Fisher SA (2010) Vascular smooth muscle phenotypic diversity and function. *Physiol Genomics* 42: 169-187.
20. Helms F, Lau S, Klingenberg M, Aper T, Haverich A, et al. (2020) Complete Myogenic Differentiation of Adipogenic Stem Cells Requires Both Biochemical and Mechanical Stimulation. *Ann Biomed Eng* 48: 913-926.
21. Webb RC (2003) Smooth muscle contraction and relaxation. *Adv Physiol Educ* 27(1-4): 201-206.
22. Zhou R, Zhu L, Fu S, Qian Y, Wang D, et al. (2016) Small Diameter Blood Vessels Bioengineered From Human Adipose-derived Stem Cells. *Sci Rep* 6: 35422.
23. Ngo P, Ramalingam P, Phillips JA, Furuta GT (2006) Collagen gel contraction assay. *Methods Mol Biol* 341: 103-109.
24. Yogi A, Rukhlova M, Charlebois C, Tian G, Stanimirovic DB, et al. (2021) Differentiation of Adipose-Derived Stem Cells into Vascular Smooth Muscle Cells for Tissue Engineering Applications. *Biomedicines* Jul 9: 797.
25. Zhang X, Simmons CA, Santerre JP (2018) Alterations of MEK1/2-ERK1/2, IFN $\gamma$  and Smad2/3 associated Signalling pathways during cryopreservation of ASCs affect their differentiation towards VSMC-like cells. *Stem Cell Res* 32: 115-125.
26. Heydarkhan-Hagvall S, Schenke-Layland K, Yang JQ, Heydarkhan S, Xu Y, et al. (2008) Human adipose stem cells: a potential cell source for cardiovascular tissue engineering. *Cells Tissues Organs* 187: 263-274.
27. Rodríguez LV, Alfonso Z, Zhang R, Leung J, Wu B (2006) Clonogenic multipotent stem cells in human adipose tissue differentiate into functional smooth muscle cells. *Proc Natl Acad Sci U S A* 103:12167-12172.
28. Jeon ES, Moon HJ, Lee MJ, Song HY, Kim YM, et al. (2006) Sphingylphosphorylcholine induces differentiation of human mesenchymal stem cells into smooth-muscle-like cells through a TGF-beta-dependent mechanism. *J Cell Sci* 119(Pt 23): 4994-5005.
29. Kim YM, Jeon ES, Kim MR, Jho SK, Ryu SW, et al. (2008) Angiotensin II-induced differentiation of adipose tissue-derived mesenchymal stem cells to smooth muscle-like cells. *Int J Biochem Cell Biol* 40: 2482-2491.
30. Kim YM, Jeon ES, Kim MR, Lee JS, Kim JH (2008) Bradykinin-induced expression of alpha-smooth muscle actin in human mesenchymal stem cells. *Cell Signal* 20: 1882-1889.
31. Mack CP (2011) Signaling mechanisms that regulate smooth muscle cell differentiation. *Arterioscler Thromb Vasc Biol* 31: 1495-1505.
32. Chen R, Dean D (2017) Mechanical properties of stem cells from different sources during vascular smooth muscle cell differentiation. *Mol Cell Biomech* 14: 153-169.
33. Travnickova M, Kasalkova NS, Sedlar A, Molitor M, Musilkova J, et al. (2021) Differentiation of adipose tissue-derived stem cells towards vascular smooth muscle cells on modified poly(L-lactide) foils. *Biomed Mater* 16: 025016.
34. Low EL, Baker AH, Bradshaw AC (2019) TGF $\beta$ , smooth muscle cells and coronary artery disease: a review. *Cell Signal* 53: 90-101.
35. Tsai S, Hollenbeck ST, Ryer EJ, Edlin R, Yamanouchi D, et al. (2009) TGF-beta through Smad3 signaling stimulates vascular smooth muscle cell proliferation and neointimal formation. *Am J Physiol Heart Circ Physiol* 297: H540-H549.
36. Yang P, Troncone L, Augur ZM, Kim SSI, McNeil ME, et al. (2020) The role of bone morphogenetic protein signaling in vascular calcification. *Bone* 141:115542.
37. Wang C, Cen L, Yin S, Liu Q, Liu W, et al. (2010) A small diameter elastic blood vessel wall prepared under pulsatile conditions from polyglycolic acid mesh and smooth muscle cells differentiated from adipose-derived stem cells. *Biomaterials* 31: 621-630.
38. Lee MJ, Kim MY, Heo SC, Kwon YW, Kim YM, et al. (2012) Macrophages regulate smooth muscle differentiation of mesenchymal stem cells via a prostaglandin F $_{2\alpha}$ -mediated paracrine mechanism. *Arterioscler Thromb Vasc Biol* 32: 2733-2740.
39. Liguori GR, Zhou Q, Liguori TTA, Barros GG, Kühn PT, et al. (2019) Directional Topography Influences Adipose Mesenchymal Stromal Cell Plasticity: Prospects for Tissue Engineering and Fibrosis. *Stem Cells Int* 2019: 5387850.
40. Abolhasani S, Rajabibazi M, Khani MM, Parandakh A, Hoseinpoor R (2020) The cooperative effects of micro-grooved topography and TGF- $\beta$ 1 on the vascular smooth muscle cell contractile protein expression of the mesenchymal stem cells. *Differentiation* 115: 22-29.
41. Parandakh A, Anbarlou A, Tafazzoli-Shadpour M, Ardeshiryajimi A, Khani MM (2019) Substrate topography interacts with substrate stiffness and culture time to regulate mechanical properties and smooth muscle differentiation of mesenchymal stem cells. *Colloids Surf B Biointerfaces* 173: 194-201.
42. Park JS, Chu JS, Tsou AD, Diop R, Tang Z, et al. (2011) The effect of matrix stiffness on the differentiation of mesenchymal stem cells in response to TGF- $\beta$ . *Biomaterials* 32: 3921-3930.
43. Teo BK, Wong ST, Lim CK, Kung TY, Yap CH, et al. (2013) Nanotopography modulates mechanotransduction of stem cells and induces differentiation through focal adhesion kinase. *ACS Nano* 7: 4785-4798.
44. Xing F, Li L, Zhou C, Long C, Wu L, et al (2019) Regulation and Directing Stem Cell Fate by Tissue Engineering Functional Microenvironments: Scaffold Physical and Chemical Cues. *Stem Cells Int* 2019: 2180925.
45. Qi YX, Han Y, Jiang ZL (2018) Mechanobiology and Vascular Remodeling: From Membrane to Nucleus. *Adv Exp Med Biol* 1097: 69-82.
46. Rashidi N, Tafazzoli-Shadpour M, Haghighipour N, Khani MM (2018) Morphology and contractile gene expression of adipose-derived mesenchymal stem cells in response to short-term cyclic uniaxial strain and TGF- $\beta$ 1. *Biomed Tech (Berl)* 63: 317-326.

47. Rabbani M, Tafazzoli-Shadpour M, Shokrgozar MA, Janmaleki M, Tey-moori M (2017) Cyclic Stretch Effects on Adipose-Derived Stem Cell Stiffness, Morphology and Smooth Muscle Cell Gene Expression. *Tissue Eng Regen Med* 14: 279-286.
48. Lau S, Klingenberg M, Mrugalla A, Helms F, Sedding D, et al. (2019) Biochemical Myogenic Differentiation of Adipogenic Stem Cells Is Donor Dependent and Requires Sound Characterization. *Tissue Eng Part A* 25: 936-948.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldoopenaccess.us/submit-manuscript>