

## Review

### Debut of Polyglandular Autoimmune Syndrome Type 2 (Schmidt Syndrome) in a Patient with Chronic Kidney Disease of Unknown Etiology in Peritoneal Dialysis

Iyad Abuward Abu-Sharkh\*, Igor Romaniouk Jakovler, Suleyka Puello Martinez, Manuel Fidalgo Díaz, Nisrine Arhda, Marta Pais Seijas and Candido Díaz Rodriguez

Department of Nephrology, University Clinical Hospital of Santiago de Compostela, Spain

#### Abstract

Schmidt Syndrome refers to the combination of autoimmune adrenal insufficiency (Addison's disease) with autoimmune thyroiditis, and is part of a larger syndrome known as autoimmune polyendocrine syndrome type II or polyglandular autoimmune syndrome type II (PAS II). Schmidt Syndrome as a 1:20,000 prevalence in general population with 3:1 ratio of females to males affected. It is autosomal dominant inheritance with variable penetrance. Associated with certain HLA antigens HLA-DR3, HLA-DR4, non HLA gene M-ICA and CTLA-4. The diagnosis of the Schmidt Syndrome is the same as that of the individual disorders. The treatment includes of primary

\*Corresponding author: Iyad Abuward Abu-Sharkh, Nephrology Department, University Clinical Hospital of Santiago de Compostela, Spain, Tel: +34-639021641; E-mail: eyadward@hotmail.com

Citation: Abuward Abu-sharkh I, Jakovler IR, Martinez SP, Díaz MF, Arhda N, et al. (2018) Debut of Polyglandular Autoimmune Syndrome Type 2 (Schmidt Syndrome) in a Patient with Chronic Kidney Disease of Unknown Etiology in Peritoneal Dialysis. J Nephrol Renal Ther 4: 019.

Received: August 03, 2018; Accepted: November 06, 2018; Published: November 21, 2018

Copyright: © 2018 Abuward Abu-sharkh I, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

hypothyroidism and the Addison disease. We present the evolution, the clinic and the treatment of a patient diagnosed with Schmidt Syndrome.

**Keywords:** Addison disease; Hashimoto's thyroiditis; Peritoneal dialysis; Polyglandular autoimmune syndrome; Schmidt Syndrome

#### Introduction

Autoimmune polyglandular syndrome is a rare disease, that is defined by the presence of two or more glandular insufficiencies caused by autoimmune mechanisms and that may be associated with other pathologies and immunological phenomena. It is characterized by the presence of circulating organospecific antibodies and lymphocytic infiltration of the affected gland. It is classified into 4 types. Polyglandular syndrome type 2 (Schmidt Syndrome) includes adrenal insufficiency associated with autoimmune thyroid disease or type 1 diabetes mellitus [1-3].

A 36-year-old woman with end-stage renal disease of unknown etiology in long-standing peritoneal dialysis who went to the emergency room for marked arterial hypotension and was subsequently diagnosed with autoimmune polyglandular syndrome. We present the evolution, the clinic and the treatment of the syndrome.

#### Case Report

A 36-year-old patient with the following background: end-stage renal failure in the peritoneal dialysis program since March 2010, diagnosed in 2009 with autoimmune thyroid disease with fine needle aspiration suggestive of Hashimoto's thyroiditis and presence of anti-TPO antibodies in blood. In September 2014, a total parathyroidectomy and bilateral total thyroidectomy was done. After the biopsy was made, she was diagnosed with classic multifocal papillary carcinoma and confirming lymphocytic thyroiditis. The patient was treated by the endocrinology service with iodine 131 without incidents. In February 2016 she went to the emergency department for presenting vomiting and dizziness, upon arrival the physical examination was: blood pressure 60/40 mmHg, skin pallor, analytically: leukocytes 5400 (4000-10000), eosinophils 8%(0.5-6), hemoglobin 12.7 g / dl (12-16), hematocrit 37.6% (37-49), platelets 224,000(138000-381000), albumin 3 mg / dl (4-5.2), glucose 95 mg / dl (74-105), urea 52 mg/dl (13-50), uric acid 3.6 mg / dl (2.4-6.4) potassium 3.6 mEq / L(3.6-4.9), lactic dehydrogenase 249 IU / ml (200-418), Sodium 127 mEq / L(137-148) thyrotropin stimulating hormone 13.68 mUI / L(0.35-5.5), T4 1.45 ng / dl(0.89-1.76), parathyroid 32 pg / ml (10-55), cortisol 2.3 µg / dl (5.27-22.45), protein C reagent 0.4 mg/dl, negative tumor markers, autoimmunity was negative.

The patient received the following peritoneal dialysis pattern: 3 exchanges of 2000 cc with physioneal 35 to 1.36%, and negative balances of -700-800 ml / day. An echocardiogram was performed with the following findings: undilated cardiac chambers. Hyperdynamic left ventricular function. Chest radiology: rule out acute pathology. Abdominal ultrasound: kidneys of small size 6 cm, difficult to delimit, intra-abdominal fluid in relation to peritoneal dialysis.

Under suspicion of an undiagnosed adrenal insufficiency syndrome, ACTH, cortisol, aldosterone and anti-alpha 21 hydroxylase levels were requested, these being negative important elevation of ACTH (1812 pg/ml) and a low cortisol (2.3 µg/dl). A Synacthen test was done and it was positive, with a clear absence of elevation of plasma cortisol. Water replacement is performed, treatment is initiated Fludrocortisone 0.1/day and Hydrocortisone 50 mg /8 hour, gradually improving the patient, and being discharged home at home with maintenance treatment 7 days later after stabilization. Subsequently, the immunological study reflects the presence of positive anti-adrenal antibodies being diagnosed the patient of autoimmune adrenal insufficiency, diabetes mellitus was ruled out, Anti GAD/64k, Anti-IA, and Anti insulin antibodies were negative. Two sisters of the patient, aged 32 and 40, were studied and were positive with anti-TPO antibodies, one of them asymptomatic and the other with clinical hypothyroidism.

## Discussion

Autoimmune polyglandular syndrome is a rare disorder of autoimmune etiology that affects two or more glands and that may be associated with more disorders of autoimmune origin. It is characterized by immunological phenomena that cause the loss of self-tolerance to the target organ, thus inducing the formation of circulating organo-specific antibodies that subsequently and in the presence of glandular lymphocytic infiltrate causes destruction of the glandular parenchyma causing failure of the affected organ. The polyglandular syndrome (autoimmune polyendocrinopathy) has been classified into 4 types (Table 1):

**Type 1:** Also known as APECED (Autoimmune Polyendocrinopathy, Candidiasis and Ectodermal dystrophy) a very rare entity, as a very low incidence 1/80000, and a very variable prevalence, is more

frequent in populations with a high degree of consanguinity [3]. It has an autosomal recessive pattern of inheritance and is associated with the mutation of the AIRE autoimmune regulatory gene [4], located on chromosome 21q22.3. Its main components are mucocutaneous candidiasis, adrenal insufficiency and hypoparathyroidism, for its diagnosis the syndrome must have at least two of its main components, it can also be associated with other autoimmune disorders such as gonadal insufficiency, vitiligo, and celiac disease alopecia among others. Its main components usually appear at 3-5 years and before 20 years.

**Type 2:** It is the most frequent of the polyglandular syndromes, affects more women 3:1, and as a prevalence of 2-4.5: 100000 inhabitants. Appears in adulthood from 3-4 decade of life, is of autosomal dominant inheritance with incomplete penetrance, it is believed that there is a polygenic predisposition of chromosome 6. In some studies has been a relationship between the appearance of the disease and the presence of alleles B8 and DR3 of HLA and in adrenal insufficiency with DR3 and DR4. This makes it necessary to rule out involvement in first-degree relatives [5]. Its main components are, involvement of the adrenal gland and thyroid gland (Schmidt Syndrome) or the adrenal gland and pancreas (carpenter syndrome) may be associated with other effects of other organs such as gonadal failure, myasthenia, celiac disease [6,7].

**Type 3:** It is the association of thyroid disease with other autoimmune diseases that affect other organs where the adrenal is not found, depending on the affected organs it is classified in / 3A (thyroid + DM1) 3B (thyroid + gastrointestinal diseases), 3B (Thyroid + skin diseases) and 3D (thyroid + collagen diseases) all of autoimmune etiopathogenesis, as a poorly defined inheritance pattern, but seems to be dominant. It appears in adults and is very rare presentation [8].

|  |   |   |                        |
|--|---|---|------------------------|
| <b>Polyglandular Autoimmune Syndrome 1</b> | <ul style="list-style-type: none"> <li>Addison's disease</li> <li>Hypoparathyroidism</li> <li>Mucocutaneous candidosis</li> </ul>         | <ul style="list-style-type: none"> <li>Hypogonadism</li> <li>Alopecia</li> <li>Keratitis</li> <li>Autoimmune hepatitis</li> <li>Vitiligo</li> <li>Alopecia</li> <li>DM1</li> <li>Chronic thyroiditis</li> <li>Malabsorption</li> <li>Pernicious anemia</li> </ul>                     | Occurs in childhood    |
| <b>Polyglandular Autoimmune Syndrome 2</b> | <ul style="list-style-type: none"> <li>Addison disease</li> <li>Thyroid disease</li> <li>Autoimmune</li> <li>Mellitus diabetes</li> </ul> | Vitiligo Atrophic Gastritis Autoimmune Hepatitis Pernicious Anemia Hypogonadism Alopecia Myasthenia gravis  | Adolescence and adults |
| <b>Polyglandular Autoimmune Syndrome 3</b> | 3A (thyroid + DM1) 3B (thyroid + gastrointestinal diseases), 3B (Thyroid + skin diseases) and 3D (thyroid + collagen diseases)            | Other endocrine affection   | Adults                 |
| <b>Polyglandular Autoimmune Syndrome 4</b> | (adrenal involvement + hypogonadism)  | <ul style="list-style-type: none"> <li>Hypogonadism</li> <li>Vitiligo</li> <li>Alopecia</li> <li>Gastritis</li> <li>Atrophic</li> <li>Anemia</li> <li>Pernicious</li> <li>Hepatitis</li> <li>Chronicle</li> <li>Hypophysitis</li> <li>DM1</li> <li>Disease</li> <li>Celiac</li> </ul> | Adults                 |

**Table 1:** Differences between polyglandular syndromes.

**Type 4:** Encompasses endocrine diseases that do not appear in previous classifications such as (adrenal involvement + hypogonadism), is rare and appears in adults, and remains poorly defined genetic patron [9].

Recall that adrenal glands are divided both anatomically and functionally into two parts: cortex and medulla. In turn, the cortex is subdivided into three zones: glomeruli, produced mainly by aldosterone, the main mineral-corticoid, which contributes to the electrolyte balance through the homeostasis of sodium and potassium. The fasciculate zone is responsible for the synthesis of glucocorticoids, such as cortisol, which affects glucose metabolism and normal cellular function [10].

With the destruction of the gland, on the one hand, low levels of cortisone stimulate ACTH-CRH, increasing levels of ACTH causing hyper pigmentation. On the other hand, the deficit of cortisol decreases gluconeogenesis, producing hypoglycemia, asthenia and muscular fatigability. As the glomerular zone is destroyed, plasma renin levels and angiotensin II, increase to maintain a normal aldosterone secretion, until finally there is a deficit in mineralocorticoid secretion [11]. The treatment is hormone replacement, and symptoms treatment [12].

## Conclusion

Autoimmune polyglandular syndrome is a rare disorder of autoimmune etiology that affects two or more glands and that may be associated with more disorders of autoimmune origin. Appears in adulthood from 3-4 decade of life with involvement of the thyroid gland. Its long latent period, is mainly due to the maintenance of the function of the adrenal glands, however, once marked destruction of these occurs, symptoms such as nausea, dizziness, often symptoms that are confused with a perennial picture treated many sometimes with volume repletion. However, sustained hypoglycemia, arterial hypotension and coloration of skin should raise suspicion of a pathology that has not yet been diagnosed. Early detection is considered essential to avoid major complications. The level of ACTH, aldosterone and imaging tests provide valuable data about the disease.

## Disclosure of Potential Conflicts of Interest

The authors declare that there is no conflict of interests.

## Research Involving Human Participants and/or Animals

The authors declare that there was no research involving human and/or animal participants.

## References

1. Garrido MJM, Ponce CG, Risueño MG, Mora A, Carrato A (2007) Syndrome pluriglandular autoinmune: Revisión. *An Med Interna (Madrid, Spain)* 24: 445-452.
2. Betterle C, Zanchetta R (2003) Update on autoimmune polyendocrine syndromes (APS). *Acta Biomed Ateneo* 74: 9-33.
3. Bahceci M, Tuzcu A, Pasa S, Ayyildiz D, Tuzcu S (2004) Polyglandular autoimmune syndrome type III accompanied by common variable immunodeficiency. *Gynecol Endocrinol* 19: 47-50.
4. Esselborn VM, Landing BH, Whitaker J, Williams RR (1956) The syndrome of familial juvenile hypoadrenocorticism, hypoparathyroidism and superficial moniliasis. *J Clin Endocrinol Metab* 16: 1374-1387.
5. Partanen J, Peterson P, Westman P, Aranko S, Krohn K (1994) Major histocompatibility complex class II and III in Addison's disease. MHC alleles do not predict autoantibody specificity and 21-hydroxylase gene polymorphism has no independent role in disease susceptibility. *Hum Immunol* 41: 135-140.
6. Zlotogora J, Shapiro MS (1992) Polyglandular autoimmune syndrome type I among Iranian Jews. *J Med Genet* 29: 824-826.
7. Muir A, She JX (1999) Advances in the genetics and immunology of autoimmune polyglandular syndrome II/III and their clinical applications. *Ann Med Interne* 150: 301-312.
8. Michels AW, Eisenbarth GS (2010) Immunologic Endocrine Disorders. *J Allergy Clin Immunol* 125: 226-237.
9. Eisenbarth GS, Gottlieb PA (2004) Autoimmune Polyendocrine Syndromes. *N Engl J Med* 350: 2068-2079.
10. Schrier RW (2006) Body water homeostasis: Clinical disorders of urinary dilution and concentration. *J Am Soc Nephrol* 17: 1820-1832.
11. Sachdev Y, Morley AR, Wilkinson R, Hall R (1977) Addison's disease with renal microangiopathy and renal failure. *Q J Med* 182: 151-162.
12. Siddiqi SS, Singh SK, Khan SA, Ishtiaq O, Pathan MF, et al. (2012) Guidelines regarding management of adrenal insufficiency in the Holy month of Ramadan. *Indian J Endocrinol Metab* 16: 519-521.



Journal of Anesthesia & Clinical Care  
Journal of Addiction & Addictive Disorders  
Advances in Microbiology Research  
Advances in Industrial Biotechnology  
Journal of Agronomy & Agricultural Science  
Journal of AIDS Clinical Research & STDs  
Journal of Alcoholism, Drug Abuse & Substance Dependence  
Journal of Allergy Disorders & Therapy  
Journal of Alternative, Complementary & Integrative Medicine  
Journal of Alzheimer's & Neurodegenerative Diseases  
Journal of Angiology & Vascular Surgery  
Journal of Animal Research & Veterinary Science  
Archives of Zoological Studies  
Archives of Urology  
Journal of Atmospheric & Earth-Sciences  
Journal of Aquaculture & Fisheries  
Journal of Biotech Research & Biochemistry  
Journal of Brain & Neuroscience Research  
Journal of Cancer Biology & Treatment  
Journal of Cardiology & Neurocardiovascular Diseases  
Journal of Cell Biology & Cell Metabolism  
Journal of Clinical Dermatology & Therapy  
Journal of Clinical Immunology & Immunotherapy  
Journal of Clinical Studies & Medical Case Reports  
Journal of Community Medicine & Public Health Care  
Current Trends: Medical & Biological Engineering  
Journal of Cytology & Tissue Biology  
Journal of Dentistry: Oral Health & Cosmesis  
Journal of Diabetes & Metabolic Disorders  
Journal of Dairy Research & Technology  
Journal of Emergency Medicine Trauma & Surgical Care  
Journal of Environmental Science: Current Research  
Journal of Food Science & Nutrition  
Journal of Forensic, Legal & Investigative Sciences  
Journal of Gastroenterology & Hepatology Research  
Journal of Gerontology & Geriatric Medicine  
Journal of Genetics & Genomic Sciences  
Journal of Hematology, Blood Transfusion & Disorders  
Journal of Human Endocrinology  
Journal of Hospice & Palliative Medical Care  
Journal of Internal Medicine & Primary Healthcare  
Journal of Infectious & Non Infectious Diseases  
Journal of Light & Laser: Current Trends  
Journal of Modern Chemical Sciences  
Journal of Medicine: Study & Research  
Journal of Nanotechnology: Nanomedicine & Nanobiotechnology  
Journal of Neonatology & Clinical Pediatrics  
Journal of Nephrology & Renal Therapy  
Journal of Non Invasive Vascular Investigation  
Journal of Nuclear Medicine, Radiology & Radiation Therapy  
Journal of Obesity & Weight Loss  
Journal of Orthopedic Research & Physiotherapy  
Journal of Otolaryngology, Head & Neck Surgery  
Journal of Protein Research & Bioinformatics  
Journal of Pathology Clinical & Medical Research  
Journal of Pharmacology, Pharmaceutics & Pharmacovigilance  
Journal of Physical Medicine, Rehabilitation & Disabilities  
Journal of Plant Science: Current Research  
Journal of Psychiatry, Depression & Anxiety  
Journal of Pulmonary Medicine & Respiratory Research  
Journal of Practical & Professional Nursing  
Journal of Reproductive Medicine, Gynaecology & Obstetrics  
Journal of Stem Cells Research, Development & Therapy  
Journal of Surgery: Current Trends & Innovations  
Journal of Toxicology: Current Research  
Journal of Translational Science and Research  
Trends in Anatomy & Physiology  
Journal of Vaccines Research & Vaccination  
Journal of Virology & Antivirals

Submit Your Manuscript: <http://www.heraldopenaccess.us/Online-Submission.php>