

Short Review

Introducing New Neurons in the Alzheimer's Disease Brain

González CA*

Facultad de Medicina y Ciencias, Universidad San Sebastián, Puerto Montt 5480000, Chile

Abstract

Alzheimer's disease is the most prevalent neurodegenerative disorder in the elderly population. The patients suffer cerebral atrophy as a consequence of extensive neuronal loss, especially in areas that play a role in memory and cognition. Cell therapies approaches have emerged as promising treatments to regenerate the brain tissue of the patients. Diverse cellular sources have been tested achieving their successful integration in the host brain, or a "bystander" effect, that reduce the pathology associated abnormalities or the activation of endogenous reparative mechanisms. Recent progress in reprogramming techniques shows that the direct conversion of glial cells into neurons *in vivo* is potentially possible in the near future, offering an interesting alternative to classic cell grafts. Regardless the technique used, the "new members" of the regenerated tissue must integrate in the pre-existing neuronal networks and survive in the toxic environment of the Alzheimer's disease brain. This is critical to achieve a long-lasting reversion of the clinical signs in the patients.

Keywords: Alzheimer's disease; Cell therapy; *In situ* reprogramming; Toxic protein aggregates

Alzheimer's Disease

Alzheimer's Disease (AD) is a progressive neurodegenerative disorder highly prevalent in the advanced age population. The typical abnormalities observed in the brain of patients include a progressive cerebral atrophy, extensive neuronal loss (especially in areas implicated in memory and cognition), synaptic alterations, neuroinflammation and the presence of intra- and extracellular protein aggregates in the form of neurofibrillary tangles and senile plaques, respectively [1]. The affected individuals experience a gradual cognitive decline, psychiatric symptoms and problems performing activities of daily living. The disease progresses from mild symptoms of memory loss to severe dementia [2,3]. The exact cause of AD is still a matter of debate. The "amyloid cascade hypothesis" holds the idea that its genesis

*Corresponding author: González CA, Facultad de Medicina y Ciencias, Universidad San Sebastián, Puerto Montt 5480000, Chile, E-mail: cesar.gonzalez@uss.cl

Citation: González CA (2021) Introducing New Neurons in the Alzheimer's Disease Brain. J Stem Cell Res Dev Ther 7: 080.

Received: August 27, 2021; **Accepted:** September 07, 2021; **Published:** September 14, 2021

Copyright: © 2021 González CA. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

is triggered by accumulation of the Amyloid Beta (A β) peptides in the brain, as a consequence of an imbalance between the secretion and clearance of this natural product of metabolism [4]. Nowadays, there is no effective therapy to treat this devastating illness since the current drugs are just palliative [5,6]. This is a very worrisome fact considering the quick aging of the world population. Another important problem is the lack of an early diagnosis method. The clinical signs associated to disease appear when the damage in the brain is already extensive. For this reason, the need to incorporate new neurons to replace the lost ones arises. To achieve this goal several strategies have been tested including the use of diverse cell types and the direct reprogramming of glial cells into neurons.

Cell Therapies

The neurons are postmitotic non-dividing cells that live many decades. Because the brain lacks the capability to regenerate neurons lost because age or disease, cell therapies are appealing approaches to overcome this problem in AD treatment context. Several cell types have been tested including fetal neural stem cells, embryonic stem cell-derived neural stem cells, Mesenchymal Stem Cells (MSCs) and induced Pluripotent Stem Cells (iPSCs). Among all these alternatives, the MSCs and the iPSCs are the most promising options [7,8].

MSCs

MSCs of diverse origin have been tested in several preclinical studies. They are relatively simple to obtain, have little immunogenicity and form no tumors [9]. The graft of these cells has a positive effect in diverse AD associated abnormalities promoting the functional recovery as a consequence of a by-stander effect rather than a cell replacement phenomenon [9,10]. The graft of bone marrow MSCs in double transgenic AD mice (APP/PSN1 mice) reduces the levels of pro-inflammatory cytokines and A β -42 peptide [11]. In another study using a transgenic AD mouse, Yokokawa and colleagues [12] reported that the transplantation of MSCs not only reduces neuroinflammation and A β -42 peptide levels but also A β plaque number in cortex and hippocampus. Additionally, in the same work, the authors show a reduction in oxidative stress, improvement of spatial memory and also evidence that soluble factors produced by MSCs improve the A β clearance by microglia. In another work, MSCs injected in the lateral ventricle in APP/PSN1 mice did not reduce the A β burden and neuronal cell loss but improved the cognitive impairment in the grafted transgenic animals which may be a consequence of an increased number of synapses in the grafted animals [13]. The same authors also found in these animals a decreased astrogliosis and microglial activation. Interestingly, the aberrant activation of microglia in AD plays a role in synapse loss in AD [14]. Additionally, Nakano and co-workers [13] provide evidence that suggests that the reduction of neuroinflammation is provoked by the suppression of NF- κ B expression by the transference of miR-146a from the MSCs into astrocytes [13].

The genetic manipulation of MSCs before the graft procedure is an interesting possibility to over-express soluble factors in the grafted cells. The signaling molecules wnt3a and CXCL1 have been over-expressed in MSCs using adenoviruses before their inoculation in a

double transgenic APP/PS1 mice [15]. These authors show, using this experimental strategy, a reduction of neuroinflammation and a significant alleviation of cognitive impairment in the grafted animals. They show evidence that this phenomenon is due to a better synaptic integrity and increased hippocampal neurogenesis reflected by the high number of proliferating Sox2⁺/Ki67⁺ neuronal precursors and DCX⁺ neuroblasts [15]. This strategy shows that the cells used in the cell therapy approach can become a source of secreted factors that may reduce disease associated abnormalities and/or promote endogenous reparative mechanisms in the receptor patients.

iPSCs

The iPSCs were first created by viral transduction of somatic cells (skin fibroblasts) with a set of transcription factors expressed in pluripotent cells [16,17]. Like Embryonic Stem Cells (ESCs), iPSCs have the ability to generate any cell type of the body including neuronal precursors [18]. Working with iPSCs not only avoid the ethical issues associated with fetal tissue or embryonic stem cells, but also have the advantage that can be generated from somatic cells of the same patient avoiding the immunological rejection upon their transplantation. The potential of these cells has been tested using neuronal precursors derived from human iPSCs (hiPSC-NPs) grafted in the lateral ventricle of neonatal mice. The transplanted cells are able to survive and migrate extensively in the mouse brain including cortex and hippocampus. Most of the cells differentiate in excitatory neurons [19]. The same authors show morphological evidence that the human neurons form synapses with the host cells suggesting their functional integration. This evidence highlights the potential of hiPSC-NPs to be considered in cell therapies strategies. Real and coworkers [20] show additional data in experiments wherein hiPSC-NPs were inoculated in adult mouse somatosensory cortex. Besides the survival and differentiation of the grafted human cells in neurons and glia, these authors show a long-distance projection of human axons and electrophysiological data that demonstrate functional synapses between human and mouse neurons.

It has been reported that iPSC-derived neurons hold a promise to treat neurodegenerative disorders. Dopaminergic neurons derived from iPSCs have been tested in rodent and primate models of Parkinson disease. The grafted animals show better dopamine levels and an improvement of neurological deficits associated to this pathology [21]. In a rat model of Huntington disease, intracerebral transplantation of hiPSC-NPs recover behavioral defects at least in part as a result of restoration of the neuron density [22]. The transplantation of iPSCs in the brain of a transgenic mice with five familial AD mutations (5XFAD) ameliorate the AD associated pathology [23]. Besides the reduction of cognitive deficits, these authors found that the iPSC treated 5XFAD animals show a reduced plaque number, lower levels of A β 40 and A β 42 and a reduction of β - and γ - secretase activities. On the other hand, the transplantation did not reduce the neuronal loss. It is interesting to mention that in this work the iPSCs were produced with a novel method: treating mouse fibroblasts with an embryonic stem cell protein extract. Upon transplantation, these cells were able to differentiate in glial cells and not in neurons suggesting that the iPSCs used in this study have a positive bystander effect on the AD pathology rather than a cell replacement phenomenon [23]. The transplantation of hiPSC-NPs in the hippocampus of a double transgenic mouse carrying two mutations in the APP precursor protein associated to familial cases of AD restored the spatial memory deficits in the animals [24]. These authors found that the grafted human cells differentiated in cholinergic and GABAergic neurons. They suggest

that the recovery of the grafted animals is a combination of a cell replacement phenomenon and a bystander effect by the human cells that stimulate the differentiation of endogenous mouse neuronal precursor in cholinergic neurons [24]. Armijo and co-workers [25], in a very recent work, grafted neuronal precursors generated from mouse iPSCs in the hippocampus of a transgenic AD mouse model carrying three mutations associated to familial AD (3XTg-AD). The grafted cells differentiate into neurons and glia. Importantly, the treated transgenic animals show, and improved learning and memory as assessed by the object location task and Barnes maze tests. This research group show evidence that this phenomenon was a consequence of the amelioration of several aspects linked to AD pathology including improved brain activity and synaptic plasticity, reduction of neuroinflammation, reduction of A β plaques, reduction of insoluble A β protein and a decreased phosphorylated tau protein level. By the other side, this work does not show strong evidence that all these beneficial effects are a consequence of the integration of transplanted cells. They claimed that the most likely mechanism involved is an indirect or "bystander" effect [25].

In situ reprogramming

Besides conversion of somatic cells into a pluripotent state as described by Yamanaka's group [16,17] differentiated somatic cells can be converted directly into another differentiated phenotypes such as cardiomyocytes, hepatocytes, endothelial cells and neurons [26,27]. Neurons have been generated *in vitro* and *in vivo* from fibroblasts and glial cells using specific transcription factors, microRNAs and small molecules [28,29]. The forced expression of three transcription factors, Ascl1, Brn2 and Myt1l (ABM) in mouse fibroblasts using lentiviral vectors, convert these cells into neurons able to generate action potentials and functional synapses [30]. Human astrocytes expressing the same transcription factors under the control of doxycycline promoter were transplanted in the striatum and hippocampus of adult rats. After the administration of doxycycline in the drinking water, the grafted cells differentiated into neurons [31]. Additionally, lentiviral vectors carrying the cDNA for ABM were injected in the striatum of GFAP-Cre transgenic mice that allow the expression of the transgene only in Cre-expressing cells. This experiment show that is possible to convert resident GFAP-expressing astrocytes in NeuN⁺ neurons [31]. Niu and co-workers [32] provide additional evidence of this phenomenon. Using a lentiviral expression system and just a single transcription factor, SOX2, they achieve the conversion of mature astrocytes of the mouse striatum in proliferating neuroblasts. The co-expression of SOX2 with BDNF and noggin or the expression of SOX2 plus the treatment of the animals with valproic acid, a histone deacetylase inhibitor, promote the differentiation of the induced neuroblasts into functional neurons integrated in the host brain [32] the process by which this reprogramming occurs is unclear. Here, we show that a distinct cellular sequence is involved in SOX2-driven *in situ* conversion of adult astrocytes to neurons. This includes ASCL1(+). These works provide evidence that the direct neuronal conversion of somatic cells *in vivo* is possible and highlight the potential of direct conversion of resident glial cells into neurons as a neuronal replacement strategy to treat neurodegenerative disorders like AD [29,33]. In an AD model, induced by intracerebral injection of streptozotocin, the inoculation in the hippocampus of lentiviral particles expressing the MicroRNA-302/367 plus the *i.p.* administration of valproic acid induced the conversion of reactive astrocytes into NeuN⁺ functional neurons [34]. In a very interesting work, Guo and colleagues [35] show the conversion of reactive glial cells into

functional neurons *in vivo* in a model of AD. In this work, the authors generated neurons from reactive astrocytes using a retroviral vector to express a single neural transcription factor, NeuroD1, under the control of GFAP promoter. The rationale behind the use of this type of vector is because retroviruses infect only proliferating cells and not dividing cells such as neurons. The AD model they used was a transgenic mice 5XFAD that have many cortical reactive astrocytes. The injection of GFAP::NeuroD1-GFP viral vector in the cortex of the transgenic animals induce the generation of NeuN+ neurons from resident reactive astrocytes. Electrophysiology tests demonstrated that these neurons are functional and integrated in the brain [35]. It is important to highlight that reactive astrogliosis is a hallmark of AD pathology and is associated with brain damaged areas [1]. The conversion of reactive astrocytes into neurons aloud the generation of new neurons just in the areas where they are needed.

It has been reported that the direct conversion of one phenotype into another is possible using small molecules. Mouse fibroblasts can be converted in functional neurons *in vitro* using a cocktail of small molecules [36]. A recent work, using genetically modified mice that aloud to follow astrocyte fate over the time, show convincing evidence that the astrocytes can be converted into functional neurons integrated in brain circuitry using a combination of small molecules applied with a mini-osmotic pump [37]. Overall, the conversion of glial cells into neurons by genetic or chemical methods have the potential to re-establish the brain neuronal population when the clinical sings of the disease became evident, holding a promise to treat AD, a neurodegenerative disorder with an increasing prevalence in the world population.

The New Neurons Must Survive in A “Toxic” Environment

The grafted or reprogrammed cells must face a diseased environment in the brain of the patients. In this scenario, those cells became exposed to neurotoxic protein aggregates. These structures are generated from A β peptides and hyperphosphorylated tau proteins that undergo a conformational change from a native and functional conformation, that is water-soluble and rich in α -helix domains, to a hydrophobic pathological β -sheet enriched conformation, that have a high tendency to aggregate. This misfolded conformation is acquired spontaneously and can be transmitted from one misfolded molecule to a native one. This template recruitment of additional native proteins generate protein aggregates of diverse size, from small oligomers to large amyloid fibers [38]. During this phenomenon, the native protein loss its normal physiological function and gain a toxic role that triggers neurodegeneration. The intervening mechanisms in this protein toxicity are still a matter of debate [39].

It has been proposed that the pathogenic aggregation of proteins can be transmitted from one cell to another, leading to the spreading of neurodegenerative process through the brain [38,39]. The first evidence that this phenomenon may be possibly came from the post-mortem analysis of brains of patients with Parkinson's disease that were grafted with human fetal mesencephalic dopaminergic neurons in the putamen and caudate nucleus. The grafted neurons display Lewy bodies and Lewy neurites, intracellular amyloid structures composed by aggregated α -synuclein, a protein that play a central role in the pathogenesis of this disease [40]. Additional evidence is provided by a work wherein cultured cells and grafted neurons in mouse brain capture exogenous α -synuclein aggregates [41] and that this phenomenon can induce the misfolding and aggregation (“seeding” effect) of the

endogenous protein [42]. Espuny-Camacho and colleagues, in a very interesting work, grafted human neuronal precursors, derived from human embryonic stem cells, in a transgenic mouse model of AD that display A β deposition and neuroinflammation. The grafted cells differentiated into neurons that survive and integrate in the host, but later on suffer neurodegeneration and death [43]. Before this event, the human neurons show AD associated abnormalities only when they are grafted in transgenic mice and not in wild type animals. Among these alterations they found dystrophic neurites with abnormal accumulation of presynaptic proteins in close association with A β plaques, immunoreactivity to specific antibodies that recognize hyperphosphorylated tau protein and transcriptional changes found in human patients [43]. The question that arises is: is it really worth the introduction of new neurons in the brain of patients if they will die over the time anyway? There are some potential strategies that may confer to the new neurons a more resistant “phenotype” and/or convert them into a source of protective molecules. This is discussed in the next section.

How to Survive in the Diseased Brain?

Monolayer neuronal cultures produced from human iPSCs derived from sporadic or familial cases of AD display AD associated abnormalities such as higher levels of A β peptide and phosphorylated tau (p-tau), endoplasmic reticulum and oxidative stress, down regulation of synaptic proteins and increased apoptosis [44-49]. These findings reveal that neurons derived from AD-specific iPSCs can reproduce AD traits spontaneously *in vitro*. Furthermore, the neurons differentiated from iPSCs produced from somatic cells obtained from familial cases of AD are more vulnerable when they are exposed to A β aggregates *in vitro* [50].

In the case of evaluation of performing a graft of hiPSC-NPs in a patient with a familial form of AD, which have an earlier onset and a more aggressive clinical outcome, there is a higher need that the grafted cells survive longer time. One exciting possible solution of this problem is the use of gene edition tools such as CRISPR/Cas9 [51]. This technique has been used in iPSCs with mutations associated to familial cases of AD to create isogenic cell lines without the mutations. Remarkably, neurons differentiated from cells with the edited genome lack the AD pathological phenotype [51]. It looks that the graft of hiPSC-NPs with the corrected mutations will increase their survival chance considering that these cells develop spontaneously a diseased phenotype that is more susceptible to suffer the toxic effect of protein aggregates. But the evidence presented by Espuny-Camacho and colleagues [43] suggests that the genome edition is not enough to confer full protection because human cells with normal genotype get AD related alterations after grafting them in the brain of AD Tg mice probably because a seeding mechanism as described before. So, the question is: how the grafted cells can avoid the seeding from diseased cells in the AD brain?

Considering that the protein aggregates, conformed by misfolded A β and tau proteins in the case of AD, are toxic to neurons, a strategy to interfere with the aggregation process seem plausible to fight against the cell-to-cell propagation of the disease. Short synthetic peptides know as β -Sheet Breakers (BSBs) prevent the aggregation of A β protein *in vitro* and inhibit the amyloidogenesis *in vivo* in a rat model of A β deposition [52]. After this first pioneering work, improved BSBs has been created with higher efficiency and increased solubility and stability [53]. Interesting, some specific A β -derived synthetic peptides not only inhibit A β aggregation but also tau aggregation as well [54] having the potential to prevent the spreading of A β and tau

pathology over the brain. The genetic manipulation of hiPSC-NPs in order they express anti-aggregation peptides is an appealing option. Using this strategy, the grafted cell may become a source of proteins that will protect themselves and the neighboring cells from the toxic effect of protein aggregates.

References

- Querfurth HW, LaFerla FM (2010) Alzheimer's disease. *N Engl J Med* 362: 329-344.
- Burns A, Iliffe S (2009) Alzheimer's disease. *BMJ* 338: b158.
- Atri A (2019) The Alzheimer's Disease Clinical Spectrum: Diagnosis and Management. *Med Clin North Am* 103: 263-293.
- Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297: 353-356.
- Masters CL, Bateman R, Blennow K, Rowe CC, Sperling RA, et al. (2015) Alzheimer's disease. *Nat Rev Dis Primers* 1: 15056.
- Norins LC (2019) Predicted economic damage from a quick, simple Alzheimer's disease cure. *Med Hypotheses* 133: 109398.
- Han F, Bi J, Qiao L, Arancio O (2020) Stem Cell Therapy for Alzheimer's Disease. *Adv Exp Med Biol* 1266: 39-55.
- Si Z, Wang X (2021) Stem Cell Therapies in Alzheimer's Disease: Applications for Disease Modeling. *J Pharmacol Exp Ther* 377: 207-217.
- Zhao X, Li D, Zhang L, Niu Y, Wang W, et al. (2021) Mesenchymal stem cell therapies for Alzheimer's disease: preclinical studies. *Metab Brain Dis* 2021.
- Qin C, Lu Y, Wang K, Bai L, Shi G, et al. (2020) Transplantation of bone marrow mesenchymal stem cells improves cognitive deficits and alleviates neuropathology in animal models of Alzheimer's disease: a meta-analytic review on potential mechanisms. *Transl Neurodegener* 9: 20.
- Wei Y, Xie Z, Bi J, Zhu Z (2018) Anti-inflammatory effects of bone marrow mesenchymal stem cells on mice with Alzheimer's disease. *Exp Ther Med* 16: 5015-5020.
- Yokokawa K, Iwahara N, Hisahara S, Emoto MC, Saito T, et al. (2019) Transplantation of Mesenchymal Stem Cells Improves Amyloid- β Pathology by Modifying Microglial Function and Suppressing Oxidative Stress. *J Alzheimers Dis* 72: 867-884.
- Nakano M, Kubota K, Kobayashi E, Chikenji TS, Saito Y, et al. (2020) Bone marrow-derived mesenchymal stem cells improve cognitive impairment in an Alzheimer's disease model by increasing the expression of microRNA-146a in hippocampus. *Sci Rep* 10: 10772.
- Subramanian J, Savage JC, Tremblay M-È (2020) Synaptic Loss in Alzheimer's Disease: Mechanistic Insights Provided by Two-Photon in vivo Imaging of Transgenic Mouse Models. *Front Cell Neurosci* 14: 592607.
- Li A, Zhao J, Fan C, Zhu L, Huang C, et al. (2020) Delivery of exogenous proteins by mesenchymal stem cells attenuates early memory deficits in a murine model of Alzheimer's disease. *Neurobiol Aging* 86: 81-91.
- Takahashi K, Yamanaka S (2006) Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors. *Cell* 126: 663-676.
- Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, et al. (2007) Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined Factors. *Cell* 131: 861-872.
- Zholudeva LV, Jin Y, Qiang L, Lane MA, Fischer I (2021) Preparation of Neural Stem Cells and Progenitors: Neuronal Production and Grafting Applications. *Methods Mol Biol* 2311: 73-108.
- Chen C, Kim W-Y, Jiang P (2016) Humanized neuronal chimeric mouse brain generated by neonatally engrafted human iPSC-derived primitive neural progenitor cells. *JCI Insight* 1: e88632.
- Real R, Peter M, Trabalza A, Khan S, Smith MA, et al. (2018) In vivo modeling of human neuron dynamics and Down syndrome. *Science* 362: eaau1810.
- Han F, Hu B (2020) Stem Cell Therapy for Parkinson's Disease. *Adv Exp Med Biol* 1266: 21-38.
- Yoon Y, Kim HS, Hong CP, Li E, Jeon I, et al. (2020) Neural Transplants From Human Induced Pluripotent Stem Cells Rescue the Pathology and Behavioral Defects in a Rodent Model of Huntington's Disease. *Front Neurosci* 14: 558204.
- Cha M-Y, Kwon Y-W, Ahn H-S, Jeong H, Lee YY, et al. (2017) Protein-Induced Pluripotent Stem Cells Ameliorate Cognitive Dysfunction and Reduce A β Deposition in a Mouse Model of Alzheimer's Disease. *Stem Cells Transl Med* 6: 293-305.
- Fujiwara N, Shimizu J, Takai K, Arimitsu N, Ueda Y, et al. (2015) Cellular and molecular mechanisms of the restoration of human APP transgenic mouse cognitive dysfunction after transplant of human iPSC cell-derived neural cells. *Exp Neurol* 271: 423-431.
- Armijo E, Edwards G, Flores A, Vera J, Shah Nawaz M, et al. (2021) Induced Pluripotent Stem Cell-Derived Neural Precursors Improve Memory, Synaptic and Pathological Abnormalities in a Mouse Model of Alzheimer's Disease. *Cells* 10: 1802.
- Kelaini S, Cochrane A, Margariti A (2014) Direct reprogramming of adult cells: avoiding the pluripotent state. *Stem Cells Cloning* 7: 19-29.
- Lee S, Kim JE, Johnson BA, Andukuri A, Yoon Y-S (2017) Direct reprogramming into endothelial cells: a new source for vascular regeneration. *Regen Med* 12: 317-320.
- Egawa N, Suzuki H, Takahashi R, Hayakawa K, Li W, et al. (2020) From in vitro to in vivo reprogramming for neural transdifferentiation: An approach for CNS tissue remodeling using stem cell technology. *J Cereb Blood Flow Metab* 40: 1739-1751.
- Yavarpour-Bali H, Ghasemi-Kasman M, Shojaei A (2020) Direct reprogramming of terminally differentiated cells into neurons: A novel and promising strategy for Alzheimer's disease treatment. *Prog Neuropsychopharmacol Biol Psychiatry* 98: 109820.
- Vierbuchen T, Ostermeier A, Pang ZP, Kokubu Y, Südhof TC, et al. (2010) Direct conversion of fibroblasts to functional neurons by defined factors. *Nature* 463: 1035-1041.
- Torper O, Pfisterer U, Wolf DA, Pereira M, Lau S, et al. (2013) Generation of induced neurons via direct conversion in vivo. *Proc Natl Acad Sci U S A* 110: 7038-7043.
- Niu W, Zang T, Smith DK, Vue TY, Zou Y, et al. (2015) SOX2 reprograms resident astrocytes into neural progenitors in the adult brain. *Stem Cell Reports* 4: 780-794.
- Torper O, Götz M (2017) Brain repair from intrinsic cell sources: Turning reactive glia into neurons. *Prog Brain Res* 230: 69-97.
- Ghasemi-Kasman M, Shojaei A, Gol M, Moghadamnia AA, Baharvand H, et al. (2018) miR-302/367-induced neurons reduce behavioral impairment in an experimental model of Alzheimer's disease. *Mol Cell Neurosci* 86: 50-57.
- Guo Z, Zhang L, Wu Z, Chen Y, Wang F, et al. (2014) In vivo direct reprogramming of reactive glial cells into functional neurons after brain injury and in an Alzheimer's disease model. *Cell Stem Cell* 14: 188-202.
- Li X, Zuo X, Jing J, Ma Y, Wang J, et al. (2015) Small-Molecule-Driven Direct Reprogramming of Mouse Fibroblasts into Functional Neurons. *Cell Stem Cell* 17: 195-203.
- Ma Y, Xie H, Du X, Wang L, Jin X, et al. (2021) In vivo chemical reprogramming of astrocytes into neurons. *Cell Discov* 7: 12.
- Davis AA, Leys CEG, Holtzman DM (2018) Intercellular Spread of Protein Aggregates in Neurodegenerative Disease. *Annu Rev Cell Dev Biol* 34: 545-568.

39. Jucker M, Walker LC (2013) Self-propagation of pathogenic protein aggregates in neurodegenerative diseases. *Nature* 501: 45-51.
40. Li J-Y, Englund E, Holton JL, Soulet D, Hagell P, et al. (2008) Lewy bodies in grafted neurons in subjects with Parkinson's disease suggest host-to-graft disease propagation. *Nat Med* 14: 501-503.
41. Hansen C, Angot E, Bergström A-L, Steiner JA, Pieri L, et al. (2011) α -Synuclein propagates from mouse brain to grafted dopaminergic neurons and seeds aggregation in cultured human cells. *J Clin Invest* 121: 715-725.
42. Luk KC, Song C, O'Brien P, Stieber A, Branch JR, et al. (2009) Exogenous alpha-synuclein fibrils seed the formation of Lewy body-like intracellular inclusions in cultured cells. *Proc Natl Acad Sci U S A* 106: 20051-20056.
43. Espuny-Camacho I, Arranz AM, Fiers M, Snellinx A, Ando K, et al. (2017) Hallmarks of Alzheimer's Disease in Stem-Cell-Derived Human Neurons Transplanted into Mouse Brain. *Neuron* 93: 1066-1081.
44. Israel MA, Yuan SH, Bardy C, Reyna SM, Mu Y, et al. (2012) Probing sporadic and familial Alzheimer's disease using induced pluripotent stem cells. *Nature* 482: 216-220.
45. Shi Y, Kirwan P, Smith J, MacLean G, Orkin SH, et al. (2012) A human stem cell model of early Alzheimer's disease pathology in Down syndrome. *Sci Transl Med* 4: 124ra29.
46. Kondo T, Asai M, Tsukita K, Kutoku Y, Ohsawa Y, et al. (2013) Modeling Alzheimer's disease with iPSCs reveals stress phenotypes associated with intracellular A β and differential drug responsiveness. *Cell Stem Cell* 12: 487-496.
47. Dashinimaev EB, Artyuhov AS, Bolshakov AP, Vorotelyak EA, Vasiliev AV (2017) Neurons Derived from Induced Pluripotent Stem Cells of Patients with Down Syndrome Reproduce Early Stages of Alzheimer's Disease Type Pathology in vitro. *J Alzheimers Dis* 56: 835-847.
48. Tew J (2019) Human iPSC application in Alzheimer's disease and Tau-related neurodegenerative diseases. *Neurosci Lett* 699: 31-40.
49. Arber C, Toombs J, Lovejoy C, Ryan NS, Paterson RW, et al. (2020) Familial Alzheimer's disease patient-derived neurons reveal distinct mutation-specific effects on amyloid beta. *Mol Psychiatry* 25: 2919-2931.
50. Armijo E, Gonzalez C, Shahnawaz M, Flores A, Davis B, et al. (2017) Increased susceptibility to A β toxicity in neuronal cultures derived from familial Alzheimer's disease (PSEN1-A246E) induced pluripotent stem cells. *Neurosci Lett* 639: 74-81.
51. Barman NC, Khan NM, Islam M, Nain Z, Roy RK, et al. (2020) CRISPR-Cas9: A Promising Genome Editing Therapeutic Tool for Alzheimer's Disease-A Narrative Review. *Neurol Ther* 9: 419-434.
52. Soto C, Sigurdsson EM, Morelli L, Kumar RA, Castaño EM, et al. (1998) Beta-sheet breaker peptides inhibit fibrillogenesis in a rat brain model of amyloidosis: implications for Alzheimer's therapy. *Nat Med* 4: 822-826.
53. Francioso A, Punzi P, Boffi A, Lori C, Martire S, et al. (2015) β -sheet interfering molecules acting against β -amyloid aggregation and fibrillogenesis. *Bioorg Med Chem* 23: 1671-1683.
54. Gorantla NV, Sunny LP, Rajasekhar K, Nagaraju PG, Cg PP, et al. (2021) Amyloid- β -Derived Peptidomimetics Inhibits Tau Aggregation. *ACS Omega* 6: 11131-11138.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>