

Review Article

Correlation between Clinical Features and Laboratory Parameters and Outcome in Acute Copper Sulphate Poisoning: A Cross-Sectional Observational Study

Dutta D¹, Gomes RR^{2**}, Hasan FMM³, Saha S² and Newaz M⁴

¹Ad-din Women's Medical College Hospital, Dhaka, Bangladesh

²Jashore Medical College Hospital, Jashore, Bangladesh

³Ad-din Sakina Women's Medical College Hospital, Jashore, Bangladesh

⁴Indoor Medical Officer, Rajshahi Medical College Hospital, Rajshahi, Bangladesh

Abstract

Background: Ingestion of copper sulphate is a common mode of poisoning in the Indian subcontinent. Cases are mainly suicidal in nature. The clinical course of the copper sulphate intoxicated patient is often complex involving intravascular hemolysis, jaundice and renal failure. The treatment is mainly supportive. Mortality is quite high in severe cases if treatment is not adequate and prompt.

Methods and materials: This cross sectional, observational, prospective study was conducted on 50 patients with acute copper sulphate poisoning attended in Medicine department of Khulna Medical College Hospital from January 2020 to June 2020.

Results: Almost all patients of copper sulphate poisoning, presented with gastrointestinal symptom like nausea 50(100%), vomiting 49 (98%) and abdominal pain 26 (52%). Development of anemia 17 (34%) and signs of hepatotoxicity like jaundice 18(36%), signs of dehydration 7 (14%), features of shock 8 (16%) and chest findings like tachypnoea, crepitations 9 (18%) were common. On investigation, Hemoglobin percentage below 10 gm/dl in 12 (24%), neutrophilic

leukocytosis 18 (36%), Platelet count below 100000/cu mm blood in 3 (6%). features of hemolysis 16 (32%), haemoglobinuria 17 (34%) and haematuria 15 (30%), S. bilirubin was elevated in 23 (46%) of patients. Serum ALT level was elevated in 17 (34%) of patients of which 13 (26%) were found as moderate to severely raised. both serum creatinine and blood urea were elevated in 16 (32%) of patients.

Conclusion: Acute copper sulphate poisoning is common in our country which is mainly suicidal in nature. As death occurs due to multi organ failure, while gastrointestinal, intravascular hemolysis, hepatic and renal toxicities significantly contribute to it. Management in High dependency unit is also required.

Keywords: Hemoglobinuria; Hematuria; Hepatotoxicity; Intravascular hemolysis; Renal failure

Introduction

Copper sulphate forms bright blue crystals containing five molecules of water [CuSO₄ .5H₂ O]. It is commonly known as "Blue Vitriol" or "Blue Stone". Usually the people of the southern region of Bangladesh uses copper sulphate as a fungicidal agent [1]. It is used chiefly for agricultural purposes as a pesticide and in leather industry. It was also being used as a precipitator in heavy metal poisoning and was used to treat gastric and topical exposure to phosphorous. It has a nauseous and metallic taste. Solutions are acid to litmus, freely soluble in water [2,3]. It is consumed mainly with suicidal intentions. Accidental poisonings have been reported from children as well [4,5].

Approximately 30% of ingested copper is absorbed from the gastrointestinal tract [6]. In blood, copper is initially albumin-bound and transported via the hepatic portal circulation to the liver where it is incorporated into ceruloplasmin. Copper is present in serum in two forms; 93% is tightly bound to ceruloplasmin and 7% is loosely bound to albumin [7]. The copper-albumin complex represents the toxicological active portion of the serum copper [2]. Systemic transport of copper from liver is primarily as ceruloplasmin, which appears to donate copper to tissues. Copper is distributed to all tissues with the highest concentrations in liver, heart, brain, kidneys and muscle. Intracellular copper is predominantly bound to metallothionein. Fecal and biliary excretion accounts for 80 percent of excreted copper. Approximately four percent is excreted in the urine [6].

In acute poisoning, albumin, rather than ceruloplasmin, binds the excess copper. The liver is the major site of deposition of copper following large ingestion. Lethal dose is about 10-20g [7]. The copper content in normal adult liver ranges from 18-45 mg/g dry weight. When the concentration of hepatic copper is greater than 50 mg/g dry weight, liver cell necrosis occurs with release of large amount of copper into the serum. This released copper is rapidly taken up by erythrocytes and results in oxidative damage and may result in hemolysis of RBCs [2]. It is proposed that free reduced copper in the cell binds to sulphhydryl groups and inactivates enzymes such as glucose-6-phosphate dehydrogenase and glutathione reductase [8]. Intravascular hemolysis appears 12-24h following ingestion of copper sulphate.

***Corresponding author:** Gomes RR, Jashore Medical College Hospital, Jashore, Bangladesh, E-mail: rrichi.dmc.k56@gmail.com

Citation: Dutta D, Gomes RR, Hasan FMM, Saha S, Newaz M (2022) Correlation between Clinical Features and Laboratory Parameters and Outcome in Acute Copper Sulphate Poisoning: A Cross-Sectional Observational Study. J Toxicol Cur Res 6: 023.

Received: May 03, 2022; **Accepted:** May 11, 2022; **Published:** May 18, 2022

Copyright: © 2022 Dutta D, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Jaundice in copper sulphate poisoning is partly hepatic in origin in addition to hemolysis [9]. Jaundice appears on the second or third day following ingestion. Liver damage has been attributed to liver mitochondrial dysfunction due to oxidized state. Intravascular hemolysis plays a major role in the pathogenesis of renal failure [8-10]. The hem pigment released due to hemolysis and direct toxic effect of copper released from lysed red cells contribute to tubular epithelial damage of the kidney. Severe vomiting, diarrhoea, lack of replacement of fluid and gastrointestinal bleed, leading to hypotension could also contribute to renal failure [10]. Renal complications are usually seen on the third or the fourth day and onwards after the poisoning [11].

The incidence of copper sulphate poisoning varies at different geographical areas depending on the local use of copper sulphate and the availability of other suicidal poisons. Diagnosis is based on history of exposure. Mainstay of treatment is supportive, including careful fluid therapy and vasopressors as needed, blood transfusion if needed and methylene in symptomatic methemoglobinemia. Chelation therapy given in most cases. The role of dialysis is limited to the management of associated renal failure. High mortality is noted in severe cases due to methemoglobinemia, hepatotoxicity and renal failure.

Materials and Methods

This cross sectional, observational, prospective study was carried out in Medicine department of Khulna Medical College Hospital (KMCH) from January, 2020 to June 2020. Total 50 cases of acute copper sulphate poisoning were selected. Data were processed and analyzed using SPSS (Statistical Package for Social Science) 15.0.

Inclusion criteria

- All patients with acute copper sulphate poisoning admitted in Medicine wards of KMCH.
- Voluntarily given consent.

Exclusion criteria

- Not willing to give informed consent.
- Patient is suffering from Wilson's disease, chronic liver disease and chronic kidney disease.

Results

Table 1 shows that almost all patients of copper sulphate poisoning, presented with gastrointestinal symptom like abdominal pain 26 (52%), nausea 50 (100%), and vomiting 49 (98%) A significant proportion of patients presented with high coloured urine 20 (40%) which developed initially and subsequently.

Presenting Feature	No. of Patient	Percentage (%)
Nausea	50	100(%)
Vomiting	49	98(%)
Abdominal pain	26	52(%)
High coloured urine	20	40(%)
Jaundice	8	16(%)

Table 1: Showing clinical presentation of the patients.

Table 2 shows that anemia 17 (34%) and jaundice 18 (36%) developed on 2nd or 3rd day following copper sulphate poisoning. Development of signs of dehydration 7 (14%), features of shock 8 (16%) and chest findings like tachypnoea, crepitations 9 (18%) were not uncommon.

Physical findings on subsequent days	No. of patients	Percentage (%)
Anemia	17	34 %
Jaundice	18	36 %
Signs of dehydration	7	14 %
Features of shock	8	16 %
Chest findings(e.g - crepitations)	9	18 %

Table 2: Showing physical findings on subsequent days during hospital stay.

Table 3 shows that anemia is a common feature. Hemoglobin percentage was found below 10 gm/dl in 12 (24%) of patients. It is due to hemolysis or blood loss. Neutrophilic Leukocytosis were found on 18 (36%) of patients. Platelet count was found below 100000/cu mm blood in a minority patient 3 (6%).

Complete Blood Count	No. of patients	Percentage (%)	
Percentage of hemoglobin	<10 gm/dl	12	24.0 %
	>10 gm/ dl	38	76.0 %
TC of WBC	>11000/cu mm	18	36.0 %
	<11000/ cu mm	32	64.0 %
DC of WBC (Percentage of Neutrophil)	>70%	18	36.0 %
	<70%	32	64.0 %
Platelet count	<100000 /cu mm	3	6.0 %
	>100000/cu mm	47	94.0 %

Table 3: Complete Blood Count among the patients.

Table 4 shows that features of hemolysis was present on 16 (32%) of patient.

PBF findings	Frequency	Percent
Features of hemolysis	16	32.0%
Normal study	34	68.0%
Total	50	100.0%

Table 4: Peripheral blood film among the patients.

Table 5 shows that hemoglobinuria and hematuria developed in 17 (34%) and 15 (30%) of the patients respectively.

Urine Analysis	No. of patients	Percentage (%)	
Albuminuria	present	5	10.0 %
	Absent	45	90.0 %
Hemoglobinuria	present	17	34.0 %
	Absent	33	66.0 %
Hematuria	present	15	30.0 %
	absent	35	70.0 %

Table 5: Urine analysis among the patients.

Table 6 shows that serum bilirubin was elevated in 23 (46%) of patients. Serum ALT level was elevated in 17 (34%) of patients of which 26% were found as moderate to severely raised. LFT were done at the 4th day of copper sulphate ingestion.

Liver Function Test		No. of patients	Percentage (%)
Serum bilirubin	Increased (>1.1 mg/dl)	23	46.0
	Normal (up to 1.1 mg/dl)	27	54.0
Serum alanine aminotransferase (ALT)	Moderate to severely raised (>80 u/l)	13	26.0
	Mildly raised (41-80 u/l)	4	8.0
	Normal (up to 40u/l)	33	66.0

Table 6: Liver function test among the patients.

Table 7 shows that both serum creatinine and blood urea were elevated in 16 (32%) of patients. RFT were done at the 4th day of copper sulphate ingestion.

Renal Function Test		No. of patients	Percentage (%)
Serum creatinine	Increased (>1.2 mg/dl)	16	32.0 %
	Normal (up to 1.2 mg/ dl)	34	68.0 %
Blood urea	Increased (>1.2 mg/dl)	16	32.0 %
	Normal (upto 1.2 mg/ dl)	34	68.0 %

Table 7: Renal function test among the patients.

Among 16 patients of raised serum creatinine, 14 patients improved and 2 patients died. Among 34 patients of normal serum creatinine, all of them (100%) improved. There is statistically significant association (p-value=0.035) between raised serum creatinine and outcome of treatment of patients in acute copper sulphate poisoning (Table 8).

Outcome of treatment of patient	Serum creatinine			P-Value
	Increased (>1.2 mg/dl)	Normal (upto 1.2 mg/ dl)	Total	
Improved	14	34	48	0.035
Death	2	0	2	
Total	16	34	50	

Table 8: Co-relation between serum creatinine and outcome of treatment.

Chi-square (χ^2) Test was employed to analyse the data. P-value <0.05 was considered statistically significant. Among 13 patients of Moderate to severely raised (>80 u/l) serum ALT, 11 patients had improved and 2 patients had died. Among 4 patients of mildly raised (41-80 u/l) serum ALT and 33 patients of normal serum creatinine, all of them (100%) had improved. There is statistically significant association (p-value=0.05) between raised serum ALT and outcome of patients in acute copper sulphate poisoning (Table 9).

Chi-square (χ^2) Test was employed to analyze the data. P-value <0.05 was considered statistically significant. Among 16 patients with features of hemolysis on peripheral blood film, 14 patients improved and 2 patients died. Among 34 patients with normal peripheral

Outcome of patient				
Serum alanine aminotransferase (ALT)	Improved	Death	Total	P-Value
Moderate to severely raised (>80 u/l)	11	2	13	0.05
Mildly raised (41-80 u/l)	4	0	4	
Normal (up to 40u/l)	33	0	33	
Total	48	2	50	

Table 9: Co-relation between serum alanine aminotransferase (ALT) and outcome of treatment of patient.

blood film, all patients (100%) improved. There is statistically significant association (p-value=0.035) between presence of features of hemolysis on peripheral blood film and outcome of patients in acute copper sulphate poisoning (Table 10).

Outcome of patient				
Peripheral blood picture	Improved	Death	Total	P-Value
Features of hemolysis	14	2	16	0.035
Normal study	34	0	34	
Total	48	2	50	

Table 10: Co-relation between features on peripheral blood picture and outcome of patient.

Chi-square (χ^2) Test was employed to analyze the data. P-value <0.05 was considered statistically significant.

Discussion

Acute copper sulphate poisoning is a common mode of poisoning in southern region of Bangladesh which are mainly suicidal in nature [1] usually the people of this region use copper sulphate as a fungicidal agent. The incidence of acute copper sulphate poisoning varies at different geographical areas depending on the local use of copper sulphate and the availability of other suicidal poisons. The lowest dose copper sulphate that is toxic when ingested is 11mg /kg [12]. Akin-tonwa et al., 1996 [13] claimed 10-20 gm. copper sulphate to be a “definitely fatal” dose. In a review of 123 cases Ahsan et al., 1994 [1] observed an “unpredictable” outcome in those consuming less than 50 gm while 100 gm was “invariably” fatal. Following ingestion of copper sulphate gastrointestinal symptoms appear first and is always present. In present study, gastrointestinal symptoms like nausea 50 (100%), and vomiting 49 (98%), abdominal pain 26 (52%) were observed in patients, which is consistent with the previous studies [SH] Mollick et al., 2011 [14], Chowdhury et al., 1961 [15], singh and singh et al., 1968 [9], Chugh KS et al., 1977 [10].

Jaundice developed on 2nd or 3rd day following ingestion of copper sulphate. Clinically it is detected in 36% patients (18 out of 50). Jaundice may be hemolytic or hepatocellular or mixed. Serum bilirubin was elevated in 23 (46%) of patients. In previous study, Wahal et al., 1963 [16] jaundice was found on 36% of patient and Agarwal SK et al., 1993 [17] jaundice was found 58% of patients. Serum ALT level was elevated in 17 (34%) of patients of which 13 were found as moderate to severely raised which is consistent with Singh M et al., 1968 [9], Ashraf I et al., 1970 [18], Wahal PK et al., 1976 [16]. In a study, SH Mollick et al., 2011 [14] stated Serum ALT & serum bilirubin were elevated in 40% and 37% patients respectively. Among

13 patients of Moderate to severely raised (>80 u/l) serum ALT, 11 patients improved and 2 patients died. Among 4 patients with mildly raised (41-80 u/l) serum ALT and 33 patients with normal serum ALT, all of them (100%) had improved. There is statistically significant association (p-value=0.05) between raised serum ALT and outcome of patients in acute copper sulphate poisoning (Table 9).

Intravascular hemolysis can start as early as within the first 24 hours since ingestion and is due to the direct oxidative damage to erythrocyte membranes. The hemolysis can be rapid and severe with drastic drops in the hemoglobin level. In this study, Hemoglobin percentage was found below 10 gm/dl in 12 (24%) of patients. Which may be due to copper induced hemolysis (increased total bilirubin and evidence from blood picture). A significant proportion of patients developed high coloured urine 20 (40%). which may be due to hemolysis or jaundice. Hemoglobinuria, hematuria and albuminuria were developed in 17 (34%), 15 (30%) and 5 (10%) of the patients respectively. In previous study, Chugh KS et al., 1977 [14] and Agarwal SK et al., 1993 [17] found intravascular hemolysis ranged from 47-65% in two case series. Among 16 patients with features of hemolysis on peripheral blood film, 14 patients improved and 2 patients died. Among 34 patients of normal peripheral blood film, all of them (100%) improved. There is statistically significant association (p-value=0.035) between presence of features of hemolysis on peripheral blood film and outcome of patients in acute copper sulphate poisoning (Table 10).

Renal complications were observed usually after 48hours. The possible mechanisms of kidney damage include; pre-renal failure due to dehydration (vomiting, diarrhea, reduced fluid intake), hemoglobinuria, sepsis, rhabdomyolysis, direct copper toxicity on proximal tubules and secondary effects of multi organ dysfunction. In this study, serum creatinine and blood urea were elevated in 16 (32%) patients. High coloured urine developed in 20 (40%) patients (it may be due to hematuria and/or hemoglobinuria). Hemoglobinuria, hematuria and albuminuria developed in 17 (34%), 15(30%) and 5 (10%) patients respectively.

In some previous studies, Dash SC et al., 1989 [8] , Chugh KS et al., 1977 [10] and Mehta A et al., 1985 [11] described that acute renal failure developed in 20-40% of patients with acute copper sulphate poisoning and urinary abnormalities detected are oliguria, anuria, albuminuria, hemoglobinuria and hematuria. Among 16 patients with raised serum creatinine, 14 patients improved and 2 patients died. Among 34 patients with normal serum creatinine, all of them (100%) improved. There is statistically significant association (p-value=0.035) between raised serum creatinine and outcome of patients in acute copper sulphate poisoning (Table 8).

Conclusion

Copper sulphate is a common form of suicidal agent with significant mortality. It is a readily available substance can be taken easily. Acute copper sulphate poisoning is a rare event and uncommon worldwide except Indian subcontinents. It is most prevalent in southern region of Bangladesh. Oral copper sulphate ingestion has significant effects on gastrointestinal, hematological, hepatic and renal system. Gastrointestinal symptoms like nausea, vomiting, abdominal pain frequently occurs and appears. Intravascular hemolysis usually developed within 12 - 24 hours of ingestion. Appearance of jaundice is common. Most common renal toxicities are hematuria, hemoglobinuria, albuminuria, elevated blood urea, elevated serum creatinine

level and acute renal failure. Although death occurs due to multi organ failure, gastrointestinal, intravascular hemolysis, hepatic and renal toxicities significantly contributes to it.

Limitations of Study

The sample size was relatively small. No follow-up could be carried out to repeat the investigations. Actual measurement of dose of copper sulphate is not possible. Advanced investigation facilities are limited. No advanced life supports were available.

Funding

Self-funded.

Conflict of Interests

None declared.

References

1. Ahasan HA, Chowdhury MA, Azhar MA, Rafiqueuddin AK (1994) Copper sulphate poisoning. *Trop Doct* 24: 52-53.
2. Barceloux DG (1999) Copper. *J Toxicol Clin Toxicol* 37: 217-230.
3. Gupta PS, Bhargava SP, Sharma ML (1962) Acute copper sulphate poisoning with special reference to its management with corticosteroid therapy. *J Assoc Physicians India* 10: 287-292.
4. Sharma NL, Singh RN, Natu NK (1967) Accidental poisoning in infancy and childhood. *J Indian Med Assoc* 48: 20-25.
5. Blundell S, Curtin J, Fitzgerald D (2003) Blue lips, coma and haemolysis. *J Paediatr Child Health* 39: 67-68.
6. Dowdy RP (1969) Copper metabolism. *Am J Clin Nutr* 22: 887-892.
7. Ellenhorn MJ (1997) Ellenhorn's Medical Toxicology: Metals and Related Compounds. In : *Diagnosis and treatment of human poisoning* 2: 1554-1556.
8. Singh M, Singh G (1968) Biochemical changes in Blood in cases of acute copper sulphate poisoning. *J Indian Med Assoc* 50: 549-554.
9. Chugh KS, Sharma BK, Singhal PC (1977) Acute renal failure following copper sulphate intoxication. *Postgrad Med J* 53: 18-23.
10. Mehta A, Patney NL, Bhati DP, Singh SP (1985) Copper sulphate poisoning - Its impact on Kidneys. *J Indian Med Assoc* 83: 108-110.
11. Mehta A, Patney NL, Bhati DP, Singh SP (1985) Copper sulphate poisoning - Its impact on Kidneys. *J Indian Med Assoc* 83:108-110.
12. <http://extoxnet.orst.edu/pips/coppersu.htm>
13. Akintonwa A, Mebadeje AFB, Odutola TA (1989) Fatal poisonings by copper sulphate ingested from "spiritualwater". *Vet Hum Toxicol* 31: 453-454.
14. Mollick SH, Mollick KA, Bakar A, Miah M (2011) Burden and outcome of acute copper sulphate poisoning in a teaching hospital Bang Med J (Khulna) 44 : 7-10.
15. Chowdhury AK, Ghosh S, Pal D (1961) Acute copper sulphate poisoning. *J Indian Med Assoc* 36: 330-336.
16. WahalPK, Lahiri B, Mathur KS, Kehar U, Wahi PN (1963) Acute copper sulphate poisoning. *J Assoc Physicians India* 11: 93-103.
17. Agarwal SK, Tiwari SC, Dash SC (1993) Spectrum of poisoning requiring hemodialysis in tertiary care hospital in India. *Int J Artif Organs* 16: 20-22.
18. Ashraf I (1970) Hepatic derangements (biochemical) in acute copper sulphate poisoning. *J Indian Med Assoc* 55: 341-342.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>