



Research Article

Advances in Fasting Aerobic Exercise Research

Jiang An Lu^{1*}, Yi Bo Wang², Yong Min Niu³, Ma Chi Cheng Bao⁴ and Zhi Yan Li¹

¹Department of Sports Human Science, Physical Education Institute, Yunnan Normal University, Kunming, China

²Department of Human anatomy and Histoembryology, College of Basic Medicine, Jinzhou Medical University, China

³Department of Neurobiology, School of Basic Medical Sciences, Kunming Medical University, Kunming, China,

⁴Department of Neurobiology and Animal, Kunming Medical University, Kunming, China

Abstract

As of 2020, obesity has become a worldwide crisis and the rate of obesity in China has exceeded 50%, there's no doubt obesity has become a major problem that plagues the healthy development of China's economy and the well-being and health of national life. Not only exercise is a good doctor, but also to solve the problem of obesity efficient means. In order to reduce the obesity rate, we have taken many exercise methods, such as resistance exercise, aerobic exercise, special exercise, but most of them have the problems of inability to adhere to and inefficiency. At this time, fasting aerobic as a new method of weight loss came into the forefront, a large number of experimental studies have been conducted on fasting aerobic abroad, and thanks to the points that have been found, fasting aerobic has been used as an effective method to cope with cardiovascular diseases, metabolic diseases and obesity abroad. On the basis of a thorough reading of domestic and international literature, the authors summarize the existing studies on fasting aerobics, hoping to point out the direction for subsequent research, so that everyone can fully understand the benefits of fasting aerobics and actively promote it.

Keywords: Fasting aerobic; Glucose and lipid metabolism; Insulin resistance

*Corresponding author: Jiang An Lu, Department of Sports Human Science, Physical Education Institute, Yunnan Normal University, Kunming, China, Tel: +86 18896652856; E-mail: 810138303@qq.com

Citation: Lu JA, Wang YB, Niu YM, Bao MCC, Li ZY (2022) Advances in Fasting Aerobic Exercise Research. J Altern Complement Integr Med 8: 257.

Received: June 20, 2022; **Accepted:** June 22, 2022; **Published:** June 29, 2022

Copyright: © 2022 Lu JA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Background

Obesity has been confirmed to be closely related to a series of cardiovascular and cerebrovascular diseases, such as coronary heart disease, diabetes, stroke, etc., which has seriously endangered human health. With the economic development of human society, "obesity" has become a global problem [1,2], and the World Health Organization officially listed "obesity" as a disease in 2020. Our people are suffering from "obesity" for a long time [3], besides, the obesity rate of our population has surpassed the United States and become the world's first obese country. Although the economic take-off in the past few decades has brought unprecedented material enrichment to Chinese people, the concept of health has not yet become popular, which leads to the rampage of hypertension, diabetes, stroke and other chronic diseases caused by obesity caused by unhealthy diet and lack of exercise in China [4]. In 2019, the number of deaths caused by chronic diseases in China accounted for 88.5% of the total death population, and has now become a major burden on the medical system and the health of Chinese people in China. In addition, owing to the increasingly serious problem of adolescent "obesity" [5], many chronic diseases such as hypertension [6], diabetes, and gout begin to show a visible tendency to adolescent group, which has become a tough problem afflicting the healthy development of Chinese society. How to carry out scientific exploration for the "obese" population and find a scientific and effective weight loss method is the starting point and desire of this review. At present, people have fully recognized the benefits of exercise for health and chronic diseases, and many exercise methods, such as aerobic exercise, resistance exercise [7], resistance plus aerobic combined exercise [8], HIIT high-frequency intermittent exercise [9], and special exercise, have begun to be gradually accepted by the public, but there is a lack of an efficient and sustainable fat reduction exercise. Resistance exercise and dedicated exercise require professional venues and devices; although the combination of aerobic and resistance exercise can efficiently lose weight, it takes too much time; HIIT exercise is effective, but it requires a good training basis and cannot be introduced with a weak foundation; aerobic exercise was first proposed by Cooper and defined by the Oxford Dictionary in 1986 as an exercise method that meets the increase and maintenance of moderate oxygen intake through exercise and is beneficial to the respiratory and circulatory systems [10,11]. In decades of practical and scientific research exploration, it has been found that it can effectively improve body shape, fasting blood glucose, blood lipids and blood pressure and recognized as a safe and effective means of weight loss [12]. With the deepening of research, Davis et al., [13] proposed in 1989 that the thermogenic effect of premeal and post-prandial exercise were totally different, and once it was introduced, it set off a wave of research on the effect of post-fasting exercise on body composition. Since then, fasting aerobic has gradually entered the public career as an efficient and convenient way to reduce fat, and is widely used as a final "tool" to reduce stubborn fat among bodybuilders and enthusiasts [14,15]. In addition, fasting aerobics has been a hot research topic in foreign sports circles and sports medicine in recent years [16]. The research on its mechanism, molecules and phenotypes has become more and more in-depth and comprehensive. This review hopes to

synthesize domestic and foreign research, summarize the current situation of fasting aerobic research, make a specific description of its mechanism, and provide a theoretical reference for future research.

Fasting Aerobic Mechanism and the Current Status of Domestic and International Research

Fasting aerobic mechanism

The fasting state refers to the physical state of fasting for more than 8 hours, while fasting aerobic refers to endurance exercise in fasting state [17]. In fasting or hungry state, the body maintains a constant blood glucose level by isogenic glucose such as lysine and glycerol [18]. The process of conversion of non-sugar compounds, pyruvate, lactate, glycerol, amino acids, etc. into glucose is called gluconeogenesis. In a prolonged fasting state, the body with no exogenous energy intake and nearly complete glycogen depletion, will preferentially choose adipose tissue for catabolism to meet the needs of vital activities. A consensus at this time suggests that in the fasting state, the body's utilization of fat is excessive [18,19], so the caloric deficit caused by fasting aerobic can cause body fat reduction in the short term and even regulate autophagy to delay aging [20,21]. Aerobic exercise alone inevitably loses more skeletal muscle during the long fat reduction process and even leads to the appearance of a fat reduction plateau [22], and Maughan argued that muscle augmentation is not suitable in the fasting state. Fasting aerobic exercise as a unique way to combine fasting and exercise, breaking the traditional concept of exercise after meals, and greatly improves the efficiency of exercise fat reduction [23].

Status of foreign research: The hot wave of fasting aerobic research, in foreign countries, has continuous for decades, the research direction is all over the molecules, physiological mechanisms, appearance, etc., and it is precisely due to the benefits of fasting aerobic found in the past research that it has been very mature in foreign countries as a healthy and effective method for the prevention and treatment of hypertension, hyperlipidemia, coronary heart disease, diabetes and other chronic diseases.

In terms of the appearance of animal studies: Mc Sheer [24] and others used 12-month-old Wistar rats for experiments and found that the weight loss effect of the experimental subjects in the fasting state was significantly greater than that of the control group; The study by Mark E Smyers et al., [25] specifically selected rats with low fat oxidation capacity and high fat oxidation capacity. The results showed that although the weight loss was not statistically significant, the "fat loss" of rats in the fasting aerobic exercise group was significantly greater than that in the feeding group, which was also confirmed by the study by Antonio Real-Hohn [26]; In terms of human phenotypic and physiological mechanism research: Ferland [27] convened 10 patients with stage II diabetes to conduct a randomized trial and found that fasting aerobic exercise had a wonderful effect on energy metabolism after the experiment; Milena Barbon [28] discovered that it could significantly improve fat combustion efficiency compared with the placebo control group after ingesting 6 g taurine to 17 healthy men; SAMARMAR [29] convened 23 healthy volunteers and conducted the experiment and found that the mixed exercise fasting aerobic group had a perfect weight reduction effect than the other single exercise groups, of which the fasting exercise group had a better fat loss effect than the mixed exercise group. Fereshteh et al., [30] convened 83 patients with nonalcoholic fatty liver disease to perform fasting exercise intervention and found that it could effectively improve the

condition; Kristin K [31] found that the satiety developed by fasting aerobic exercise facilitated long-term weight loss in 61 healthy volunteers after an eight-week exercise intervention, which is instructive for people's long-term weight loss; it's interesting that Victoria J [32], after experiments with 12 people, found that gastrointestinal function, hunger and appetite regulating hormones were insensitive to low-intensity activities, which suggested that people needed to perform a certain degree of high-intensity stimulation. Jenna B. Gillen performed thigh muscle biopsy after fasting aerobic exercise intervention experiments in 16 volunteers and found that fasting training-induced increased mitochondrial capacity and increased maximum activity of citrate lyase and b-hydroxycoenzyme a dehydrogenase [33]; even in Alessio Nencioni [34] on metabolic molecules and clinical applications of fasting and cancer discovered fasting can alter cancer factors. Almost all of their studies found that fasting aerobic was effective in reducing blood glucose, blood pressure, blood lipids, and insulin resistance levels in "obese" people. In summary, fasting aerobic has enough advantages from the perspective of human physical status and physiological indicators to allow us to adhere to and use.

Domestic research status: Domestic research on fasting aerobics started late and is relatively small, but abundant results have still been achieved at the phenotypic, physiological and molecular levels. In terms of performance and physiological studies, researchers in China have conducted extensive studies on patients with metabolic syndrome, diabetes, and stroke, and have achieved good efficacy in terms of physiological parameters: Body Mass Index (BMI), Triacyl Glycerides (TG), Systolic Blood Pressure (SBP), Diastolic Blood Pressure (DBP), Fasting Plasma Glucose (Fasting blood-glucose, FPG), 2-h plasma glucose (2-h protein-cholesterol), Total Cholesterol (TC), Low-Density Lipoprotein- Cholesterol (LDL-C) and other parameters closely related to metabolic diseases of the body have achieved effective control and decrease; meanwhile, High-Density Lipoprotein-Cholesterol (HDL-C) has increased. Recent studies have shown that the consumption of triglycerides in one hour has only 50% of the fat-burning effect in the fasting aerobic state, which can have higher levels of blood glycerol and fatty acid release, stimulate higher metabolic levels [35], and have a positive feedback on the treatment of patients with nonalcoholic fatty liver disease and type 2 diabetes. In molecular research, fasting aerobic has not been done, but many aerobic levels have emerged that can provide guidance for fasting aerobic. Ji Lili found that vascular aortic molecules LOX-1, NF- κ Bp65 and caspase-3 protein expression increased; Wang Xiaoqiang, Li Xun and others found that exercise intervention miR-7/PDX-1/GLUT2-GCK with dietary plan can effectively improve islet function; Wang Tianyuan, Wang Xiaohui found the factors PPAR α , PPAR γ and PPARs after aerobic exercise intervention in diabetic rats for 4 weeks, 6 days a week, 60 min a day! / σ expression increased; Li Jun, Feng Lijie found that inflammatory factors decreased, Nrf2/ARE signaling was activated, and vascular oxidative stress was relieved after eight weeks of swimming exercise in diabetic rats; Zhang Qiang performed intervention experiments by dividing rats into ketogenic diet, ketogenic diet and aerobic exercise groups, ketogenic diet and high-frequency intermittent exercise groups and found that aerobic exercise may accelerate fat oxidation in liver tissue of diabetic mice by activating AMPK/ACC/CPT1A signaling pathway and slow down the process of fatty liver development, while high-intensity intermittent exercise did not rely on this regulatory pathway. Various domestic studies also lay a certain foundation for the future development of fasting aerobic. With the understanding of the public and researchers, fasting aerobic will certainly set off an upsurge in China in the future.

Benefits of Fasting Aerobic

Improving Insulin Sensitivity and Insulin Resistance

Insulin sensitivity: In recent years, a large number of domestic and international studies have shown that fasting aerobic exercise can improve insulin sensitivity, but glucose and insulin metabolism under fasting exercise is a very complex and interesting thing. In the fasting state, the body's glycogen is consumed overnight and blood glucose is at a low level, when glucagon, which promotes catabolism, is high and insulin is low. It can promote glycogenolysis and gluconeogenesis, significantly increase blood glucose; in addition, it can also activate lipase, promote the notice of lipolysis to strengthen fatty acid oxidation, so that ketone bodies increase.

For example, the application of different relative exercise intensities significantly affected blood glucose and insulin concentrations. In trained cyclists, elevated blood glucose and insulin concentrations (above anaerobic threshold at 86% of VO₂ peak) were found during high-intensity exercise in the fasted state (as opposed to exercise in the fed state). In contrast, during moderate-intensity endurance exercise (below the anaerobic threshold at 79% of peak oxygen), fasting was comparable to fed state blood glucose and insulin concentrations. Thus, changes in blood glucose levels during exercise depend, at least in part, on exercise intensity. It is well-known that high-intensity exercise is associated with greater epinephrine and growth hormone synthesis/release, which will stimulate hepatic glucose output. Blood glucose concentrations may increase during high-intensity exercise (above the anaerobic threshold) due to greater glucose production rather than glucose utilization (in skeletal muscle cells).

Lei Yu develops insulin resistance through an 8-week diet in rats. Aerobic exercise can inhibit or reduce the occurrence of insulin resistance in the body, acting through activation of IL-6 and AMPK, IL-6 may reduce blood glucose by activating GLUT4 mRNA; Guo Yin conducted a 4-week exercise intervention experiment in 126 obese adolescents at Shanghai Peak Training Camp and found that aerobic exercise can effectively improve the body morphology of obese children and adolescents, improve lipid metabolism disorders and insulin resistance, and prevent metabolic syndrome; Gui Shukang performed an 8-week aerobic exercise intervention in mice with treadmill and found that aerobic exercise can reduce the concentration of insulin and IGF-1 in serum and significantly improve insulin sensitivity in insulin-resistant mice; Sun Min found that insulin resistance is related to Extracellular Matrix Protein (EMC); Wang Zhengqing and others performed a 12-week aerobic exercise intervention in diabetic patients and found that 12-week aerobic exercise can effectively improve the metabolic abnormalities such as hyperglycemia, low insulin sensitivity, and insulin resistance in IGR population, regulate the secretory function of islet β cells, and reduce the risk factors of cardiovascular disease. It normalized blood glucose in 62.4% of patients with Impaired Glucose Regulation (IGR). JA [36] conducted a study of 154 overweight subjects and found that 170 min of exercise per week better improved insulin sensitivity than 120 min of exercise, and there was a relationship between insulin sensitivity and sedentary.

Insulin resistance: Insulin can promote the uptake and utilization of glucose by tissues and cells, accelerate the synthesis of glucose into glycogen, store in liver and muscle, inhibit gluconeogenesis, promote the conversion of glucose into fatty acids, store in adipose tissue, resulting in decreased blood glucose levels. In insulin deficiency, the blood glucose concentration increases, and once the renal glucose

threshold is exceeded, sugar will appear in the urine, causing diabetes. Insulin resistance, on the other hand, refers to a decrease in the ability of insulin to maintain normoglycemia, which produces biological effects below normal levels, or a decrease in tissue responsiveness to insulin. It is mainly manifested in the reduced utilization of glucose uptake by peripheral tissues, especially adipose tissue and skeletal muscle, as well as the effect of reducing hepatic glucose output [37]. At present, obesity is considered to be the most important cause of insulin resistance, especially central obesity, which is mainly related to long-term insufficient exercise and excessive dietary energy intake. Patients with type 2 diabetes are diagnosed with obesity and insulin resistance is a more important risk factor leading to coronary heart disease or stroke. A large number of studies have shown that aerobic exercise can reduce insulin resistance in obese people and people with metabolic diseases, including college students, elderly patients with IGF, patients with IGF, obese patients, patients with early diabetes and elderly patients with diabetes, all of which have achieved good improvement results; Karen Van [38] found that systemic glucose tolerance and insulin sensitivity were significantly improved during fasting aerobic exercise intervention experiments on clerkships.

Improvement of diabetes and metabolic diseases

Type 2 diabetes is a chronic metabolic non-communicable disease characterized by chronic hyperglycemia and insulin resistance caused by various factors such as genes, environment and lifestyle [39]. Although insulin resistance and β -cell failure are considered to be the main causes of the development of type 2 diabetes, a large body of evidence suggests that diabetes is a systemic inflammatory metabolic disease. At present, diabetes is still incurable, so it is also known as chronic cancer.

Diabetes mellitus is currently the most serious chronic metabolic disease in China. It is reported that as of 2013, diabetic patients in China have accounted for 11.6% of the total population. Cardiovascular disease and insulin resistance caused by obesity caused by lack of exercise and unhealthy diet are the main causes of diabetes mellitus. Active and effective exercise therapy is currently being sought worldwide to improve diabetes and related metabolic diseases. Kristin I. Stanford [40] found that maternal exercise is especially important compared with the parental exercise after conducting an exercise intervention experiment in a diabetic rat model, and maternal exercise can significantly improve the ability of offspring to metabolically healthy, and the good glucose tolerance produced by this exercise intervention can be transmitted from mother to offspring, thereby counteracting obesity and diabetes; Michael [41] concluded that aerobic exercise can effectively improve diabetic conditions after META analysis of previous studies; Alissa A [42] conducted a 12-week randomized aerobic experiment in thirty obese patients and found that fasting blood glucose increased after glucose sodium co-transporter 2 (SGLT2) inhibition and insulin sensitivity increased after (SGLT2) inhibition after endurance exercise training, suggesting that Glucose Sodium co-Transporter 2 (SGLT2) induced an increase in insulin sensitivity.

Aerobic capacity

Endurance exercise training in the fasted state significantly improves basal muscle fat transport and oxidative capacity, as evidenced by a significant increase in basal muscle fatty acid translocase CD36 (FAT/CD36) protein and Fatty Acid Binding Protein (FABP) content, which is superior to exercise training in the fed state in this regard;

Alexandra [43] conducted a META analysis of more than 10,400 participants and finally found that the fat oxidative capacity was enhanced in the fasting aerobic group; recent studies also showed that fasting aerobic can improve gastrointestinal sensitivity, obese people due to long-term massive excess diet lead to reduced gastrointestinal sensitivity to food, while small-intensity exercise does not; high-intensity exercise can increase tissue oxygen supply, while leading to increased reactive oxygen species production [44].

Weight loss

Obesity is one of the main culprits leading to cardiovascular disease. According to the 2020 Report on Nutrition and Chronic Diseases of Chinese Residents, overweight and obese people account for 50% in China. People have been making various attempts to reduce the obesity rate, and since people proposed that exercise is the slogan of good medicine, various exercise methods have emerged, but most of them have a threshold and cannot adhere to it for a long time, and since then fasting aerobic has come into front.

Past theories and practices have demonstrated that fasting aerobic brings greater benefits to weight loss than ordinary aerobic training. This is also the same as the results of the Michelle RJospe experiment, in which fasting aerobic decreased more significantly than fed aerobic weight through a 12-month participant self-supervision experiment, and because this experiment was performed by patient self-supervision it also reflected its sustainability from the side.

Past theories and practices have demonstrated that fasting aerobic brings greater benefits to weight loss than ordinary aerobic training. This is also the same as the results of the Michelle RJospe experiment, in which fasting aerobic decreased more significantly than fed aerobic weight through a 12-month participant self-supervision experiment, and because this experiment was performed by patient self-supervision it also reflected its sustainability from the side.

The Su-JongKim-Dorner [45] study found that optimistic psychological attitudes had a positive effect on insulin resistance through a study of 108 participants in the United States, and I suspect that mood pleasure may produce certain factors that activate insulin function, such as lowering cortisol; there is a very interesting study recently, through a group experiment of 873 students with obesity and health, and finally found that high levels of cardiopulmonary health function can counteract the harm caused by hyperlipidemia [46], and if you really cannot reduce fat, it may be a good choice to improve cardiopulmonary capacity.

Autonomic function

Heart Rate Variability (HRV), which reflects the function of cardiac autonomic nervous system, can reflect the control and regulation of heart by nervous system and reflect the degree of variation between successive sinus beats and cardiac cycle in obese people. This feature is the result of the common regulation of human neurohumoral and cardiovascular systems. Not only has the odds of cardiovascular disease, hypertension, type II diabetes, in excessive weight predisposed obese individuals, will increase, but also will face a higher risk of death compare with normal. The disorders of autonomic nervous system in obese people have become the focus of attention in the health field, and recent studies have been found that obese people gain 10% of body weight and have reduced parasympathetic regulation; while obese or overweight people lose 10% of body weight and have increased parasympathetic nerves, and the increase in body

weight leads to inhibition of parasympathetic nerves, enhancement of sympathetic nerves, and disruption of the balance of cardiac rhythm, which is likely to be caused by disturbances in their own ANS (autonomic nervous system) function in obese people and is an important cause of cardiovascular disease. Guan Yuming et al., studied heart rate variability and vagus nerve in adolescent obese people, and the results showed that in people with reduced heart rate variability, the corresponding vagus nerve function was also reduced; Gao Arning observed the relationship between Body Mass Index(BMI) and heart rate variability in 70 obese children and found that BMI was positively correlated with Triglyceride (TG) and High-Density Lipoprotein Cholesterol (LDL-C) in obese children; BMI was negatively correlated with SDNN, SDANN, and RMSSD, and TG was negatively correlated with SDNN, SDANN, and RMSSD. The results showed that obese children may cause autonomic nerve damage due to high TG; Zhang Jingyang et al., found that the content of metabolic indicators in obese children was significantly correlated with heart rate variability indicators SDNN, SDANN, RMSSD, body cytokines, body adipocytokines and other indicators after observing 260 obese children. In summary, heart rate variability in obese children is closely related to the body's lipid metabolism, and excessive obesity can lead to abnormal cardiac function. Recent studies have shown that metabolism affects a variety of pathways in the human body, MarkP [47] proposed a hypothesis on the basis of previous experiments, that light fasting can improve nerve reactivity and brain health, improve brain resistance to injury and disease; simultaneously, he also proposed a conjecture which based on his long-term research on diet and brain nerves that excessive intake will damage some metabolism-related pathways which will affect people's cognitive and neurological function [48].

Exercise is an auxiliary tool in the treatment of T2D, abundant time-tested evidence can prove aerobic exercise can have unique positive effects on autonomic function while improving glycemic control [49,50] and insulin resistance [50,51]. Long-term moderate-intensity aerobic exercise can improve cardiac autonomic regulation and baroreflex function in T2D [52,53] is a consensus in the field of research on the effects of exercise on autonomic function in the fasting state of T2D. The effects of aerobic exercise and resistance training on autonomic function during fasting and glucose load in patients with type 2 diabetes were studied in China, and the results showed that aerobic capacity, muscle mass, glycosylated hemoglobin, fasting blood glucose, HOMA2-0, body weight, BMI, HDL-C, LDL-C and Framingham risk score and Heart Rate Variability (HRV) were improved [54-56]. In summary, exercise can improve cognition and autonomic function.

Other benefits

Historic breakthrough has been discovered not long ago. Though we still can't figure out the mechanism, that can't obstruct the enthusiasm about the detection of the combination of rapamycin and fasting aerobic exercise have a magical effect in against anti-aging [20]; Jenna B [33] found by biopsy that fasting training-induced increased mitochondrial capacity and increased maximum activity of citrate lyase and b-hydroxycoenzyme a dehydrogenase.

Problems and Prospects

The current domestic research focuses on the aerobic exercise level and has not been advanced to the fasting aerobic stage. Fasting aerobic as a method for the treatment of cardiovascular diseases and obesity has been very popular abroad. There are many advantages of

fasting aerobic itself, such as the previously mentioned increase of high density lipoprotein, insulin sensitivity, gastrointestinal sensitivity, insulin resistance, low-density lipoprotein, blood pressure, blood lipids, fasting blood glucose, body weight, body fat rate, waist circumference and so on. In addition to its many advantages of improving physical conditions, the cost performance of fasting aerobic is also very prominent. In terms of economy, compared with resistance and special sports requiring complete sites and professional devices, fasting aerobic requires only a simple treadmill or even a home bicycle to perform fasting aerobic; half an hour after the start of the morning will not occupy too much time.

There are few existing fasting aerobic studies in China, and its mechanistic concept is still vague and has great potential, and it is sincere to hope that domestic researchers can carry out more meaningful research in the future.

Tip: The fasting aerobic benefits addressed throughout this article are presented in figure 1




Figure 1: The fasting aerobic benefits.

References

- Wang Y, Wang L, Qu W (2017) New national data show alarming increase in obesity and noncommunicable chronic diseases in China. *European journal of clinical nutrition* 71: 149-150.
- Wang Y, Lobstein T (2006) Worldwide trends in childhood overweight and obesity. *International journal of pediatric obesity. Int J Pediatr Obes* 1: 11-25.
- Wang Y, Mi J, Shan XY, Wang QJ, Ge KY (2007) Is China facing an obesity epidemic and the consequences? The trends in obesity and chronic disease in China. *Int J Obes (Lond)* 31: 177-188.
- Estes C, Anstee QM, Lobstein M, Bantel H, Bellentani S, et al. (2018) Modeling NAFLD disease burden in China, France, Germany, Italy, Japan, Spain, United Kingdom, and United States for the period 2016-2030. *Journal of hepatology* 69: 896-904.
- Steinbeck KS, Lister NB, Gow ML, Baur LA (2018) Treatment of adolescent obesity. *Nature reviews Endocrinology* 14: 331-344.
- Seeman T, Hamdani G, Mitsnefes M (2019) Hypertensive crisis in children and adolescents. *Pediatric nephrology (Berlin, Germany)* 34: 2523-2537.
- Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, et al. (2018) The Physical Activity Guidelines for Americans. *Jama* 320: 2020-2028.
- Ho SS, Dhaliwal SS, Hills AP, Pal S (2012) The effect of 12 weeks of aerobic, resistance or combination exercise training on cardiovascular risk factors in the overweight and obese in a randomized trial. *BMC Public Health* 12: 704.
- Batacan RB, Duncan MJ, Dalbo VJ, Tucker PS, Fenning AS (2017) Effects of high-intensity interval training on cardiometabolic health: a systematic review and meta-analysis of intervention studies. *British journal of sports medicine* 51: 494-503.
- Fisher JP, Young CN, Fadel PJ (2015) Autonomic adjustments to exercise in humans. *Comprehensive Physiology* 5: 475-512.
- McKenzie DC (2012) Respiratory physiology: adaptations to high-level exercise. *British journal of sports medicine* 46: 381-384.
- Gollisch KS, Brandauer J, Jessen N, Toyoda T, Nayer A, et al. (2009) Effects of exercise training on subcutaneous and visceral adipose tissue in normal- and high-fat diet-fed rats. *Am J Physiol Endocrinol Metab* 297: 495-504.
- Davis JM, Sadri S, Sargent RG, Ward D (1989) Weight control and caloric expenditure: thermogenic effects of pre-prandial and post-prandial exercise. *Addict Behav* 14: 347-351.
- Robinson SL, Mansell AL, Gillibrand G, Ryan AS, Bannock L (2015) A nutrition and conditioning intervention for natural bodybuilding contest preparation: case study. *Journal of the International Society of Sports Nutrition* 12: 20.
- Hartgens F, Rietjens G, Keizer HA, Kuipers H, Wolffenbuttel BH (2004) Effects of androgenic-anabolic steroids on apolipoproteins and lipoprotein (a). *British journal of sports medicine* 38: 253-259.
- Hartgens F, Rietjens G, Keizer HA, Kuipers H, Wolffenbuttel BH (2004) Effects of androgenic-anabolic steroids on apolipoproteins and lipoprotein (a). *British journal of sports medicine* 38: 253-259.
- Meng H, Zhu L, Varkaneh HK, Heitor OS, Tinsley GM, et al. (2020) Effects of intermittent fasting and energy-restricted diets on lipid profile: A systematic review and meta-analysis. *Nutrition* 77: 110801.
- Secor SM, Carey HV (2016) Integrative Physiology of Fasting. *Comprehensive Physiology* 6: 773-825.
- Tinsley GM, La Bounty PM (2015) Effects of intermittent fasting on body composition and clinical health markers in humans. *Nutrition reviews* 73: 661-674.
- Blagosklonny MV (2019) Fasting and rapamycin: diabetes versus benevolent glucose intolerance. *Cell Death Dis* 10: 607.
- Kristensen CM, Olsen MA, Jessen H, Brandt N, Meldgaard JN, et al. (2018) PGC-1 α in exercise and fasting-induced regulation of hepatic UPR in mice. *Pflugers Archiv: European journal of physiology* 470: 1431-1447.
- Maughan RJ (2010) Fasting and sport: an introduction. *British journal of sports medicine* 44: 473-475.
- Li L, Wang Z, Zuo Z (2013) Chronic intermittent fasting improves cognitive functions and brain structures in mice. *PLoS One* 8: 66069.
- Jospe MR, Roy M, Brown RC, Haszard JJ, Jones KM, et al. (2020) Intermittent fasting, Paleolithic, or Mediterranean diets in the real world: exploratory secondary analyses of a weight-loss trial that included choice of diet and exercise. *The American journal of clinical nutrition* 111: 503-514.
- Smyers ME, Koch LG, Britton SL, Wagner JG, Novak CM (2021) Enhanced weight and fat loss from long-term intermittent fasting in obesity-prone, low-fitness rats. *Physiol Behav* 230: 113280.
- Hohn AR, Navegantes C, Ramos K, Filho DR, Cahué F, et al. (2018) The synergism of high-intensity intermittent exercise and every-other-day intermittent fasting regimen on energy metabolism adaptations includes hexokinase activity and mitochondrial efficiency. *PLoS One* 13: 0202784.
- Ferland A, Broderick TL, Nadeau A, Simard S, Martin J, et al. (2007) Impact of fasting and postprandial state on plasma carnitine concentrations during aerobic exercise in type 2 diabetes. *Acta Diabetol* 44: 114-120.

28. Carvalho MB, Brandao CFC, Fassini PG, Bianco TM, Batitucci G, et al. (2020) Taurine Supplementation Increases Post-Exercise Lipid Oxidation at Moderate Intensity in Fasted Healthy Males. *Nutrients* 12: 1540.
29. Chacaroun S, Borowik A, Escamilla YGIV, Doutreleau S, Wuyam B, et al. (2020) Hypoxic Exercise Training to Improve Exercise Capacity in Obese Individuals. *Med Sci Sports Exerc* 52: 1641-1649.
30. Aliasghari F, Izadi A, Gargari BP, Ebrahimi S (2017) The Effects of Ramadan Fasting on Body Composition, Blood Pressure, Glucose Metabolism, and Markers of Inflammation in NAFLD Patients: An Observational Trial. *J Am Coll Nutr* 36: 640-645.
31. Hoddy KK, Gibbons C, Kroeger CM, Trepanowski JF, Barnosky A, et al. (2016) Changes in hunger and fullness in relation to gut peptides before and after 8 weeks of alternate day fasting. *Clinical nutrition* 35: 1380-1385.
32. McIver VJ, Mattin L, Evans GH, Yau AMW (2019) The effect of brisk walking in the fasted versus fed state on metabolic responses, gastrointestinal function, and appetite in healthy men. *Int J Obes* 43: 1691-1700.
33. Gillen JB, Percival ME, Ludzki A, Tarnopolsky MA, Gibala MJ (2013) Interval training in the fed or fasted state improves body composition and muscle oxidative capacity in overweight women. *Obesity* 21: 2249-2255.
34. Nencioni A, Caffa I, Cortellino S, Longo VD (2018) Fasting and cancer: molecular mechanisms and clinical application. *Nature reviews Cancer* 18: 707-719.
35. Enevoldsen LH, Simonsen L, Macdonald IA, Bülow J (2004) The combined effects of exercise and food intake on adipose tissue and splanchnic metabolism. *J Physiol* 561: 871-882.
36. Houmard JA, Tanner CJ, Slentz CA, Duscha BD, McCartney JS, et al. (1985) Effect of the volume and intensity of exercise training on insulin sensitivity. *Journal of applied physiology* 96: 101-106.
37. Cheng S, Ge J, Zhao C, Le S, Yang Y, et al. (2017) Effect of aerobic exercise and diet on liver fat in pre-diabetic patients with non-alcoholic-fatty-liver-disease: A randomized controlled trial. *Scientific reports* 7: 15952.
38. Proeyen KV, Szlufcik K, Nielens H, Pelgrim K, Deldicque L, et al. (2010) Training in the fasted state improves glucose tolerance during fat-rich diet. *J Physiol* 588: 4289-4302.
39. Yang SH, Dou KF, Song WJ (2010) Prevalence of diabetes among men and women in China. *N Engl J Med* 362: 2425-2426.
40. Stanford KI, Takahashi H, So K, Wagner ABA, Prince NB, et al. (2017) Maternal Exercise Improves Glucose Tolerance in Female Offspring. *Diabetes* 66: 2124-2136.
41. Fedewa MV, Gist NH, Evans EM, Dishman RK (2014) Exercise and insulin resistance in youth: a meta-analysis. *Pediatrics* 133: 163-174.
42. Newman AA, Grimm NC, Wilburn JR, Schoenberg HM, Tripathi SRJ, et al. (2019) Influence of Sodium Glucose Cotransporter 2 Inhibition on Physiological Adaptation to Endurance Exercise Training. *The Journal of clinical endocrinology and metabolism* 104: 1953-1966.
43. Vieira AF, Costa RR, Macedo RC, Coconcelli L, Kruegel LF (2016) Effects of aerobic exercise performed in fasted v. fed state on fat and carbohydrate metabolism in adults: a systematic review and meta-analysis. *The British journal of nutrition* 116: 1153-1164.
44. Wiecek M, Maciejczyk M, Szymura J, Szygula Z (2015) Changes in oxidative stress and acid-base balance in men and women following maximal-intensity physical exercise. *Physiol Res* 64: 93-102.
45. Dornier SJK, McKenzie COS, Poth M, Deuster PA (2009) Psychological and physiological correlates of insulin resistance at fasting and in response to a meal in African Americans and Whites. *Ethn Dis* 19: 104-110.
46. Ruiz JR, Rizzo NS, Ortega FB, Loit HM, Veidebaum T, et al. (2007) Markers of insulin resistance are associated with fatness and fitness in school-aged children: the European Youth Heart Study. *Diabetologia* 50: 1401-1408.
47. Mattson MP, Moehl K, Ghena N, Schmaedick M, Cheng A (2018) Intermittent metabolic switching, neuroplasticity and brain health. *Nature reviews Neuroscience* 19: 63-80.
48. Mattson MP (2019) An Evolutionary Perspective on Why Food Overconsumption Impairs Cognition. *Trends Cogn Sci* 23: 200-212.
49. Reddy R, Wittenberg A, Castle JR, Youssef J, Stone KW, et al. (2019) Effect of Aerobic and Resistance Exercise on Glycemic Control in Adults With Type 1 Diabetes. *Canadian journal of diabetes* 43: 406-414.
50. Yang Z, Scott CA, Mao C, Tang J, Farmer AJ (2014) Resistance exercise versus aerobic exercise for type 2 diabetes: a systematic review and meta-analysis. *Sports medicine* 44: 487-499.
51. Goulopoulou S, Baynard T, Franklin RM, Fernhall B, Carhart R, et al. (2010) Exercise training improves cardiovascular autonomic modulation in response to glucose ingestion in obese adults with and without type 2 diabetes mellitus. *Metabolism: clinical and experimental* 59: 901-910.
52. Goit RK, Paudel BH, Khadka R, Roy RK, Shrestwa MK (2014) Mild-to-moderate intensity exercise improves cardiac autonomic drive in type 2 diabetes. *Journal of diabetes investigation* 5: 722-727.
53. Goit RK, Pant BN, Shrestwa MK (2018) Moderate intensity exercise improves heart rate variability in obese adults with type 2 diabetes. *Indian heart journal* 70: 486-491.
54. Masroor S, Bhati P, Verma S, Khan M, Hussain ME (2018) Heart Rate Variability following Combined Aerobic and Resistance Training in Sedentary Hypertensive Women: A Randomised Control Trial. *Indian heart journal* 70: 28-35.
55. Kelley GA, Kelley KS, Roberts S, Haskell W (2011) Efficacy of aerobic exercise and a prudent diet for improving selected lipids and lipoproteins in adults: a meta-analysis of randomized controlled trials. *BMC medicine* 9: 74.
56. Ratajczak M, Skrypnik D, Krutki P, Karolkiewicz J (2020) Effects of an Indoor Cycling Program on Cardiometabolic Factors in Women with Obesity vs. Normal Body Weight. *International journal of environmental research and public health* 17: 8718.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldoopenaccess.us/submit-manuscript>